

## Ca<sup>2+</sup> Signaling Domains Responsible For Cardiac Hypertrophy and Arrhythmias

Steven R. Houser

Ca<sup>2+</sup> activates and regulates multiple processes in every cell type. In the mammalian heart, cyclic fluctuations in cytosolic [Ca<sup>2+</sup>] induce and regulate the strength of cardiac contraction (termed “contractile” [Ca<sup>2+</sup>]). In addition, changes in Ca<sup>2+</sup> appear to be centrally involved in normal and pathological signaling (termed “signaling” [Ca<sup>2+</sup>]) that regulates myocyte growth, hypertrophy, apoptosis, and necrosis.<sup>1</sup> Whether or not contractile and signaling [Ca<sup>2+</sup>] are derived from common or distinct sources and are constrained to unique cellular microdomains is not established.<sup>2</sup> What is clear is that cardiovascular diseases including hypertension and myocardial infarction are associated with alterations in contractile and possibly signaling [Ca<sup>2+</sup>] that are centrally involved in pathological cardiac hypertrophy, heart failure progression,<sup>1</sup> and lethal cardiac arrhythmias.<sup>3</sup> Defining the sources of signaling Ca<sup>2+</sup> involved in the induction of pathological hypertrophy and the bases of dysregulated contractile [Ca<sup>2+</sup>] in cardiovascular disease should identify novel ways to treat heart disease.

In this issue of *Circulation Research*, 2 independent reports address fundamental aspects of alterations in signaling and contractile [Ca<sup>2+</sup>]. Chiang et al<sup>4</sup> have studied the idea that Ca<sup>2+</sup> influx through voltage operated  $\alpha$ 1H (Ca<sub>v</sub>3.2) T-type Ca<sup>2+</sup> channels (TTCCs) is the source of the signaling [Ca<sup>2+</sup>] that activates the calcineurin (Cn)-NFAT (nuclear factor of activated T cells) signaling cascade and induces pathological cardiac hypertrophy in pressure overload. In a separate report, Terentyev et al<sup>5</sup> explore the idea that microRNA (miR)-1, a muscle-specific microRNA that increases in abundance in cardiac disease,<sup>6</sup> causes dysregulated contractile [Ca<sup>2+</sup>] and induces single cell arrhythmias. These 2 reports are provocative and, if independently confirmed, will have identified novel mechanisms for abnormalities in the signaling and contractile [Ca<sup>2+</sup>] that cause hypertrophy and sudden death.

Almost 20 years ago, we<sup>7</sup> and others<sup>8</sup> showed that TTCCs are reexpressed in adult ventricular myocytes after pressure overload. TTCCs are expressed in fetal/neonatal heart but are not normally found in the adult ventricular myocyte. We speculated that the Ca<sup>2+</sup> influx through these channels was involved in the induced cardiac hypertrophy.<sup>7</sup> The report by Chiang et al<sup>4</sup> explores this idea in TTCC knockout (KO)

mouse models. There are 3 TTCC genes, and 2 ( $\alpha$ 1G [Ca<sub>v</sub>3.1] and  $\alpha$ 1H [Ca<sub>v</sub>3.2]) are found in the heart.<sup>9</sup> Ca<sub>v</sub>3.1<sup>10</sup> and 3.2 KO<sup>11</sup> animals, each of which is viable with modest basal phenotypes,<sup>10,11</sup> were used. The authors make the provocative observation that thoracic aortic constriction (TAC) induces cardiac hypertrophy in the Ca<sub>v</sub>3.1 KO and control animals, but not in Ca<sub>v</sub>3.2 KO. Ca<sub>v</sub>3.2 KO animals had similar degrees of pressure overload after TAC, documenting a similar degree of stress. The inability of TAC to induce hypertrophy in Ca<sub>v</sub>3.2 KO appeared to be attributable to the fact that Cn-mediated nuclear NFAT translocation, which is known to induce pathological hypertrophy,<sup>12</sup> was not activated in these animals. Surprisingly, the fetal gene program activated with pathological hypertrophy was induced by TAC in Ca<sub>v</sub>3.2 KO without left ventricular hypertrophy.

These are provocative results that, if confirmed, will change thinking in the field. These results suggest that most if not all of NFAT mediated pathological hypertrophy is induced by a very small influx of Ca<sup>2+</sup> through reexpressed  $\alpha$ 1H TTCCs. These new findings also suggest that Cn-NFAT signaling is not influenced by changes in the amplitude and duration of the systolic [Ca<sup>2+</sup>] transient (contractile [Ca<sup>2+</sup>]). Contractility in Ca<sub>v</sub>3.2 KO mice after TAC must be greater than in controls which develop left ventricular hypertrophy, because Ca<sub>v</sub>3.2 KO hearts are generating high pressures with less cardiac mass. Therefore, the systolic Ca<sup>2+</sup> must be greater in Ca<sub>v</sub>3.2 KO TAC myocytes than in control TAC hearts, yet there was no activation of Cn-NFAT signaling. These results are different from those that have linked the activation of Cn-NFAT signaling with increases in either the rate or amplitude of the cytoplasmic (contractile) [Ca<sup>2+</sup>] transient in skeletal<sup>13</sup> and cardiac muscle.<sup>2,14</sup>

The report by Chiang et al<sup>4</sup> also suggests that Ca<sup>2+</sup> activated Cn-NFAT signaling does not play a role in the activation of the fetal gene program after TAC. Their studies show no activation of Cn-NFAT signaling in Ca<sub>v</sub>3.2 KO animals after TAC, but the fetal gene program was induced. In fact, the induction was greater than in controls after TAC. These results suggest that NFAT nuclear translocation has no role in the activation of these well studied fetal genes. Such results are in stark contrast to studies that have shown equally convincing data documenting that block of NFAT nuclear translocation eliminates agonist and pressure overload induced hypertrophy and the activation of the fetal gene program.<sup>12</sup> Because these data sets seem mutually exclusive this topic clearly needs additional study.


The provocative study by Chiang et al<sup>4</sup> suggests that pressure overload causes hypertrophy by inducing the expression of Ca<sub>v</sub>3.2 TTCCs. A very small Ca<sup>2+</sup> influx through these channels would need to enter a specialized subsar-

The opinions expressed in this editorial are not necessarily those of the editors or of the American Heart Association.

From the Temple University School of Medicine, Philadelphia, Pa.  
Correspondence to Steven R. Houser, PhD, FAHA, Laura H. Carnell Professor of Physiology, Temple University School of Medicine, 3400 N Broad St, Philadelphia, PA 19140. E-mail steven.houser@temple.edu  
(*Circ Res.* 2009;104:413-415.)

© 2009 American Heart Association, Inc.

*Circulation Research* is available at <http://circres.ahajournals.org>  
DOI: 10.1161/CIRCRESAHA.109.193821



**Figure.** Illustration depicting the novel signaling pathways in cardiac myocytes reported in this issue of *Circulation Research*.  $Ca^{2+}$  primarily enters cardiac myocytes via LTCCs and induces a larger amount of  $Ca^{2+}$  release from the SR, by activation of  $Ca^{2+}$  release channels (RyR). The report by Chiang et al<sup>4</sup> suggests that the  $Ca^{2+}$  required for induction of cardiac hypertrophy is not derived from the  $Ca^{2+}$  that induces cardiac contraction. Their data suggest that hypertrophic  $Ca^{2+}$  is very small and enters the cell exclusively via TTCCs that are expressed in response to cardiac stress. In this study, changes in the global  $Ca^{2+}$  transient did not activate the Cn-NFAT signaling cascade that induces turns on hypertrophic genes. This implies that a very tiny  $Ca^{2+}$  influx through TTCCs activates Cn (a phosphatase) within a portion of the cell that is protected from large changes in the global  $Ca^{2+}$

transient. The reported data are such that Cn would need to be constrained to this microdomain and dephosphorylate NFAT locally before it can translocate to the nucleus. The report by Terentyev et al<sup>5</sup> suggests that miR-1 influences a phosphatase localized near the junctional SR and this specifically alters CaMKII-mediated phosphorylation of RyR. It appears that modest phosphorylation of RyR at either S2808 (a PKA site) or S2814 (a CaMKII site) produces an increase in SR  $Ca^{2+}$  release (from green to blue in the illustration) without arrhythmias. This study also suggests that hyperphosphorylation of RyR at both S2808 and S2814 is needed to induce the abnormal spontaneous and evoked SR  $Ca^{2+}$  release associated with arrhythmias (black tracing in the cartoon). These new reports suggest that  $Ca^{2+}$  signaling in cardiac myocytes can be controlled locally by constraining downstream mediators to specific microdomains and locally regulating the phosphorylation state of  $Ca^{2+}$ -handling proteins.

colemmal signaling domain that is not influenced by large changes in contractile  $[Ca^{2+}]_i$ , where it exclusively activates Cn-NFAT signaling cascades. These new results suggest that pathological hypertrophy is induced through a highly specialized signaling  $[Ca^{2+}]_i$  microdomain that protects Cn-NFAT signaling from changes in contractile  $Ca^{2+}$  and causes pathological hypertrophy without activation of the fetal gene program. These results also exclude a role for TRPC, IP3R, and L-type  $Ca^{2+}$  channels (LTCCs) as a source of  $Ca^{2+}$  regulating cardiac hypertrophy and Cn-NFAT activity, in contrast to numerous reports.<sup>2</sup>

The second  $Ca^{2+}$  centric report in this issue of *Circulation Research*, by Terentyev et al,<sup>5</sup> identified a novel role for miR-1 in the regulation of contractile  $Ca^{2+}$ . Increasing miR-1 in cardiac myocytes caused alterations in the properties of the systolic  $Ca^{2+}$  transient, sarcoplasmic reticulum (SR)  $Ca^{2+}$  loading, and spontaneous and evoked SR  $Ca^{2+}$  release. When myocytes were exposed to catecholamines (isoproterenol [ISO]), only miR-1 myocytes demonstrated arrhythmogenic  $Ca^{2+}$  release. These results suggest that when miR-1 is increased in the diseased heart, catecholamine stress could induce life-threatening arrhythmias.

A novel aspect of this study was that the authors identified that miR-1 targets a regulatory subunit (B56 $\alpha$ ) of protein phosphatase (PP)2A, leading to reduced PP2A activity and increased phosphorylation of PP2A target proteins. Interestingly, only the phosphorylation state of specific  $Ca^{2+}$ /calmodulin kinase (CaMK)II phosphorylation sites were in-

creased in miR-1 myocytes, and inhibition of CaMKII with KN93 reversed dysregulated  $Ca^{2+}$  handling. These results add to the growing body of work linking persistent activation of CaMKII to cardiac dysfunction.<sup>15</sup> The authors concluded that hyperphosphorylation of the SR  $Ca^{2+}$  release channel (ryanodine receptor [RyR]) at a known CaMKII site (S2814) alters RyR function and is responsible for arrhythmogenic SR  $Ca^{2+}$  release in the presence of catecholamines (Figure).

The idea that either protein kinase (PK)A or CaMKII mediated phosphorylation of RyR can induce SR  $Ca^{2+}$  leak and cardiac arrhythmias is a contentious topic<sup>16</sup> and, in my view, this new study does not resolve critical issues. Although the authors have shown dysregulated  $Ca^{2+}$  in miR-1 myocytes as well as alterations in RyR phosphorylation at RyR S2814, a cause and effect relationship between these 2 miR-1 effects was not proven.

The effects of miR-1 on myocyte  $Ca^{2+}$  handling were complex and varied with conditions. In quiescent miR-1 myocytes, RyR S2814 and LTCC phosphorylation were increased, spark activity (an index of RyR activity) was enhanced and SR  $Ca^{2+}$  loading was reduced. The authors conclude that RyR phosphorylation at S2814 enhances RyR opening to cause diastolic SR  $Ca^{2+}$  "leak," which reduces SR  $Ca^{2+}$  loading. In voltage-clamped myocytes L-type  $Ca^{2+}$  current and  $Ca^{2+}$  transient amplitude were increased at positive potentials, suggestive of increased excitation-contraction coupling gain. ISO failed to further increase L-type  $Ca^{2+}$  current and the amplitude of the  $Ca^{2+}$  transient in

miR-1 myocytes did not increase and was smaller than in controls. miR-1 myocytes field stimulated at 1 Hz had systolic Ca<sup>2+</sup> transients that were much larger than in controls and SR Ca<sup>2+</sup> loading was now normalized. Why increases in RyR phosphorylation at S2814 would unload the SR in voltage-clamped and quiescent myocytes and maintain SR Ca<sup>2+</sup> load when these myocytes are paced is unclear, and suggest other unmeasured factors contribute to miR-1 effects on myocyte contractile Ca<sup>2+</sup>. Like most new findings, there are many issues to be resolved in future studies.

ISO induced arrhythmogenic Ca<sup>2+</sup> release only in miR-1 myocytes. The authors conclude that this resulted from hyperphosphorylation of RyR at S2814. To me, this conclusion is not fully justified. RyR S2814 phosphorylation is increased in miR-1 myocytes under control conditions and arrhythmogenic Ca<sup>2+</sup> release is not present. This suggests that CaMKII-mediated phosphorylation of RyR at S2814 is not sufficient to induce single cell arrhythmias. Adding ISO to miR-1 cells induced arrhythmias but did not cause further increases in RyR S2814 phosphorylation or the phosphorylation of the LTCCs, so it is unclear how phosphorylation at S2814 alone can be responsible for the induction of arrhythmias. ISO increased PKA-mediated phosphorylation of RyR at S2808, suggesting that hyperphosphorylation of RyR at this site could be the factor that precipitates arrhythmogenic Ca<sup>2+</sup> signaling. However, after inhibition of CaMKII with Kn93, RyR S2814 phosphorylation was reduced, RyR S2808 remained hyperphosphorylated, and arrhythmogenic Ca<sup>2+</sup> transients were eliminated. These observations suggest that hyperphosphorylation of RyR at S2808 also is not sufficient to induce Ca<sup>2+</sup> release mediated arrhythmias in miR-1 myocytes. Therefore, neither CaMKII phosphorylation of RyR S2814 nor PKA phosphorylation of RyR S2808 alone appear to be sufficient to produce the alterations in RyR behavior that underlie arrhythmogenic SR Ca<sup>2+</sup> release. Hyperphosphorylation of both RyR S2814 and S2808 appear to be necessary for this process. Fortunately the model systems to test these ideas are available and hopefully these issues can be resolved.

In summary, 2 new articles in this issue of *Circulation Research* have identified novel mechanisms for inducing pathological hypertrophy and arrhythmias in cardiac myocytes by altering signaling and contractile Ca<sup>2+</sup>. New studies will need to confirm these results and determine whether the responsible molecules are good targets for novel therapies for pathological cardiac hypertrophy and arrhythmias.

### Sources of Funding

Supported by NIH grant HL33920.

### Disclosures

None.

### References

- Bers DM. Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol.* 2008;70:23–49.
- Houser SR, Molkenin JD Does contractile Ca<sup>2+</sup> control calcineurin-NFAT signaling and pathological hypertrophy in cardiac myocytes? *Sci Signal.* 2008;1:pe31.
- Anderson ME. Multiple downstream proarrhythmic targets for calmodulin kinase II: moving beyond an ion channel-centric focus. *Cardiovasc Res.* 2007;73:657–666.
- Chiang C-S, Huang C-H, Chieng H, Chang Y-T, Chang D, Chen J-J, Chen Y-C, Chen Y-H, Shin H-S, Campbell KP, Chen C-C. The Ca<sub>v</sub>3.2 T-Type Ca<sup>2+</sup> channel is required for pressure overload-induced cardiac hypertrophy in mice. *Circ Res.* 2009;104:514–521.
- Terentyev D, Belevych AE, Terentyeva R, Martin MM, Malana GE, Kuhn DE, Abdellatif M, Feldman DS, Elton TS, Gyorke S. *miR-1* overexpression enhances Ca<sup>2+</sup> release and promotes cardiac arrhythmogenesis by targeting PP2A regulatory subunit B56 $\alpha$  and causing CaMKII-dependent hyperphosphorylation of RyR2. *Circ Res.* 2009;104:522–530.
- Thum T, Galuppo P, Wolf C, Fiedler J, Kneitz S, van Laake LW, Doevendans PA, Mummery CL, Borlak J, Haverich A, Gross C, Engelhardt S, Ertl G, Bauersachs J. MicroRNAs in the human heart: a clue to fetal gene reprogramming in heart failure. *Circulation.* 2007;116:258–267.
- Nuss HB, Houser SR. T-type Ca<sup>2+</sup> current is expressed in hypertrophied adult feline left ventricular myocytes. *Circ Res.* 1993;73:777–782.
- Martinez ML, Heredia MP, Delgado C. Expression of T-type Ca(2+) channels in ventricular cells from hypertrophied rat hearts. *J Mol Cell Cardiol.* 1999;31:1617–1625.
- Vassort G, Talavera K, Alvarez JL. Role of T-type Ca<sup>2+</sup> channels in the heart. *Cell Calcium.* 2006;40:205–220.
- Mangoni ME, Traboulsie A, Leoni AL, Couette B, Marger L, Le Quang K, Kupfer E, Cohen-Solal A, Vilar J, Shin HS, Escande D, Charpentier F, Nargeot J, Lory P. Bradycardia and slowing of the atrioventricular conduction in mice lacking CaV3.1/alpha1G T-type calcium channels. *Circ Res.* 2006;98:1422–1430.
- Chen CC, Lamping KG, Nuno DW, Barresi R, Prouty SJ, Lavoie JL, Cribbs LL, England SK, Sigmund CD, Weiss RM, Williamson RA, Hill JA, Campbell KP. Abnormal coronary function in mice deficient in alpha1H T-type Ca<sup>2+</sup> channels. *Science.* 2003;302:1416–1418.
- Wilkins BJ, Molkenin JD. Calcium-calcineurin signaling in the regulation of cardiac hypertrophy. *Biochem Biophys Res Commun.* 2004;322:1178–1191.
- Shen T, Liu Y, Randall WR, Schneider MF. Parallel mechanisms for resting nucleocytoplasmic shuttling and activity dependent translocation provide dual control of transcriptional regulators HDAC and NFAT in skeletal muscle fiber type plasticity. *J Muscle Res Cell Motil.* 2006;27:405–411.
- Colella M, Grisan F, Robert V, Turner JD, Thomas AP, Pozzan T. Ca<sup>2+</sup> oscillation frequency decoding in cardiac cell hypertrophy: role of calcineurin/NFAT as Ca<sup>2+</sup> signal integrators. *Proc Natl Acad Sci U S A.* 2008;105:2859–2864.
- Zhang R, Khoo MS, Wu Y, Yang Y, Grueter CE, Ni G, Price EE Jr, Thiel W, Guatimosim S, Song LS, Madu EC, Shah AN, Vishnivetskaya TA, Atkinson JB, Gurevich VV, Salama G, Lederer WJ, Colbran RJ, Anderson ME. Calmodulin kinase II inhibition protects against structural heart disease. *Nat Med.* 2005;11:409–417.
- Venetucci LA, Trafford AW, O'Neill SC, Eisner DA. The sarcoplasmic reticulum and arrhythmogenic calcium release. *Cardiovasc Res.* 2008;77:285–292.

KEY WORDS: cardiac hypertrophy ■ arrhythmias ■ sudden death ■ Ca<sup>2+</sup> handling

# Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## Ca<sup>2+</sup> Signaling Domains Responsible For Cardiac Hypertrophy and Arrhythmias Steven R. Houser

*Circ Res.* 2009;104:413-415

doi: 10.1161/CIRCRESAHA.109.193821

*Circulation Research* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2009 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circres.ahajournals.org/content/104/4/413>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation Research* is online at:  
<http://circres.ahajournals.org/subscriptions/>