

# Letter to the Editor

Letters to the Editor will be published, if suitable, as space permits. They should not exceed 1000 words (typed double-spaced) in length and may be subject to editing or abridgment.

## STIM1/Orai1, $I_{CRAC}$ , and Endothelial SOC

To the Editor:

Store-operated calcium ( $Ca^{2+}$ ) entry (SOCE) is an evolutionarily conserved pathway of regulated  $Ca^{2+}$  entry into cells whereby the depletion of  $Ca^{2+}$  from internal stores signals the activation of plasma membrane SOC channels.<sup>1,2</sup> Nonselective canonical transient receptor potential channel (TRPC) proteins were investigated as potential components of SOC channels in mammals. However, the resulting literature offer no consensus among investigators, and the role of TRPCs in encoding SOC remains a contentious issue.<sup>2</sup> In our recent study,<sup>3</sup> we challenged the role of TRPCs in endothelial cell (EC) SOCE by showing evidence for  $I_{CRAC}$  in ECs encoded by STIM1 and Orai1 independently of TRPC1 and TRPC4. In a letter to *Circulation Research*,<sup>4</sup> Beech pointed out the discrepancies between previously published data and ours and contended: (1) “that although the biophysical properties of the  $I_{CRAC}$  recorded in divalent-free (DVF) solutions are convincing, its small size in the presence of extracellular  $Ca^{2+}$  and intracellular BAPTA makes its existence uncertain and that observation of current with BAPTA in the pipette is not proof that it is store-operated”; and (2) we “failed to provide confidence for the knockdown of TRPC1.” We reject these contentions and maintain that  $I_{CRAC}$  encoded by STIM1/Orai1 accounts for EC SOCE based on the following evidence:

First, it has been clearly established that  $I_{CRAC}$  can be activated to the same extent with any means that deplete the stores; these include: BAPTA or inositol-1,4,5-trisphosphate ( $IP_3$ ) dialysis through the pipette; SERCA inhibitors or ionomycin applied to the bath; sensitizing  $IP_3$  receptors to resting levels of  $IP_3$  with thimerosal; and loading of stores with the membrane-permeable metal  $Ca^{2+}$  chelator TPEN.<sup>1,2</sup> We maintain that all of our recordings in which store depletion was elicited by either thapsigargin or BAPTA in 10 mmol/L external  $Ca^{2+}$  consistently produced currents with similar sizes, displaying the biophysical and pharmacological characteristics of  $I_{CRAC}$  (total block by 1 to 10  $\mu$ mol/L gadolinium or 30  $\mu$ mol/L 2-APB).

Second, HEK293 cells show almost undetectable endogenous  $I_{CRAC}$  in  $Ca^{2+}$ -containing solutions that is revealed only in divalent-free (DVF) solutions<sup>5</sup>; yet in these cells,  $I_{CRAC}$  is largely responsible for SOCE and is encoded by Orai1.<sup>6</sup> Thorough comparisons of SOC currents in HEK293 cells, RBL cells, and ECs by Fasolato and Nilius<sup>7</sup> failed to detect nonselective SOC currents in these cells and described  $I_{CRAC}$  currents in ECs with similar size to ours. Furthermore, estimations of  $Ca^{2+}$  entry in human umbilical vein ECs in the presence of 10 mmol/L extracellular  $Ca^{2+}$  suggests that when entry through highly  $Ca^{2+}$ -selective  $I_{CRAC}$  occurs (considering  $Ca^{2+}$  fraction of the current 100%;  $P_{Ca}/P_{Na} \approx 1000$ ),  $\approx 1$  pA is needed to reach global micromolar  $Ca^{2+}$  concentrations.<sup>8</sup> On the other hand, TRPCs are bona fide nonselective channels; a major portion of their currents is carried by sodium. There is no evidence that TRPCs conduct  $Ca^{2+}$  under native conditions; the agonist-induced  $Ca^{2+}$  entry measured by Fura2 imaging when TRPCs are expressed in HEK293 cells might correspond to an artifact of overexpression.<sup>9</sup> Beech<sup>4</sup> refers to the study by Freichel et al<sup>10</sup> describing  $I_{CRAC}$ -like currents in ECs from wild-type mice but not in TRPC4

knockout mice; however, Beech but did not elaborate on how these  $Ca^{2+}$ -selective currents resembling  $I_{CRAC}$  could be reconciled with either nonselective TRPC4 channels as their molecular correlates or nonselective “SOC” currents recorded by other groups. Based on  $Ca^{2+}$ -imaging experiments using ectopic expression of Orai and TRPCs in HEK293 cells, a recent report proposed an attractive model where Orai/TRPC complexes would mediate SOCE.<sup>11</sup> However, it is not clear how the ectopic expression in HEK293 cells relates to native conditions, and the data reported would need to be confirmed under voltage-clamp conditions. Although the results by Freichel et al on TRPC4 knockout mice are quite puzzling,<sup>10</sup> this same group questioned a role for TRPCs as a component of  $I_{CRAC}$  in a subsequent perspective.<sup>12</sup>

Third, as is the case with any scientific question, it is best to use a combination of experimental approaches. Although patch clamp is the most direct and unambiguous method for studying SOC, when SOC currents appear very small, the advantage of fluorescent indicators over direct current measurements is their greater sensitivity to small  $Ca^{2+}$  changes. The use of several SOCE inhibitors at similar concentrations for patch-clamp recordings and Fura2 imaging strengthens the results. We would like to point out that in our study: (1) SOCE measured with Fura2 imaging and  $I_{CRAC}$  had similar pharmacological profiles; (2) most importantly, SOCE in response to thapsigargin and to agonist (thrombin) had similar pharmacological profiles and was both essentially abrogated on STIM1 and Orai1 knockdown; (3) in these cases, SOCE was rescued by STIM1 and Orai1 expression; and (4) SOCE in ECs has one of the most peculiar characteristics of Orai1 (potentiation with 5  $\mu$ mol/L 2-APB and block with 30 to 50  $\mu$ mol/L). Taken together, these results strongly argue that EC SOCE is mediated by  $I_{CRAC}$  encoded by STIM1/Orai1.

Fourth, in our study, we ruled out the involvement of TRPC1 and TRPC4 in SOCE and  $I_{CRAC}$  using 2 independent small interfering (si)RNA sequences.<sup>3</sup> Beech<sup>4</sup> questioned the specificity of the TRPC1 antibody used in our study. However, the same anti-TRPC1 antibody was used in studies cited by Beech,<sup>4</sup> from groups that proposed TRPC1 as a SOC component.<sup>13</sup> In our study, 2 independent siRNA against TRPC1 significantly decreased the TRPC1 mRNA (see Figure 7A in our article) and reduced the TRPC1 protein band to the level of background; Figure 7B in our article shows a statistical analysis of the anti-TRPC1 Western blots where densitometry of TRPC1 bands relative to  $\beta$ -actin is depicted.<sup>3</sup> Beech argued that only a small fraction of previous studies relied on inhibitory antibodies as evidence for a role of TRPCs in SOCE. We strongly disagree with this statement; to our knowledge, there is no molecular evidence that store depletion activates TRPC currents when intracellular  $Ca^{2+}$  is buffered, and only 2 studies reported attempts at TRPC knockdown in ECs, and both used the less specific technique of antisense RNA.<sup>14,15</sup> In these studies, currents were recorded with low intracellular buffering (1.15 mmol/L EGTA or 1 mmol/L BAPTA)<sup>14,15</sup> such that  $Ca^{2+}$ -activated TRPC-mediated currents could have been a confounding factor.

Five, in the original siRNA screens identifying STIM1 and Orai1 as components of SOCE by 4 independent groups, none showed involvement of the 28 members of the larger TRP family.<sup>6</sup> Studies by Gudermann and colleagues showed that smooth muscle isolated from aorta and cerebral arteries of TRPC1 knockout mice possess SOCE currents that were comparable to those recorded in cells from wild-type mice.<sup>16</sup> Furthermore, the replenishment of Ca<sup>2+</sup> stores is normal in central neurons from TRPC3 knockout mice and TRPC1/4/6 triple knockout mice,<sup>17</sup> arguing that Ca<sup>2+</sup> entry through TRPC channels is not required for refilling of internal stores.

In fact, the biophysical and pharmacological heterogeneity of the presumed nonselective "SOC" currents recorded by different groups is quite astonishing (see elsewhere,<sup>2</sup> including references therein). Many of the discrepancies between different groups and controversies regarding TRPCs encoding SOC channels may originate from differences in techniques, patch-clamp protocols, and concentrations of pharmacological reagents used. Our study<sup>3</sup> used a combination of patch clamp and Fura2 imaging with 2 pharmacological inhibitors at similar concentrations in both techniques. We used molecular tools to knockdown in parallel in 2 different EC types several molecular candidates of the SOCE pathway (STIM1, Orai1, TRPC1, and TRPC4). Therefore, we believe our study provides significant insights into the mechanism of SOCE in ECs. It is certainly conceivable that TRPCs in ECs, as well as smooth muscle, carry out important signaling functions, albeit through mechanisms not necessarily dependent on calcium-store depletion.

### Sources of Funding

Research in the laboratory of the author is supported by NIH grant K22ES014729.

### Disclosures

None.

**Mohamed Trebak**

Center for Cardiovascular Sciences  
Albany Medical College  
Albany, NY  
E-mail: [trebakm@mail.amc.edu](mailto:trebakm@mail.amc.edu)

1. Parekh AB, Penner R. Store depletion and calcium influx. *Physiol Rev.* 1997;77:901–930.
2. Parekh AB, Putney JW Jr. Store-operated calcium channels. *Physiol Rev.* 2005;85:757–810.
3. Abdullaev IF, Bisailon JM, Potier M, Gonzalez JC, Motiani RK, Trebak M. Stim1 and Orai1 mediate CRAC currents and store-operated calcium

- entry important for endothelial cell proliferation. *Circ Res.* 2008;103:1289–1299.
4. Beech DJ. Harmony and discord in endothelial calcium entry. *Circ Res.* 2009;104:e22–e23.
5. DeHaven WI, Smyth JT, Boyles RR, Putney JW Jr. Calcium inhibition and calcium potentiation of Orai1, Orai2, and Orai3 calcium release-activated calcium channels. *J Biol Chem.* 2007;282:17548–17556.
6. Vig M, Peinelt C, Beck A, Koomoa DL, Rabah D, Koblan-Huberson M, Kraft S, Turner H, Fleig A, Penner R, Kinet JP. CRACM1 is a plasma membrane protein essential for store-operated Ca<sup>2+</sup> entry. *Science.* 2006;312:1220–1223.
7. Fasolato C, Nilius B. Store depletion triggers the calcium release-activated calcium current (ICRAC) in macrovascular endothelial cells: a comparison with Jurkat and embryonic kidney cell lines. *Pflugers Arch.* 1998;436:69–74.
8. Oike M, Gericke M, Droogmans G, Nilius B. Calcium entry activated by store depletion in human umbilical vein endothelial cells. *Cell calcium.* 1994;16:367–376.
9. Rosker C, Graziani A, Lukas M, Eder P, Zhu MX, Romanin C, Groschner K. Ca(2+) signaling by TRPC3 involves Na(+) entry and local coupling to the Na(+)/Ca(2+) exchanger. *J Biol Chem.* 2004;279:13696–13704.
10. Freichel M, Suh SH, Pfeifer A, Schweig U, Trost C, Weissgerber P, Biel M, Philipp S, Freise D, Droogmans G, Hofmann F, Flockerzi V, Nilius B. Lack of an endothelial store-operated Ca<sup>2+</sup> current impairs agonist-dependent vasorelaxation in TRP4<sup>-/-</sup> mice. *Nat Cell Biol.* 2001;3:121–127.
11. Liao Y, Plummer NW, George MD, Abramowitz J, Zhu MX, Birnbaumer L. A role for Orai in TRPC-mediated Ca<sup>2+</sup> entry suggests that a TRPC:Orai complex may mediate store and receptor operated Ca<sup>2+</sup> entry. *Proc Natl Acad Sci U S A Mar 3.* 2009;106:3202–3206.
12. Nilius B. Store-operated Ca<sup>2+</sup> entry channels: still elusive! *Sci STKE.* 2004;2004:pe36.
13. Ahmmed GU, Mehta D, Vogel S, Holinstat M, Paria BC, Tiruppathi C, Malik AB. Protein kinase Cα phosphorylates the TRPC1 channel and regulates store-operated Ca<sup>2+</sup> entry in endothelial cells. *J Biol Chem.* 2004;279:20941–20949.
14. Brough GH, Wu S, Cioffi D, Moore TM, Li M, Dean N, Stevens T. Contribution of endogenously expressed Trp1 to a Ca<sup>2+</sup>-selective, store-operated Ca<sup>2+</sup> entry pathway. *FASEB J.* 2001;15:1727–1738.
15. Paria BC, Vogel SM, Ahmmed GU, Alamgir S, Shroff J, Malik AB, Tiruppathi C. Tumor necrosis factor-α-induced TRPC1 expression amplifies store-operated Ca<sup>2+</sup> influx and endothelial permeability. *Am J Physiol Lung Cell Mol Physiol.* 2004;287:L1303–L1313.
16. Dietrich A, Kalwa H, Storch U, Mederos y Schnitzler M, Salanova B, Pinkenburg O, Dubrovskaya G, Essin K, Gollasch M, Birnbaumer L, Gudermann T. Pressure-induced and store-operated cation influx in vascular smooth muscle cells is independent of TRPC1. *Pflugers Arch.* 2007;455:465–477.
17. Hartmann J, Dragicevic E, Adelsberger H, Henning HA, Sumser M, Abramowitz J, Blum R, Dietrich A, Freichel M, Flockerzi V, Birnbaumer L, Konnerth A. TRPC3 channels are required for synaptic transmission and motor coordination. *Neuron.* 2008;59:392–398.

# Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## STIM1/Orai1, *I<sub>CRAC</sub>*, and Endothelial SOC Mohamed Trebak

*Circ Res.* 2009;104:e56-e57

doi: 10.1161/CIRCRESAHA.109.196105

*Circulation Research* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2009 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circres.ahajournals.org/content/104/9/e56>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation Research* is online at:  
<http://circres.ahajournals.org/subscriptions/>