

This Review is the Introduction of a new thematic series on **Wnts in Cardiovascular Development and Disease**, which includes the following articles:

An Updated Overview on Wnt Signaling Pathways: A Prelude for More

The Multiple Phases and Faces of Wnt Signaling During Cardiac Differentiation and Development

Wnt Signaling in Vascular Progenitor Cells and Angiogenesis

Wnt Signaling in Cardiac Hypertrophy and Remodeling

Wnt Signaling in Heart Failure and Aging

Wnt Signaling and Stem Cells

Michael Kühl, Guest Editor

## An Updated Overview on Wnt Signaling Pathways A Prelude for More

Tata Purushothama Rao, Michael Kühl

**Abstract:** Growth factor signaling is required for cellular differentiation, tissue morphogenesis, and tissue homeostasis.

Misregulation of intracellular signal transduction can lead to developmental defects during embryogenesis or particular diseases in the adult. One family of growth factors important for these aspects is given by the Wnt proteins. In particular, Wnts have important functions in stem cell biology, cardiac development and differentiation, angiogenesis, cardiac hypertrophy, cardiac failure, and aging. Knowledge of growth factor signaling during differentiation will allow for improvement of targeted differentiation of embryonic or adult stem cells toward functional cardiomyocytes or for understanding the basis of diseases. Our major aim here is to provide a state of the art review summarizing our present knowledge of the intracellular Wnt-mediated signaling network. In particular, we provide evidence that the subdivision into canonical and noncanonical Wnt signaling pathways solely based on the identity of Wnt ligands or Frizzled receptors is not appropriate anymore. We thereby deliver a solid base for further upcoming articles of a review series focusing on the role of Wnt proteins on different aspects of cardiovascular development and dysfunction. (*Circ Res.* 2010;106:1798-1806.)

**Key Words:** Wnt ■ frizzled ■ canonical ■ noncanonical

Wnt proteins are secreted glycoproteins acting as short or long range signaling molecules. To trigger a cellular response and to activate intracellular signal transduction, they bind to receptors of the Frizzled family and several coreceptors such as lipoprotein receptor-related protein (LRP)-5/6, Ryk, or Ror2.<sup>1,2</sup> In humans, 19 members of the Wnt family and 10 Frizzled receptors are known. Based on different biological readouts, Wnt ligands, as well as Frizzled receptors, were subdivided into different subclasses, eg, some Wnts are able to induce secondary body axes when injected into *Xenopus* embryos or to transform C57 mg cells, whereas other Wnts regulate cell adhesion and morphogenetic movements.

### Wnt Proteins Activate Apparently Distinct Signaling Pathways

#### Canonical Wnt/ $\beta$ -Catenin Signaling

On a molecular level, these different biological activities are mediated through different intracellular signaling pathways (Figure 1). Axis duplication and cell transformation are based on the activation of the canonical Wnt pathway that involves the multifunctional protein  $\beta$ -catenin. In the absence of Wnt,  $\beta$ -catenin is targeted to a multimeric protein complex called destruction complex for its phosphorylation. This is achieved by CK1-mediated phosphorylation at Ser45, followed by

Original received March 5, 2010; revision received April 23, 2010; accepted May 11, 2010.

From the Institute for Biochemistry and Molecular Biology, Ulm University, Germany.

Correspondence to Michael Kühl, Institute for Biochemistry and Molecular Biology, Ulm University, Albert-Einstein-Allee 11, D-89081 Ulm, Germany. E-mail michael.kuehl@uni-ulm.de

(*Circ Res.* 2010;106:1798-1806.)

© 2010 American Heart Association, Inc.

*Circulation Research* is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.110.219840

Ser33, Ser37, and Thr41 phosphorylation by glycogen synthase kinase (GSK)3 $\beta$ .<sup>3</sup> This phosphorylation targets  $\beta$ -catenin for  $\beta$ -Trcp-mediated ubiquitination and its subsequent degradation by the proteasome. Deletion of these phosphorylation sites either by point mutations (eg, S33A) or by larger deletions, therefore, result in stabilization of  $\beta$ -catenin. A widely used mouse model is a floxed exon 3 of  $\beta$ -catenin allowing a conditional deletion of exon 3 that codes for these amino acids. This mouse model shows an increased  $\beta$ -catenin stability and T-cell factor (TCF)/lymphocyte enhancer factor (LEF)-dependent transcription.<sup>4</sup>

In the presence of some Wnt ligands, a cascade of events initiated at the plasma membrane by the binding of Wnt to the cysteine-rich domain of Frizzled receptors results in the disassembly of the destruction complex consisting of axin, adenomatous polyposis coli (APC), and GSK3 $\beta$  and the stabilization of  $\beta$ -catenin. This process also involves the phosphoprotein Dishevelled. Cytoplasmic  $\beta$ -catenin accumulates and is eventually imported into the nucleus, where it serves as a transcriptional coactivator of transcription factors of the TCF/LEF family. In mice, deletion of exon 3 to 6 results in a  $\beta$ -catenin loss-of-function situation because this

Non-standard Abbreviations and Acronyms	
APC	adenomatous polyposis coli
CaMKII	calcium/calmodulin-dependent kinase II
GSK	glycogen synthase kinase
JNK	jun N-terminal kinase
HDAC	histone deacetylase
LEF	lymphocyte enhancer factor
LRP	lipoprotein receptor-related protein
NFAT	nuclear factor of activated T cells
NLK	nemo-like kinase
PK	protein kinase
TCF	T-cell factor

truncated form of  $\beta$ -catenin is not able to function in transcriptional activation.<sup>4</sup> TCF/LEF target genes are then involved in regulating cell proliferation, stem cell maintenance, or differentiation. Interestingly,  $\beta$ -catenin can also interact with other transcription factors such as Prop1, Oct-




Figure 1. Overview of canonical (left) and noncanonical Wnt signaling pathways (right) as discussed in the text. CaCN indicates calcineurin.

3/4, Pitx2, or Mitf.<sup>5–8</sup> This pathway has been linked to cardiac development<sup>9</sup> and angiogenesis,<sup>10</sup> which are 2 aspects of cardiovascular development that will be discussed in 2 follow-up reviews.

### **$\beta$ -Catenin As a Multifunctional Protein**

One must keep in mind, however, that  $\beta$ -catenin is a multifunctional protein and interacts with additional proteins such as cadherins.<sup>11</sup> This interaction with  $\beta$ -catenin is essential for the function of cadherins in cell adhesion by establishing a link to the actin cytoskeleton. In this context, the closely related  $\gamma$ -catenin (plakoglobin) can also replace  $\beta$ -catenin. This interaction of cadherins with  $\beta$ -catenin is also tightly regulated by phosphorylation. Phosphorylation of  $\beta$ -catenin at Tyr654 through src kinase, as well as Thr112 and Thr120 by protein kinase (PK)D1, enhances the interaction with E-cadherin.<sup>12</sup> Conversely, phosphorylation of E-cadherin at Ser834, Ser836, and Ser842 by CK2 and GSK3 $\beta$  enhances its interaction with  $\beta$ -catenin. In contrast, phosphorylation at Tyr831 and Tyr860 by src kinase and Ser846 by CK1 inhibits the interaction of E-cadherin with  $\beta$ -catenin.<sup>13</sup> Taken together, these data indicate that the role of  $\beta$ -catenin in Wnt-mediated signal transduction and cell adhesion is highly dependent on its phosphorylation status at multiple sites.<sup>14,15</sup>

Members of the cadherin and catenin families are also required for adhesion between cardiomyocytes. This also includes the desmosomal cadherins desmocollin and desmoglein, the catenins plakoglobin ( $\gamma$ -catenin) and plakophilin2, or desmoplakin, which links the cadherin/catenin complex to intermediate filaments. Mutations in these components are the cause of arrhythmogenic right ventricular cardiomyopathies. In combination with woolly hair and palmoplantar keratoderma this congenital disease is also called Naxos disease.<sup>16,17</sup> This disease is characterized by ventricular arrhythmias and a progressive loss of cardiomyocytes that are replaced by fibrofatty tissue. Recent data suggested that this phenotype is not only caused by defects in cell adhesion. Loss of desmoplakin results in nuclear import of  $\gamma$ -catenin (plakoglobin) that negatively interferes with Wnt/ $\beta$ -catenin signaling.<sup>18</sup> Fine-tuned regulation of cadherin-mediated cell adhesion is also required for epithelial–mesenchymal or mesenchymal–epithelial transitions. The interplay of cadherin-mediated cell adhesion and Wnt signaling has recently also been reviewed in detail by others.<sup>14</sup>

### **$\beta$ -Catenin–Independent Noncanonical Wnt Pathways**

In contrast, all Wnt signaling activities that are apparently independent of  $\beta$ -catenin constitute different noncanonical Wnt signaling pathways. Depending on the major intracellular mediators used, those are called the Wnt/jun N-terminal kinase (JNK) or Wnt/calcium pathway, respectively. The Wnt/JNK pathway has a high degree of overlap with the planar cell polarity pathway originally described in *Drosophila*.<sup>19</sup> It involves activation of small GTPases of the rho family including rac, cdc42, and rho and further downstream protein kinases such as JNK or rho kinase. In this branch, Frizzled and dishevelled function in concert with other proteins to set up cellular polarity by asymmetrical and

polarized protein localization. Those include classic and novel components of the planar cell polarity pathway like vangl/Strabismus,<sup>20–22</sup> Celsr,<sup>23</sup> Prickle,<sup>24</sup> or PTK7<sup>25</sup> in a context-dependent manner. Most of these proteins have been shown to regulate polarized cell movements (eg, during gastrulation, neural crest migration, or cardiac outflow tract development) and planar polarity of epithelial cells (eg, wing hair development in *Drosophila* or stereociliary bundles in the vertebrate cochlea). Furthermore, noncanonical Wnt signaling has also been linked to ciliogenesis.<sup>26</sup>

The existence of a calcium-mediated pathway that is directly activated by Wnt proteins was originally described in zebrafish and *Xenopus*.<sup>27,28</sup> These pathways have been linked to cardiac development in several settings,<sup>29–33</sup> as discussed in more detail in another review of this series. A long-lasting debate whether the observed intracellular release of calcium is a direct response or reflects some indirect effects was solved by showing that purified Wnt5a is able to activate calcium signaling in different cell culture models.<sup>34–37</sup> This process is very rapid and depends on heterotrimeric G proteins.<sup>36</sup> The released Ca<sup>2+</sup> then activates calcium dependent enzymes like calcium/calmodulin-dependent kinase (CaMK)II, PKC or calcineurin.<sup>38,39</sup> In particular the regulation of PKC by Wnt ligands is important for cardiac differentiation.<sup>29,40,41</sup> PKC comes in different isoforms and a detailed analysis suggested PKC $\delta$  to play an important role during noncanonical Wnt signaling during cardiac differentiation.<sup>40</sup>


Through calcineurin the Wnt/calcium pathway connects to NFAT (nuclear factor of activated T cells) transcription factor<sup>35,42</sup> and gene expression. Interestingly, cardiac hypertrophy involves, beside other signaling pathways, calcineurin and its downstream target NFAT.<sup>43</sup> The involvement of both canonical and noncanonical Wnt signaling during cardiac hypertrophy and cardiac remodeling will therefore be discussed in great detail in a forthcoming review of this series.

On the other hand CaMKII can lead to activation of a nemo-like kinase (NLK), which interferes with  $\beta$ -catenin signaling.<sup>44,45</sup> Although not formally proven, the activation of CaMKII suggests that noncanonical Wnt signaling could modify the activity of histone deacetylase (HDAC). In particular, HDAC4 and -5 are regulated through CaMKII or CaMKIV,<sup>46–50</sup> and this is required for the expression of early cardiac genes such as Nkx2.5, MEF2C, or GATA-4.<sup>46</sup> This would not only provide an additional link to gene regulation through noncanonical Wnt signaling but suggests that a misregulation of noncanonical Wnt signaling could result in pathophysiological conditions in the heart that had been linked to CaMKII or HDACs.<sup>48,51</sup> One interesting finding here is that noncanonical Wnt signaling has been shown to modulate the activity of a histone lysine methyltransferase in a different setting.<sup>52</sup> Here, noncanonical Wnt signaling through the CaMKII-NLK axis results in H3-K9 methylation of histones to regulate gene expression.

### **Wnt Signaling at the Plasma Membrane**

#### **Wnt and Frizzleds: Monomers, Heterodimers, and Homodimers**

Based on the biological assays mentioned, Wnts were subdivided into canonical or noncanonical Wnts, and the same



**Figure 2. Wnt signaling at the membrane.** Relative concentrations of different components determine pathway specificity.

holds true for Frizzled receptors,<sup>39</sup> eg, Wnt1, Wnt3a, or Wnt8 were considered to activate canonical Wnt signaling, whereas Wnt5a or Wnt11 are best known for their ability to trigger noncanonical Wnt signaling. This model of signaling specificity has been challenged by the finding that so-called noncanonical Wnt ligands can activate  $\beta$ -catenin signaling in the presence of appropriate Frizzled receptors and vice versa. “Canonical” Wnt3a has the ability to activate PKC during bone formation<sup>53</sup> and the bona fide “noncanonical” Wnt5a couples to canonical Wnt signaling in the presence of Frizzled-5.<sup>54</sup> Just recently noncanonical Wnt11 was shown to activate  $\beta$ -catenin signaling during axis specification in *Xenopus*.<sup>55</sup> These data indicate that differences in affinities of different Wnt and Frizzleds and their local concentrations (in vivo but also experimentally) would determine to which signaling branch a particular Wnt couples. Extending these earlier findings, it was recently shown that Wnt5a and Wnt11 proteins can interact physically<sup>56</sup> and thereby induce  $\beta$ -catenin signaling. Tyrosyl protein sulfotransferase-1-dependent *O*-sulfation of specific tyrosine residues of Wnt11 is required for this interaction with Wnt5a.<sup>57</sup> Similarly, Frizzled-4 has been shown to activate either  $\beta$ -catenin signaling<sup>58</sup> or noncanonical Wnt signaling branches.<sup>59</sup> There is also evidence for Frizzled dimers and multimers.<sup>60–62</sup>

These findings indicate that the analysis of a particular Wnt ligand or Frizzled receptor in a certain biological context always requires an investigation of how the Wnt signal is transduced. However, this leaves with challenging questions: How is Wnt signaling initiated at the plasma membrane? And how is specificity achieved?

### LRP Phosphorylation Generates a Docking Site for Axin

Activation of canonical Wnt/ $\beta$ -catenin signaling requires the interaction of a Wnt ligand with Frizzleds and LRP-5/6. The major outcome of pathway activation is the disintegration of the destruction complex, the release of  $\beta$ -catenin and its subsequent stabilization. This is achieved by binding of dishevelled and axin to Frizzleds and phosphorylated LRP-5/6, respectively (Figure 2). Wnt-induced phosphorylation of LRP-5/6 is mediated by the concerted action of GSK3 $\beta$  and CK1.<sup>63,64</sup> A recent report indicated that in *Drosophila*, a cyclin-dependent kinase (CDK), L63, or its vertebrate homolog PFTK also phosphorylates LRP-5/6.<sup>65</sup> This is an interesting finding because Wnt/ $\beta$ -catenin itself regulates the cell cycle by directly promoting c-Myc and cyclinD1 expression. Furthermore,  $\beta$ -catenin is required for centrosome

separation during mitosis.<sup>66</sup> Apparently, there is an intricate feedback between the cell cycle machinery and Wnt signaling to maintain a high-proliferative state as found in stem cells or tumor cells. Interestingly, some reports indicated that Wnt/ $\beta$ -catenin signaling is required to regulate cardiac progenitor cell proliferation, an issue that is also discussed in detail in a forthcoming review of this series.<sup>67,68</sup> A recent report also claims that Wnt3a stimulates the formation of phosphatidylinositol-4,5-bisphosphates [PtdIns(4,5)P<sub>2</sub>] through Frizzled and dishevelled, the latter of which directly interacts with and activates PIP5KI.<sup>69</sup> PIP5KI then subsequently can phosphorylate LRP5/6. In addition, G protein-coupled receptor kinase 5/6 is able to phosphorylate LRP-5/6.<sup>70</sup> Some evidence suggests that LRP6 also antagonizes noncanonical Wnt signaling in vivo, possibly via competition for Wnt ligands<sup>71</sup> or by an unknown mechanism.<sup>72</sup>

### Local Changes in pH Are Involved in Wnt Signaling

Frizzled binding of dishevelled as part of noncanonical Wnt signaling is pH-dependent.<sup>73</sup> The authors screened for molecules involved in dishevelled membrane recruitment and found that a Na<sup>+</sup>/H<sup>+</sup> exchange activity of the plasma membrane exchanger, Nhe2, is a key player. Just recently, this model has been extended by showing that a local change in pH is also required for Wnt/ $\beta$ -catenin signaling. This is mediated by the prorenin receptor and a vacuolar H<sup>+</sup>ATPase.<sup>74</sup>

### Things at the Membrane Are More Complex: The Involvement of G Proteins

Frizzleds are 7-transmembrane proteins. Those are often associated with heterotrimeric G proteins to initiate intracellular signal transduction. Whereas noncanonical Wnt signaling has been linked to G proteins for a long time,<sup>27,38,75</sup> the lack of genetic evidence was used by some to argue against a role of G proteins in canonical Wnt signal transduction. These days, however, we have mounting evidence that they are involved in a key position. The first findings were obtained by experiments using inducible Frizzled receptors that indicated a requirement for G proteins in both canonical and noncanonical signaling branches, requiring G<sub>αo</sub> and G<sub>αq</sub> or G<sub>αo</sub> and G<sub>αt</sub>, respectively.<sup>76,77</sup> Moreover, activation of a TCF/ $\beta$ -catenin-dependent reporter<sup>78</sup> is blocked by small interfering RNA targeting different G<sub>α</sub> subunits<sup>79</sup> and Frizzled, dishevelled, and axin can directly interact with some G-protein subunits.<sup>80,81</sup> Also, genetic evidence argues for a requirement of G proteins in Wnt signaling.<sup>82</sup>

### Dkk1: Beyond a Wnt Inhibitor

Dkk1 is among the best-characterized inhibitors of the canonical Wnt pathway. Dkk1 itself is a target gene of Wnt/ $\beta$ -catenin signaling, thereby establishing a negative-feedback loop. The Dickkopf family of secreted proteins is conserved among all vertebrates. They are also present in some invertebrates, such as urochordates and ascidians, but are noticeably missing in *Caenorhabditis elegans* and *Drosophila*.<sup>83</sup> Dkk1 consists of 2 cysteine-rich domains at the N and C terminus, respectively. The C-terminal cysteine-rich domain is shown to be responsible for the Wnt inhibitory function.<sup>84,85</sup> Dkk1 inhibits the formation of a ternary complex

consisting of LRP5/6, Frizzled, and the Wnt ligand (Figure 2).<sup>86</sup> Others indicated that Dkk1, together with single pass transmembrane proteins of the Kremen family, mediates LRP5/6 internalization.<sup>87</sup> In the absence of Dkk1, however, Kremen activates Wnt signaling through LRP-5/6.<sup>88</sup> This is an interesting finding because it would allow for a steep gradient of pathway activation in the presence of a Dkk1 protein gradient.<sup>89</sup> Interestingly, inhibiting canonical Wnt signaling by Dkk1 resulted in an activation of noncanonical JNK signaling in some studies.<sup>29,90,91</sup> The N-terminal domain of Dkk1 has additional, not yet characterized signaling activities independent of  $\beta$ -catenin regulation. Interestingly, this activity is involved in regulating cardiogenesis.<sup>92</sup>

Internalization of LRP-5/6 is not only a mechanism to inhibit Wnt signaling by reducing the accessibility of LRP-5/6 for Wnt ligands, but is also required for Wnt signaling. This difference is mainly based on the mechanism by which LRP-5/6 is internalized. Whereas Dkk-induced clathrin-mediated internalization of LRP5/6<sup>93</sup> results in pathway inhibition, Wnt stimulates a caveolin-mediated LRP5/6 internalization that is involved in normal Wnt signaling.<sup>93</sup>

### Atypical Receptor Kinases in Wnt Signaling

Members of the Ryk and Ror families are single span transmembrane receptors with an intracellular tyrosine kinase domain. They can interact with Wnt proteins through their extracellular cysteine-rich domain in the case of Ror proteins or their Wif domain in the case of Ryk proteins.<sup>94–98</sup> With respect to signal transduction, Ror2 has been placed upstream of small rho-GTPases and JNK,<sup>99,100</sup> whereas src is involved in Ryk signaling.<sup>101</sup> Interestingly, Ror2-mediated signal transduction also inhibits  $\beta$ -catenin signaling in a RTK-dependent manner.<sup>96,102</sup> Mutations in Ror2 result in Robinow syndrome, which is characterized by abnormal morphogenesis in the face and limb defects. Many patients also have heart defects,<sup>103</sup> further highlighting the role of noncanonical Wnt signaling in cardiac development.

### Extracellular Modulation of Wnt Signaling

Wnt/ $\beta$ -catenin signaling can also be modulated by extracellular ligands other than Wnts. R-Spondins form a small family with 4 members. Initially, R-Spondin was shown to directly interact with Frizzled-8 and LRP5/6, thereby positively regulating  $\beta$ -catenin signaling.<sup>104</sup> Furthermore, R-Spondin positively regulates canonical Wnt signaling by competing with Dkk1 for binding to Kremen and LRP5/6.<sup>105,106</sup> R-Spondin3 supports angioblast development and vasculogenesis through  $\beta$ -catenin signaling.<sup>107</sup> Norrin is another extracellular protein that can activate  $\beta$ -catenin signaling through interaction with Frizzled-4.<sup>58,108</sup> Signaling through norrin and Fz-4 for example has been linked to vasculogenesis.<sup>108</sup> Furthermore, the extracellular collagen triple helix repeat-containing protein 1 (Cthrc1) interacts with some Wnts and Frizzleds and supports Wnt-Fz-Ror2 complex formation and at the same reduces Wnt-Fz-LRP complex formation. Thus, Cthrc1 might favor noncanonical Wnt signaling (Figure 2).<sup>109</sup>

An additional level of regulation comes in form of secreted Wnt inhibitors, such as Wif proteins and secreted frizzled-related proteins (Sfrps). Wifs and Sfrps can directly bind to

Wnt proteins in the extracellular space, thereby affecting receptor occupancy and, ultimately, the cellular response.<sup>110</sup> Finally, more and more evidence indicate a role for heparin sulfate proteoglycans (HSPG) in the transport, stabilization, and presentation of Wnt.<sup>2</sup> In *Drosophila*, absence of *Dally*, an HSPG,<sup>111</sup> and mutations in genes encoding enzymes that modify HSPG<sup>112</sup> generate phenotypes similar to *wingless* mutants.

In summary, all these reports consistently support the requirement for coordinated changes at the plasma membrane for Wnt mediated signal transduction. The major challenge in the field is to describe how all of these different pieces fit together in a unifying model of receptor activation explaining signaling specificity in vivo.

### Nuclear Components of $\beta$ -Catenin Signaling: From Transcription Factors to Chromatin Remodeling

#### Regulating Nuclear Localization of $\beta$ -Catenin

The mechanisms underlying the nuclear import and export of  $\beta$ -catenin are not well understood but apparently do not require a nuclear localization signal.<sup>113</sup> A recent report indicated that the nuclear import of  $\beta$ -catenin is mediated by Wnt-induced activation of a Rac-JNK pathway by phosphorylation of  $\beta$ -catenin at Ser191 and Ser605.<sup>114</sup> On the other hand, APC and axin were shown to function in nuclear export of  $\beta$ -catenin.<sup>115,116</sup> Furthermore, RanBP3 has been shown to be involved in nuclear export of  $\beta$ -catenin.<sup>117</sup> In the nucleus, Bcl9 and pygopus play a role in retention of  $\beta$ -catenin.<sup>118</sup> In detailed fluorescence recovery after photobleaching experiments, it was recently shown that APC, axin, Bcl9, and TCF factors influence the cytoplasmic/nuclear localization of  $\beta$ -catenin mainly on the level of retention rather than active transport.<sup>118</sup>

#### $\beta$ -Catenin and TCF/LEF Transcription Factors

In the absence of Wnt, TCF/LEF transcription factors bind to Wnt response elements, facilitating the recruitment of corepressors such as Groucho, CtBP, and HDACs to the particular genomic region. In the presence of Wnt, however,  $\beta$ -catenin is imported into the nucleus and binds to TCF/LEF, thereby replacing transcriptional repressors and recruiting additional transcriptional coactivators, in particular chromatin remodeling complexes such as swi/SNF. Those are brought to the TCF/ $\beta$ -catenin complex by Bcl-9/legless and pygopus.<sup>119</sup> Several posttranslational modifications like phosphorylation, sumoylation, ubiquitination, and acetylation modulate the potential of TCF/LEF transcription factors to interact with nuclear coactivators, repressors, or DNA.<sup>120</sup> For example, the phosphorylation of TCF/LEF by activated NLK/Nemo is thought to diminish the DNA-binding affinity of the  $\beta$ -catenin/TCF/LEF complex, thereby affecting transcriptional regulation of Wnt target genes.<sup>44,45</sup>

Interestingly, in some cases, a repressive function of TCF/ $\beta$ -catenin complexes has been described. In *Drosophila*, this apparently involves a different Wnt responsive element, whereas in vertebrates, the well-known consensus sequence is used.<sup>121</sup> These data indicate that the molecular mechanisms underlying the exchange of transcriptional repressors by

transcriptional activators within a TCF/LEF-mediated transcriptional complex are far from being understood.

### Intracellular Signaling Pathways: Toward an Integrative View of Wnt Signaling

The historical model of Wnt signaling assumed the existence of 3 completely separate signaling pathways. Within the last decade, however, many examples were found demonstrating that individual mediators of Wnt signaling are involved in several branches. This includes certain Frizzled receptors that obviously have the ability to elicit different intracellular responses,<sup>58,59</sup> G proteins that are involved in different signaling branches,<sup>27,28,75–82</sup> and continues with the protein disheveled that also links to all three signaling branches.<sup>122</sup> Furthermore, some components seem to favor one signaling pathway at the expense of the other. Finally, as mentioned above, it recently became evident that  $\beta$ -catenin signaling also relies on a simultaneous activation of the Wnt/JNK branch.<sup>114</sup> Given all of these issues, it seems more appropriate to talk of a Wnt signaling network of interwoven signaling branches. This new model of Wnt signaling has been discussed recently in more detail.<sup>1,123</sup> Depending on the cellular context, eg, concentrations of individual components, this would allow for the preferred activation of certain aspects of the network. This, however, has direct implications on the interpretation of experimental results because the status of the network must be analyzed in detail in each experimental setting and biological model.

### Outlook

These discussions indicate that Wnt proteins have multiple functions on a cellular level, including regulation of gene expression via different mechanisms (and thus differentiation) and cell cycle and proliferation, as well as other nongenomic responses such as cell migration or cilia formation. As indicated, a critical analysis of Wnt function in a given context should always include an investigation of mediators used. A simple assignment based on the identity of Wnt ligands or Frizzled receptors is not sufficient. In the forthcoming reviews of this series, different authors will focus on different aspects of Wnt signaling during cardiovascular development and dysfunction, in particular, during cardiac differentiation and development, angiogenesis, and cardiac remodeling, as well as cardiac failure and aging.

### Sources of Funding

Work in our laboratory is supported by the Deutsche Forschungsgemeinschaft (DFG).

### Disclosures

None.

### References

1. Kestler HA, Kuhl M. From individual Wnt pathways towards a Wnt signalling network. *Philos Trans R Soc Lond B Biol Sci.* 2008;363:1333–1347.
2. Logan CY, Nusse R. The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol.* 2004;20:781–810.
3. Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, Zhang Z, Lin X, He X. Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell.* 2002;108:837–847.

4. Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 2006;34:D140–D144.
5. Olson LE, Tollkuhn J, Scafoglio C, Kronen A, Zhang J, Ohgi KA, Wu W, Taketo MM, Kemler R, Grosschedl R, Rose D, Li X, Rosenfeld MG. Homeodomain-mediated beta-catenin-dependent switching events dictate cell-lineage determination. *Cell.* 2006;125:593–605.
6. Takao Y, Yokota T, Koide H. Beta-catenin up-regulates Nanog expression through interaction with Oct-3/4 in embryonic stem cells. *Biochem Biophys Res Commun.* 2007;353:699–705.
7. Kioussi C, Briata P, Baek SH, Rose DW, Hamblet NS, Herman T, Ohgi KA, Lin C, Gleiberman A, Wang J, Brault V, Ruiz-Lozano P, Nguyen HD, Kemler R, Glass CK, Wynshaw-Boris A, Rosenfeld MG. Identification of a Wnt/Dvl/beta-Catenin → Pitx2 pathway mediating cell-type-specific proliferation during development. *Cell.* 2002;111:673–685.
8. Schepsky A, Bruser K, Gunnarsson GJ, Goodall J, Hallsson JH, Goding CR, Steingrimsson E, Hecht A. The microphthalmia-associated transcription factor Mitf interacts with beta-catenin to determine target gene expression. *Mol Cell Biol.* 2006;26:8914–8927.
9. Tzahor E. Wnt/beta-catenin signaling and cardiogenesis: timing does matter. *Dev Cell.* 2007;13:10–13.
10. Goodwin AM, D'Amore PA. Wnt signaling in the vasculature. *Angiogenesis.* 2002;5:1–9.
11. Bullions LC, Levine AJ. The role of beta-catenin in cell adhesion, signal transduction, and cancer. *Curr Opin Oncol.* 1998;10:81–87.
12. Du C, Jaggi M, Zhang C, Balaji KC. Protein kinase D1-mediated phosphorylation and subcellular localization of beta-catenin. *Cancer Res.* 2009;69:1117–1124.
13. Dupre-Crochet S, Figueroa A, Hogan C, Ferber EC, Bialucha CU, Adams J, Richardson EC, Fujita Y. Casein kinase 1 is a novel negative regulator of E-cadherin-based cell-cell contacts. *Mol Cell Biol.* 2007;27:3804–3816.
14. Heuberger J, Birchmeier W. Interplay of cadherin-mediated cell adhesion and canonical wnt signaling. *Cold Spring Harb Perspect Biol.* 2010;2:a002915.
15. Daugherty RL, Gottardi CJ. Phospho-regulation of Beta-catenin adhesion and signaling functions. *Physiology (Bethesda).* 2007;22:303–309.
16. McKoy G, Protonotarios N, Crosby A, Tsatsopoulou A, Anastasakis A, Coonar A, Norman M, Baboonian C, Jeffery S, McKenna WJ. Identification of a deletion in plakoglobin in arrhythmic right ventricular cardiomyopathy with palmoplantar keratoderma and woolly hair (Naxos disease). *Lancet.* 2000;355:2119–2124.
17. Sen-Chowdhry S, Morgan RD, Chambers JC, McKenna WJ. Arrhythmic cardiomyopathy: etiology, diagnosis, and treatment. *Annu Rev Med.* 2010;61:233–253.
18. Garcia-Gras E, Lombardi R, Giocondo MJ, Willerson JT, Schneider MD, Khoury DS, Marian AJ. Suppression of canonical Wnt/beta-catenin signaling by nuclear plakoglobin recapitulates phenotype of arrhythmic right ventricular cardiomyopathy. *J Clin Invest.* 2006;116:2012–2021.
19. Simons M, Mlodzik M. Planar cell polarity signaling: from fly development to human disease. *Annu Rev Genet.* 2008;42:517–540.
20. Kibar Z, Torban E, McDearmid JR, Reynolds A, Berghout J, Mathieu M, Kirillova I, De Marco P, Merello E, Hayes JM, Wallingford JB, Drapeau P, Capra V, Gros P. Mutations in VANGL1 associated with neural-tube defects. *N Engl J Med.* 2007;356:1432–1437.
21. Kibar Z, Vogan KJ, Groulx N, Justice MJ, Underhill DA, Gros P. Ltap, a mammalian homolog of Drosophila Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat Genet.* 2001;28:251–255.
22. Montcouquiol M, Rachel RA, Lanford PJ, Copeland NG, Jenkins NA, Kelley MW. Identification of Vangl2 and Scrib1 as planar polarity genes in mammals. *Nature.* 2003;423:173–177.
23. Curtin JA, Quint E, Tspouri V, Arkell RM, Cattanach B, Copp AJ, Henderson DJ, Spurr N, Stanier P, Fisher EM, Nolan PM, Steel KP, Brown SD, Gray IC, Murdoch JN. Mutation of Celsr1 disrupts planar polarity of inner ear hair cells and causes severe neural tube defects in the mouse. *Curr Biol.* 2003;13:1129–1133.
24. Das G, Jenny A, Klein TJ, Eaton S, Mlodzik M. Diego interacts with Prickle and Strabismus/Van Gogh to localize planar cell polarity complexes. *Development.* 2004;131:4467–4476.
25. Lu X, Borchers AG, Jolicoeur C, Rayburn H, Baker JC, Tessier-Lavigne M. PTK7/CCK-4 is a novel regulator of planar cell polarity in vertebrates. *Nature.* 2004;430:93–98.
26. Ross AJ, May-Simera H, Eichers ER, Kai M, Hill J, Jagger DJ, Leitch CC, Chapple JP, Munro PM, Fisher S, Tan PL, Phillips HM, Leroux MR, Henderson DJ, Murdoch JN, Copp AJ, Eliot MM, Lupski JR, Kemp DT, Dollfus H, Tada M, Katsanis N, Forge A, Beales PL. Disruption of Bardet-Biedl syndrome ciliary proteins perturbs planar cell polarity in vertebrates. *Nat Genet.* 2005;37:1135–1140.
27. Slusarski DC, Corces VG, Moon RT. Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature.* 1997;390:410–413.
28. Slusarski DC, Yang-Snyder J, Busa WB, Moon RT. Modulation of embryonic intracellular Ca<sup>2+</sup> signaling by Wnt-5A. *Dev Biol.* 1997;182:114–120.
29. Pandur P, Lasche M, Eisenberg LM, Kuhl M. Wnt-11 activation of a non-canonical Wnt signalling pathway is required for cardiogenesis. *Nature.* 2002;418:636–641.
30. Eisenberg CA, Gourdie RG, Eisenberg LM. Wnt-11 is expressed in early avian mesoderm and required for the differentiation of the quail mesoderm cell line QCE-6. *Development.* 1997;124:525–536.
31. Eisenberg CA, Eisenberg LM. WNT11 promotes cardiac tissue formation of early mesoderm. *Dev Dyn.* 1999;216:45–58.
32. Koyanagi M, Haendeler J, Badorf C, Brandes RP, Hoffmann J, Pandur P, Zeiher AM, Kuhl M, Dimmeler S. Non-canonical Wnt signaling enhances differentiation of human circulating progenitor cells to cardiomyogenic cells. *J Biol Chem.* 2005;280:16838–16842.
33. Terami H, Hidaka K, Katsumata T, Iio A, Morisaki T. Wnt11 facilitates embryonic stem cell differentiation to Nkx2.5-positive cardiomyocytes. *Biochem Biophys Res Commun.* 2004;325:968–975.
34. Kremenevskaja N, von Wasielewski R, Rao AS, Schoff C, Andersson T, Brabant G. Wnt-5a has tumor suppressor activity in thyroid carcinoma. *Oncogene.* 2005;24:2144–2154.
35. Dejmeck J, Safholm A, Kamp Nielsen C, Andersson T, Leandersson K. Wnt-5a/Ca<sup>2+</sup>-induced NFAT activity is counteracted by Wnt-5a/Yes-Cdc42-casein kinase 1alpha signaling in human mammary epithelial cells. *Mol Cell Biol.* 2006;26:6024–6036.
36. Ma L, Wang HY. Suppression of cyclic GMP-dependent protein kinase is essential to the Wnt/cGMP/Ca<sup>2+</sup> pathway. *J Biol Chem.* 2006;281:30990–31001.
37. Saefholm A, Leandersson K, Dejmeck J, Nielsen CK, Villoutreix BO, Andersson T. A formylated hexapeptide ligand mimics the ability of Wnt-5a to impair migration of human breast epithelial cells. *J Biol Chem.* 2006;281:2740–2749.
38. Kuhl M, Sheldahl LC, Malbon CC, Moon RT. Ca(2+)/calmodulin-dependent protein kinase II is stimulated by Wnt and Frizzled homologs and promotes ventral cell fates in Xenopus. *J Biol Chem.* 2000;275:12701–12711.
39. Kuhl M, Sheldahl LC, Park M, Miller JR, Moon RT. The Wnt/Ca<sup>2+</sup> pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet.* 2000;16:279–283.
40. Koyanagi M, Iwasaki M, Haendeler J, Leitges M, Zeiher AM, Dimmeler S. Wnt5a increases cardiac gene expressions of cultured human circulating progenitor cells via a PKC delta activation. *PLoS One.* 2009;4:e5765.
41. Belema Bedada F, Technau A, Ebelt H, Schulze M, Braun T. Activation of myogenic differentiation pathways in adult bone marrow-derived stem cells. *Mol Cell Biol.* 2005;25:9509–9519.
42. Saneyoshi T, Kume S, Amasaki Y, Mikoshiba K. The Wnt/calcium pathway activates NF-AT and promotes ventral cell fate in Xenopus embryos. *Nature.* 2002;417:295–299.
43. Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signalling pathways. *Nat Rev Mol Cell Biol.* 2006;7:589–600.
44. Ishitani T, Kishida S, Hyodo-Miura J, Ueno N, Yasuda J, Waterman M, Shibuya H, Moon RT, Ninomiya-Tsuji J, Matsumoto K. The TAK1-NLK mitogen-activated protein kinase cascade functions in the Wnt-5a/Ca(2+) pathway to antagonize Wnt/beta-catenin signaling. *Mol Cell Biol.* 2003;23:131–139.
45. Ishitani T, Ninomiya-Tsuji J, Nagai S-i, Nishita M, Meneghini M, Barker N, Waterman M, Bowerman B, Clevers H, Shibuya H, Matsumoto K. The TAK1-NLK-MAPK-related pathway antagonizes signalling between beta-catenin and transcription factor TCF. *Nature.* 1999;399:798–802.
46. Karamboulas C, Swedan A, Ward C, Al-Madhoun AS, Wilton S, Boisvenue S, Ridgeway AG, Skerjanc IS. HDAC activity regulates entry of mesoderm cells into the cardiac muscle lineage. *J Cell Sci.* 2006;119:4305–4314.

47. Zhang T, Kohlhaas M, Backs J, Mishra S, Phillips W, Dybkova N, Chang S, Ling H, Bers DM, Maier LS, Olson EN, Brown JH. CaMKI-Delta isoforms differentially affect calcium handling but similarly regulate HDAC/MEF2 transcriptional responses. *J Biol Chem.* 2007; 282:35078–35087.
48. Backs J, Song K, Bezprozvannaya S, Chang S, Olson EN. CaM kinase II selectively signals to histone deacetylase 4 during cardiomyocyte hypertrophy. *J Clin Invest.* 2006;116:1853–1864.
49. Backs J, Backs T, Bezprozvannaya S, McKinsey TA, Olson EN. Histone deacetylase 5 acquires calcium/calmodulin-dependent kinase II responsiveness by oligomerization with histone deacetylase 4. *Mol Cell Biol.* 2008;28:3437–3445.
50. Little GH, Bai Y, Williams T, Poizat C. Nuclear calcium/calmodulin-dependent protein kinase IIdelta preferentially transmits signals to histone deacetylase 4 in cardiac cells. *J Biol Chem.* 2007;282: 7219–7231.
51. Passier R, Zeng H, Frey N, Naya FJ, Nicol RL, McKinsey TA, Overbeek P, Richardson JA, Grant SR, Olson EN. CaM kinase signaling induces cardiac hypertrophy and activates the MEF2 transcription factor in vivo. *J Clin Invest.* 2000;105:1395–1406.
52. Takada I, Mihara M, Suzawa M, Ohtake F, Kobayashi S, Igarashi M, Youn MY, Takeyama K, Nakamura T, Mezaki Y, Takezawa S, Yogiashi Y, Kitagawa H, Yamada G, Takada S, Minami Y, Shibuya H, Matsumoto K, Kato S. A histone lysine methyltransferase activated by non-canonical Wnt signalling suppresses PPAR-gamma transactivation. *Nat Cell Biol.* 2007;9:1273–1285.
53. Tu X, Joeng KS, Nakayama KI, Nakayama K, Rajagopal J, Carroll TJ, McMahon AP, Long F. Noncanonical Wnt signaling through G protein-linked PKCdelta activation promotes bone formation. *Dev Cell.* 2007; 12:113–127.
54. He X, Saint-Jeannet JP, Wang Y, Nathans J, Dawid I, Varmus H. A member of the Frizzled protein family mediating axis induction by Wnt-5A. *Science.* 1997;275:1652–1654.
55. Tao Q, Yokota C, Puck H, Kofron M, Birsoy B, Yan D, Asashima M, Wylie CC, Lin X, Heasman J. Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell.* 2005;120:857–871.
56. Cha SW, Tadjuidje E, Tao Q, Wylie C, Heasman J. Wnt5a and Wnt11 interact in a maternal Dkk1-regulated fashion to activate both canonical and non-canonical signaling in *Xenopus* axis formation. *Development.* 2008;135:3719–3729.
57. Cha SW, Tadjuidje E, White J, Wells J, Mayhew C, Wylie C, Heasman J. Wnt11/5a complex formation caused by tyrosine sulfation increases canonical signaling activity. *Curr Biol.* 2009;19:1573–1580.
58. Ye X, Wang Y, Cahill H, Yu M, Badea TC, Smallwood PM, Peachey NS, Nathans J. Norrin, frizzled-4, and Lrp5 signaling in endothelial cells controls a genetic program for retinal vascularization. *Cell.* 2009;139: 285–298.
59. Robitaille J, MacDonald ML, Kaykas A, Sheldahl LC, Zeisler J, Dube MP, Zhang LH, Singaraja RR, Guernsey DL, Zheng B, Siebert LF, Hoskin-Mott A, Trese MT, Pimstone SN, Shastry BS, Moon RT, Hayden MR, Goldberg YP, Samuels ME. Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy. *Nat Genet.* 2002;32:326–330.
60. Junge HJ, Yang S, Burton JB, Paes K, Shu X, French DM, Costa M, Rice DS, Ye W. TSPAN12 regulates retinal vascular development by promoting Norrin- but not Wnt-induced FZD4/beta-catenin signaling. *Cell.* 2009;139:299–311.
61. Carron C, Pascal A, Djiane A, Boucaut JC, Shi DL, Umbhauer M. Frizzled receptor dimerization is sufficient to activate the Wnt/beta-catenin pathway. *J Cell Sci.* 2003;116:2541–2550.
62. Kaykas A, Yang-Snyder J, Heroux M, Shah KV, Bouvier M, Moon RT. Mutant Frizzled 4 associated with vitreoretinopathy traps wild-type Frizzled in the endoplasmic reticulum by oligomerization. *Nat Cell Biol.* 2004;6:52–58.
63. Davidson G, Wu W, Shen J, Bilic J, Fenger U, Stanek P, Glinka A, Niehrs C. Casein kinase 1 gamma couples Wnt receptor activation to cytoplasmic signal transduction. *Nature.* 2005;438:867–872.
64. Zeng X, Tamai K, Doble B, Li S, Huang H, Habas R, Okamura H, Woodgett J, He X. A dual-kinase mechanism for Wnt co-receptor phosphorylation and activation. *Nature.* 2005;438:873–877.
65. Davidson G, Shen J, Huang YL, Su Y, Karaulanov E, Bartscherer K, Hassler C, Stanek P, Boutros M, Niehrs C. Cell cycle control of wnt receptor activation. *Dev Cell.* 2009;17:788–799.
66. Bahmanyar S, Kaplan DD, Deluca JG, Giddings TH Jr, O'Toole ET, Winey M, Salmon ED, Casey PJ, Nelson WJ, Barth AI. beta-Catenin is a Nek2 substrate involved in centrosome separation. *Genes Dev.* 2008; 22:91–105.
67. Ai D, Fu X, Wang J, Lu MF, Chen L, Baldini A, Klein WH, Martin JF. Canonical Wnt signaling functions in second heart field to promote right ventricular growth. *Proc Natl Acad Sci U S A.* 2007;104:9319–9324.
68. Kwon C, Arnold J, Hsiao EC, Taketo MM, Conklin BR, Srivastava D. Canonical Wnt signaling is a positive regulator of mammalian cardiac progenitors. *Proc Natl Acad Sci U S A.* 2007;104:10894–10899.
69. Pan W, Choi SC, Wang H, Qin Y, Volpicelli-Daley L, Swan L, Lucast L, Khoo C, Zhang X, Li L, Abrams CS, Sokol SY, Wu D. Wnt3a-mediated formation of phosphatidylinositol 4,5-bisphosphate regulates LRP6 phosphorylation. *Science.* 2008;321:1350–1353.
70. Chen M, Philipp M, Wang J, Premont RT, Garrison TR, Caron MG, Lefkowitz RJ, Chen W. G Protein-coupled receptor kinases phosphorylate LRP6 in the Wnt pathway. *J Biol Chem.* 2009;284:35040–35048.
71. Bryja V, Andersson ER, Schambony A, Esner M, Bryjova L, Biris KK, Hall AC, Kraft B, Cajanek L, Yamaguchi TP, Buckingham M, Arenas E. The extracellular domain of Lrp5/6 inhibits noncanonical Wnt signaling in vivo. *Mol Biol Cell.* 2009;20:924–936.
72. Tahinci E, Thorne CA, Franklin JL, Salic A, Christian KM, Lee LA, Coffey RJ, Lee E. Lrp6 is required for convergent extension during *Xenopus* gastrulation. *Development.* 2007;134:4095–4106.
73. Simons M, Gault WJ, Gotthardt D, Rohatgi R, Klein TJ, Shao Y, Lee HJ, Wu AL, Fang Y, Satlin LM, Dow JT, Chen J, Zheng J, Boutros M, Mlodzik M. Electrochemical cues regulate assembly of the Frizzled/Dishevelled complex at the plasma membrane during planar epithelial polarization. *Nat Cell Biol.* 2009;11:286–294.
74. Cruciat CM, Ohkawara B, Acebron SP, Karaulanov E, Reinhard C, Ingelfinger D, Boutros M, Niehrs C. Requirement of prorenin receptor and vacuolar H<sup>+</sup>-ATPase-mediated acidification for Wnt signaling. *Science.* 2007;317:459–463.
75. Penzo-Mendez A, Umbhauer M, Djiane A, Boucaut JC, Riou JF. Activation of Gbetagamma signaling downstream of Wnt-11/Xfz7 regulates Cdc42 activity during *Xenopus* gastrulation. *Dev Biol.* 2003;257: 302–314.
76. Liu X, Liu T, Slusarski DC, Yang-Snyder J, Malbon CC, Moon RT, Wang H. Activation of a frizzled-2/beta-adrenergic receptor chimera promotes Wnt signaling and differentiation of mouse F9 teratocarcinoma cells via Galphao and Galphat. *Proc Natl Acad Sci U S A.* 1999;96:14383–14388.
77. Liu T, Liu X, Wang H, Moon RT, Malbon CC. Activation of rat frizzled-1 promotes Wnt signaling and differentiation of mouse F9 teratocarcinoma cells via pathways that require Galpha(q) and Galpha(o) function. *J Biol Chem.* 1999;274:33539–33544.
78. Stemmler LN, Fields TA, Casey PJ. The regulator of G protein signaling domain of axin selectively interacts with Galpha12 but not Galpha13. *Mol Pharmacol.* 2006;70:1461–1468.
79. Liu T, DeCostanzo AJ, Liu X, Wang H, Hallagan S, Moon RT, Malbon CC. G protein signaling from activated rat frizzled-1 to the beta-catenin-Lef-Tcf pathway. *Science.* 2001;292:1718–1722.
80. Liu X, Rubin JS, Kimmel AR. Rapid, Wnt-induced changes in GSK3beta associations that regulate beta-catenin stabilization are mediated by Galpha proteins. *Curr Biol.* 2005;15:1989–1997.
81. Angers S, Thorpe CJ, Biechele TL, Goldenberg SJ, Zheng N, MacCoss MJ, Moon RT. The KLHL12-Cullin-3 ubiquitin ligase negatively regulates the Wnt-beta-catenin pathway by targeting Dishevelled for degradation. *Nat Cell Biol.* 2006;8:348–357.
82. Katanaev VL, Ponzicelli R, Semeriva M, Tomlinson A. Trimeric G protein-dependent frizzled signaling in *Drosophila*. *Cell.* 2005;120: 111–122.
83. Niehrs C. Function and biological roles of the Dickkopf family of Wnt modulators. *Oncogene.* 2006;25:7469–7481.
84. Bafico A, Liu G, Yaniv A, Gazit A, Aaronson SA. Novel mechanism of Wnt signalling inhibition mediated by Dickkopf-1 interaction with LRP6/Arrow. *Nat Cell Biol.* 2001;3:683–686.
85. Brott BK, Sokol SY. Regulation of Wnt/LRP signaling by distinct domains of Dickkopf proteins. *Mol Cell Biol.* 2002;22:6100–6110.
86. Semenov MV, Tamai K, Brott BK, Kuhl M, Sokol S, He X. Head inducer Dickkopf-1 is a ligand for Wnt coreceptor LRP6. *Curr Biol.* 2001;11:951–961.
87. Mao B, Wu W, Davidson G, Marhold J, Li M, Mechler BM, Delius H, Hoppe D, Stanek P, Walter C, Glinka A, Niehrs C. Kremen proteins are



- Dickkopf receptors that regulate Wnt/beta-catenin signalling. *Nature*. 2002;417:664–667.
88. Hassler C, Cruciat CM, Huang YL, Kuriyama S, Mayor R, Niehrs C. Kremen is required for neural crest induction in *Xenopus* and promotes LRP6-mediated Wnt signaling. *Development*. 2007;134:4255–4263.
  89. Cselenyi CS, Lee E. Context-dependent activation or inhibition of Wnt-beta-catenin signaling by Kremen. *Sci Signal*. 2008;1:pe10.
  90. Lee AY, He B, You L, Xu Z, Mazieres J, Reguart N, Mikami I, Batra S, Jablons DM. Dickkopf-1 antagonizes Wnt signaling independent of beta-catenin in human mesothelioma. *Biochem Biophys Res Commun*. 2004;323:1246–1250.
  91. Canapero L, Huang YL, Staudt N, Masasumi T, Ahrendt R, Kazanskaya O, Niehrs C, Houart C. Dickkopf-1 regulates gastrulation movements by coordinated modulation of Wnt/beta-catenin and Wnt/PCP activities, through interaction with the Dally-like homolog Knypek. *Genes Dev*. 2007;21:465–480.
  92. Korol O, Gupta RW, Mercola M. A novel activity of the Dickkopf-1 amino terminal domain promotes axial and heart development independently of canonical Wnt inhibition. *Dev Biol*. 2008;324:131–138.
  93. Yamamoto H, Sakane H, Yamamoto H, Michiue T, Kikuchi A. Wnt3a and Dkk1 regulate distinct internalization pathways of LRP6 to tune the activation of beta-catenin signaling. *Dev Cell*. 2008;15:37–48.
  94. Schmitt AM, Shi J, Wolf AM, Lu CC, King LA, Zou Y. Wnt-Ryk signalling mediates medial-lateral retinotectal topographic mapping. *Nature*. 2006;439:31–37.
  95. Inoue T, Oz HS, Wiland D, Gharib S, Deshpande R, Hill RJ, Katz WS, Sternberg PW. *C. elegans* LIN-18 is a Ryk ortholog and functions in parallel to LIN-17/Frizzled in Wnt signaling. *Cell*. 2004;118:795–806.
  96. Mikels AJ, Nusse R. Purified Wnt5a protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol*. 2006;4:e115.
  97. Lu W, Yamamoto V, Ortega B, Baltimore D. Mammalian Ryk is a Wnt coreceptor required for stimulation of neurite outgrowth. *Cell*. 2004;119:97–108.
  98. Green JL, Inoue T, Sternberg PW. The *C. elegans* ROR receptor tyrosine kinase, CAM-1, non-autonomously inhibits the Wnt pathway. *Development*. 2007;134:4053–4062.
  99. Oishi I, Suzuki H, Onishi N, Takada R, Kani S, Ohkawara B, Koshida I, Suzuki K, Yamada G, Schwabe GC, Mundlos S, Shibuya H, Takada S, Minami Y. The receptor tyrosine kinase Ror2 is involved in non-canonical Wnt5a/JNK signalling pathway. *Genes Cells*. 2003;8:645–654.
  100. Schambony A, Wedlich D. Wnt-5A/Ror2 regulate expression of XPAPC through an alternative noncanonical signaling pathway. *Dev Cell*. 2007;12:779–792.
  101. Wouda RR, Bansraj MR, de Jong AW, Noordermeer JN, Fradkin LG. Src family kinases are required for WNT5 signaling through the Derailed/RYK receptor in the *Drosophila* embryonic central nervous system. *Development*. 2008;135:2277–2287.
  102. Billiard J, Way DS, Seestaller-Wehr LM, Moran RA, Mangine A, Bodine PV. The orphan receptor tyrosine kinase Ror2 modulates canonical Wnt signaling in osteoblastic cells. *Mol Endocrinol*. 2005;19:90–101.
  103. Al-Ata J, Paquet M, Teebi AS. Congenital heart disease in Robinow syndrome. *Am J Med Genet*. 1998;77:332–333.
  104. Kim KA, Zhao J, Andarmani S, Kakitani M, Oshima T, Binnerts ME, Abo A, Tomizuka K, Funk WD. R-Spondin proteins: a novel link to beta-catenin activation. *Cell Cycle*. 2006;5:23–26.
  105. Binnerts ME, Kim KA, Bright JM, Patel SM, Tran K, Zhou M, Leung JM, Liu Y, Lomas WE III, Dixon M, Hazell SA, Wagle M, Nie WS, Tomasevic N, Williams J, Zhan X, Levy MD, Funk WD, Abo A. R-Spondin1 regulates Wnt signaling by inhibiting internalization of LRP6. *Proc Natl Acad Sci U S A*. 2007;104:14700–14705.
  106. Kim KA, Wagle M, Tran K, Zhan X, Dixon MA, Liu S, Gros D, Korver W, Yonkovich S, Tomasevic N, Binnerts M, Abo A. R-Spondin family members regulate the Wnt pathway by a common mechanism. *Mol Biol Cell*. 2008;19:2588–2596.
  107. Kazanskaya O, Ohkawara B, Heroult M, Wu W, Maltry N, Augustin HG, Niehrs C. The Wnt signaling regulator R-spondin 3 promotes angioblast and vascular development. *Development*. 2008;135:3655–3664.
  108. Xu Q, Wang Y, Dabdoub A, Smallwood PM, Williams J, Woods C, Kelley MW, Jiang L, Tasman W, Zhang K, Nathans J. Vascular development in the retina and inner ear: control by Norrin and Frizzled-4, a high-affinity ligand-receptor pair. *Cell*. 2004;116:883–895.
  109. Yamamoto S, Nishimura O, Misaki K, Nishita M, Minami Y, Yonemura S, Tarui H, Sasaki H. Cthrc1 selectively activates the planar cell polarity pathway of Wnt signaling by stabilizing the Wnt-receptor complex. *Dev Cell*. 2008;15:23–36.
  110. Bovolenta P, Esteve P, Ruiz JM, Cisneros E, Lopez-Rios J. Beyond Wnt inhibition: new functions of secreted Frizzled-related proteins in development and disease. *J Cell Sci*. 2008;121:737–746.
  111. Lin X, Perrimon N. Dally cooperates with *Drosophila* Frizzled 2 to transduce Wingless signalling. *Nature*. 1999;400:281–284.
  112. Baeg GH, Lin X, Khare N, Baumgartner S, Perrimon N. Heparan sulfate proteoglycans are critical for the organization of the extracellular distribution of Wingless. *Development*. 2001;128:87–94.
  113. Fagotto F, Gluck U, Gumbiner BM. Nuclear localization signal-independent and importin/karyopherin-independent nuclear import of beta-catenin. *Curr Biol*. 1998;8:181–190.
  114. Wu X, Tu X, Joeng KS, Hilton MJ, Williams DA, Long F. Rac1 activation controls nuclear localization of beta-catenin during canonical Wnt signaling. *Cell*. 2008;133:340–353.
  115. Henderson BR, Fagotto F. The ins and outs of APC and beta-catenin nuclear transport. *EMBO Rep*. 2002;3:834–839.
  116. Cong F, Varmus H. Nuclear-cytoplasmic shuttling of Axin regulates subcellular localization of beta-catenin. *Proc Natl Acad Sci U S A*. 2004;101:2882–2887.
  117. Hendriksen J, Fagotto F, van der Velde H, van Schie M, Noordermeer J, Fornerod M. RanBP3 enhances nuclear export of active (beta)-catenin independently of CRM1. *J Cell Biol*. 2005;171:785–797.
  118. Kriehoff E, Behrens J, Mayr B. Nucleo-cytoplasmic distribution of beta-catenin is regulated by retention. *J Cell Sci*. 2006;119:1453–1463.
  119. Willert K, Jones KA. Wnt signaling: is the party in the nucleus? *Genes Dev*. 2006;20:1394–1404.
  120. Kikuchi A, Kishida S, Yamamoto H. Regulation of Wnt signaling by protein-protein interaction and post-translational modifications. *Exp Mol Med*. 2006;38:1–10.
  121. Hoppler S, Kavanagh CL. Wnt signalling: variety at the core. *J Cell Sci*. 2007;120:385–393.
  122. Sheldahl LC, Slusarski DC, Pandur P, Miller JR, Kuhl M, Moon RT. Dishevelled activates Ca<sup>2+</sup> flux, PKC, and CamKII in vertebrate embryos. *J Cell Biol*. 2003;161:769–777.
  123. van Amerongen R, Nusse R. Towards an integrated view of Wnt signaling in development. *Development*. 2009;136:3205–3214.

# Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## An Updated Overview on Wnt Signaling Pathways: A Prelude for More Tata Purushothama Rao and Michael Kühl

*Circ Res.* 2010;106:1798-1806

doi: 10.1161/CIRCRESAHA.110.219840

*Circulation Research* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2010 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circres.ahajournals.org/content/106/12/1798>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation Research* is online at:  
<http://circres.ahajournals.org/subscriptions/>