

Biochemistry, Physiology, and Pathophysiology of NADPH Oxidases in the Cardiovascular System

Bernard Lassègue,* Alejandra San Martín,* Kathy K. Griendling

Abstract: The NADPH oxidase (Nox) enzymes are critical mediators of cardiovascular physiology and pathophysiology. These proteins are expressed in virtually all cardiovascular cells, and regulate such diverse functions as differentiation, proliferation, apoptosis, senescence, inflammatory responses and oxygen sensing. They target a number of important signaling molecules, including kinases, phosphatases, transcription factors, ion channels, and proteins that regulate the cytoskeleton. Nox enzymes have been implicated in many different cardiovascular pathologies: atherosclerosis, hypertension, cardiac hypertrophy and remodeling, angiogenesis and collateral formation, stroke, and heart failure. In this review, we discuss in detail the biochemistry of Nox enzymes expressed in the cardiovascular system (Nox1, 2, 4, and 5), their roles in cardiovascular cell biology, and their contributions to disease development. (*Circ Res.* 2012;110:1364-1390.)

Key Words: NADPH oxidases ■ vascular smooth muscle ■ endothelial cells ■ cardiomyocytes ■ atherosclerosis ■ hypertension ■ cardiac hypertrophy ■ cardiovascular disease ■ reactive oxygen species ■ redox

In the dozen years since the discovery that the neutrophil respiratory burst NADPH oxidase was only one member of a family of homologous enzymes, tremendous progress has been made in understanding the role of these proteins in the cardiovascular system. The NADPH oxidase (Nox) family is composed of 7 catalytic subunits termed Nox1-5 and Duox1 and Duox2 (for Dual Oxidase), regulatory subunits p22phox, p47phox, Noxo1, p67phox, Noxa1, p40phox, and the major binding partner Rac. Of these, the Nox1, 2, 4, and 5 enzymes are expressed in cardiovascular tissues and not only participate in normal vascular and cardiac function but also contribute to the development of cardiovascular disease.

Nox enzymes are heteroprotein complexes (except Nox5) with very specific regulatory mechanisms, tissue and subcellular patterns of expression, downstream targets, and functions. They are found in endothelial cells (ECs), vascular smooth muscle cells (VSMCs), macrophages, adventitial fibroblasts, cardiac myocytes and fibroblasts, as well as adipocytes and stem cells. These proteins work independently or in concert to control not only cell survival, growth and death but also cell-specific functions such as differentiation, angiogenesis, and contraction. When acutely or chronically upregulated, they have been implicated in hypertension, atherosclerosis, heart failure, ischemia reperfusion injury, and cardiac remodeling, but upregulation can be physiologically advantageous, as in angiogenesis and collateral formation. The molecular mechanisms that differentially control the

expression, activation and cellular pathways linked to Nox enzymes, as well as the physiological and pathophysiological contexts in which they exert their effects, are the focus of this review.

Biochemistry

Nox2

Historically, the first discovered member of the NADPH oxidase family,¹ also known as gp91phox, Nox2 is the catalytic subunit of an enzymatic complex responsible for the phagocyte respiratory burst. On activation, Nox2 uses NADPH to reduce molecular oxygen to superoxide anion, which, in concert with its metabolites, is used by phagocytes to destroy invading microorganisms. Nox2 is also found throughout the cardiovascular system (Table 1), where its expression and activity are much lower than in phagocytes,²⁻⁴ consistent with signaling functions described in a following section. However, in pathological circumstances, excess Nox2 can lead to oxidative stress and disease development (see Physiology and Pathophysiology In Vivo). We will first summarize salient features of Nox2, mostly derived from studies conducted in phagocytes, that apply to the cardiovascular enzyme. This enzyme will later serve as a reference against which to compare other NADPH oxidases.

The structure of Nox2 is not known in complete detail, but current models are based on homology to proteins of known architecture, mapped epitopes of monoclonal antibodies and

Original received December 22, 2011; revision received February 9, 2012; accepted March 9, 2012. In February 2012, the average time from submission to first decision for all original research papers submitted to *Circulation Research* was 13.77 days.

From the Division of Cardiology, Department of Medicine, Emory University, Atlanta, GA.

*B.L. and A.S.M. contributed equally.

Correspondence to Kathy K. Griendling, PhD, Emory University, Division of Cardiology, 319 WMB, 1639 Pierce Dr, Atlanta, GA 30322. E-mail kgriend@emory.edu

© 2012 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.111.243972

Non-standard Abbreviations and Acronyms

AGEs	advanced glycation end-products
Ang II	Angiotensin II
AP-1	activator protein 1
ASK1	apoptosis signal-regulating kinase 1
AT1R	Angiotensin II type 1 receptor
bFGF	basic fibroblast growth factor
BMP4	bone morphogenetic protein 4
CGD	Chronic granulomatous disease
CRP	C-reactive protein
Duox	dual oxidase
EGF	epidermal growth factor
ECs	endothelial cells
eNOS	endothelial nitric oxide synthase
EPCs	endothelial progenitor cells
ER	endoplasmic reticulum
ERK	extracellular signal-regulated kinase
ET-1	endothelin 1
G/GM-CSF	granulocyte/granulocyte macrophage colony-stimulating factor
GSK3	glycogen synthase kinase 3
Hsp	heat shock protein
IL	interleukin
iNOS	inducible nitric oxide synthase
IQGAP1	IQ motif-containing GTPase-activating protein 1
JAK	Janus kinase
JNK	c-Jun N-terminal kinase
LDLR	low-density lipoprotein receptor
LPS	lipopolysaccharide
MAPK	mitogen-activated protein kinase
MCP-1	monocyte chemotactic protein-1
MEF2C	myocyte enhancer factor-2C
MI	myocardial infarction
miRNA	microRNA
MMP	matrix metalloproteinase
NF-κB	nuclear factor kappa B
NO	nitric oxide
Nox	NADPH oxidase
Nox5S/L	Nox5 short/long
Noxa	Nox activator
Noxo	Nox organizer
oxLDL	oxidized low-density lipoprotein
Ox-PAPC	oxidized 1-palmitoyl-2-arachidonoyl-sn-glycerol-3-phosphocholine
PAI-1	plasminogen activator inhibitor-1
PDGF	platelet-derived growth factor
PDI	protein disulfide isomerase
PI3K	phosphoinositide 3-kinase
PKA/C	protein kinase A/C
Poldip2	polymerase delta-interacting protein 2
PPAR	peroxisome proliferator-activated receptor
PTP1B	protein tyrosine phosphatase 1B
PVN	paraventricular nucleus
ROS	reactive oxygen species
SAPK	stress-activated protein kinase

SERCA	sarco/endoplasmic reticulum calcium ATPase
SH2/3	Src homology 2/3
SHP-2	SH2 domain tyrosine phosphatase 2
SHR	spontaneously hypertensive rat
si/shRNA	small interfering/short hairpin RNA
SRF	serum response factor
STAT	signal-transducer and activator of transcription
TASK-1	TWIK-related acid-sensitive potassium channel
TGF	transforming growth factor
TLR	Toll-like receptor
TNF	tumor necrosis factor
TPR	tetratricopeptide repeat
VCAM	vascular cell adhesion molecule
VSMCs	vascular smooth muscle cells
WKY	Wistar-Kyoto rats

mutations from X-linked chronic granulomatous disease (CGD) patients or site-directed mutagenesis.⁵ Briefly, Nox2 is composed of two main domains of equal sizes with very different properties. The amino-terminal moiety includes six transmembrane α -helices (numbered I–VI) connected by 5 loops (numbered A through E, Figure 1). Because both amino- and carboxy-termini are cytosolic, 3 loops are extracellular and include consensus asparagine glycosylation sites, whereas the other 2 are intracellular and accessible to cytosolic regulators. The degree of Nox2 glycosylation and apparent molecular mass differ between species. In Western blots of human neutrophil proteins, Nox2 is detected as a smear from about 55 to 91 kDa. Transmembrane helices III and V each include 2 conserved histidine residues that coordinate iron atoms at the center of 2 heme molecules.⁶ The cytosolic carboxy-terminal moiety of Nox2 constitutes a dehydrogenase domain that includes consensus binding sites for its NADPH substrate and FAD cofactor. On activation, 2 electrons are transferred from cytosolic NADPH to FAD and in succession across the membrane, via redox changes in heme irons. Finally, each electron reduces a molecule of oxygen to a superoxide radical, which is subsequently released outside the cell or in a topologically equivalent compartment, such as a vesicle lumen. Superoxide is the first reactive oxygen species (ROS) in a cascade of metabolites including hydrogen peroxide and peroxynitrite.⁷ A charge compensation mechanism, required to balance electron transport by Nox2 and sustain its activity, is provided by a voltage-gated proton channel⁸ and the chloride/proton antiporter CIC-3.⁹

Tight regulation, critical to avoid excessive production of deleterious superoxide, is evident from the large number of proteins involved in oxidase assembly (Figure 1). These include Nox2 itself, p22phox, p47phox, p67phox, and p40phox, all essential subunits whose mutations can cause CGD.^{10,11} Also crucial is Rac GTPase, which binds p67phox and the dehydrogenase domain of Nox2.¹² In the resting state, Nox2 and p22phox form an inactive membrane complex known as cytochrome b558,^{1,5} whereas the other dormant

Table 1. Selected Examples of Characterization and Interventions Affecting Nox Subunits in Cardiovascular Cells

	Cell/Tissue Distribution	Upregulators	Transcription Factors	Subcellular Localization	Genetic Models	Inhibitors	
Nox1	ECs	Ang II	AP-1 ³⁶¹	Plasma membrane	Overexpressor ²⁸⁴	Apocynin	
	VSMCs	PDGF	ATF-1 ³⁶²	Caveolae ¹⁰⁵	Knockout ^{170,282}	DPI	
	Adventitial fibroblasts	PGF _{2α} , LDL ⁴	MEF2B ³⁶³	Endosomes ¹⁰⁶		GK-136901	
	Whole vessels ⁴	TNF-α ¹⁹⁶	NFκB ³⁶⁴	Perinuclear ⁷⁴		ML171	
	Cardiomyocytes ³⁵⁶	Oscillatory shear stress	STAT1/3 ³⁵⁹				Nox2ds-tat ³⁶⁵
		BMP4 ³⁵⁷					
Aldosterone + salt ³⁵⁸							
IFN-γ ³⁵⁹							
ET-1 ³⁶⁰							
	T3 ¹⁷⁷						
	Urokinase ¹⁷⁸						
	Oxidized LDL ¹⁸⁰						
	Vascular injury ⁴						
Nox2	ECs	Ang II, ET-1, TGF-β, IFN-γ, oxidized LDL ⁴		Plasma membrane ³⁶⁹	Overexpressor ²⁹⁵ Knockout ³⁷¹	Apocynin	
	VSMCs			Perinuclear ^{2,74,370}		DPI	
	Adventitial fibroblasts	Oscillatory shear stress ^{232,257}		Nuclear pore ⁷⁵		Nox2ds-tat	
	Whole vessels ⁴	Aldosterone + salt ³⁶⁷				VAS2870 ³⁶⁵	
	Cardiomyocytes ³⁶⁶	Ischemia ³⁶⁸					
	Vascular injury ⁴						
Nox4	ECs	TGF-β ^{235,236,237}	E2F ³⁷³	ER ^{255,369}	Overexpressor ^{139,332}	DPI	
	VSMCs	Thromboxane A2	NFκB ³⁶⁴	Perinuclear ⁷⁴	Knockout ^{332,342,352,378}	GK-136901	
	Adventitial fibroblasts	TNF-α ^{145,372}	STAT1/3 ³⁵⁹	Nucleus ^{105,129,217,369}		Plumbagin	
	Whole vessels ⁴	IFN-γ ³⁵⁹	HIF-1α ³⁷⁴	Mitochondria ^{139,352,369}		VAS2870 ³⁶⁵	
	Cardiac fibroblasts ²³⁶	Urotensin ¹¹⁶	Nrf2 ^{375,376}	Focal adhesions ^{105,229}			
	Cardiomyocytes ³⁵²	Urokinase ¹⁷⁸	Oct-1 ³⁷⁶	Stress fibers ²²⁹			
	Heart ³⁶⁹	Oscillatory shear stress ²³²			Cytoplasm ³⁷⁷		
		Hypoxia ²³⁸					
Hyperoxia ²¹⁷							
	Vascular injury ⁴						
Nox5	ECs ^{153,372}	Ang II		Plasma membrane		DPI ³⁶⁵	
	VSMCs ^{146,154}	ET-1 ¹⁵³		Intracellular ^{155,162}			
	Cardiac fibroblasts ²³⁶	Thromboxane A2					
		TNF-α ³⁷²					
	Atherosclerosis ³²⁴						
p22phox	ECs	Ang II ^{4,379}	AP-1 ³⁷⁹	Plasma membrane	Overexpressor ²⁹⁴		
	VSMCs	Thrombin ³⁸⁰	STAT1/3 ³⁵⁹	Caveolae	Point mutation ^{121,167}		
	Adventitial fibroblasts	PDGF	NFκB ³⁸⁴	Focal adhesions ¹⁰⁵			
	Whole vessels ⁴	TGF-β ⁴		Perinuclear ^{2,370}			
	Cardiomyocytes ³⁶⁶	Urotensin ¹¹⁶			Nucleus ^{105,217}		
		TNF-α ^{4,145,379}			Nuclear pore ⁷⁵		
		IFN-γ ³⁸¹					
		LPS ³⁸²					
		Pulsatile shear stress ⁴					
		Stretch + Ang II ³⁸³					
	Aldosterone + salt ³⁶⁷						
	Vascular injury ⁴						

(Continued)

Table 1. Continued

	Cell/Tissue Distribution	Upregulators	Transcription Factors	Subcellular Localization	Genetic Models	Inhibitors
p47phox	ECs	Ang II ^{4,379}	HBP1 ⁶⁷	Perinuclear ³⁷⁰	Spontaneous null ³⁵	Apocynin
	VSMCs	Thrombin ⁴	Ets-1 ³⁸⁵	Nuclear pore ⁷⁵	Knockout ^{36,37}	Nox2ds-tat ³⁶⁵
	Adventitial fibroblasts	TNF- α ^{145,379}	STAT1/3 ³⁵⁹	Actin fibers ³²		
	Whole vessels ⁴	Aldosterone + salt ³⁶⁷				
	Cardiomyocytes ³⁶⁶					
p67phox	ECs	Ang II ^{4,379}	STAT1/3 ³⁵⁹	Perinuclear ³⁷⁰	Spontaneous mutant (requires confirmation) ⁵⁷	
	VSMCs	TNF- α ^{145,379}		Actin fibers ³²		
	Adventitial fibroblasts					
	Whole vessels ⁴					
	Cardiomyocytes ³⁶⁶					
p40phox	ECs	Ang II ²⁵	HBP1 ⁶⁷		Point mutation	
	VSMCs				Knockout ⁶⁵	
	Whole vessels ^{4,67}					
Nox1	ECs ⁸⁶				Spontaneous null ⁹¹	
	Whole vessels ⁸⁵					
Noxa1	ECs ^{86,145,386}	Atherosclerosis ⁸³			Knockout ⁹⁷	
	VSMCs ^{83,84}					
	Whole vessels ⁸⁵					

subunits form a trimer in the cytosol with p67phox linking p47phox and p40phox.¹³ Signaling cascades activate Rac, p47phox and p40phox, allowing translocation of cytosolic subunits to the membrane and association with the cytochrome to form an active enzyme.^{1,5} Phosphorylation of Nox2 by protein kinase C (PKC) also promotes activity and recruitment of cytosolic subunits.¹⁴ We will now summarize essential features of these regulatory subunits.

Numerous structural and functional studies revealed that p47phox is composed of an N-terminal PX domain, 2 central SH3 domains, an autoinhibitory region and a C-terminal proline-rich region that binds p67phox.⁵ At rest, intramolecular folding masks the PX domain and the SH3 tandem, which binds to the autoinhibitory region.¹⁵ In leukocytes and transfected cells, unfolding of p47phox is triggered by PKC phosphorylation of serine residues, including 303, 304, and 328 in the autoinhibitory region.^{16–20} Other kinases that can phosphorylate p47phox on serine and tyrosine may contribute to oxidase activation.^{21–23} Similar phosphorylation events occur in vascular cells^{24–26} as well as tyrosine phosphorylation by Src.^{27,28} Once exposed, the PX domain allows translocation of p47phox to the membrane by binding to phosphoinositide 3-kinase (PI3K) lipid products^{20,29} and actin fibers,^{30–32} whereas the SH3 domains interact with the cytosolic C-terminal proline-rich region of p22phox.^{33,34} Thus, PKC, PI3K, and Src can cause oxidase assembly by recruiting p47phox bound to p67phox and p40phox. Because p47phox is an essential component of the oxidase in vivo, many pathophysiological studies have taken advantage of a spontaneous null mutant,³⁵ as well as 2 independent knockout mouse lines.^{36,37} These models are useful, but it should be remembered that p47phox can also be part of the Nox1-based oxidase (see below).

Functionally, p67phox is indispensable for Nox2 activation, whereas p47phox can be omitted from cell-free as-

says.^{38,39} Results from many studies indicate that p67phox is composed of an N-terminal region including 4 tetratricopeptide repeat (TPR) motifs, followed by an activation domain, an SH3 domain, a PB1 domain, and finally a second SH3 domain at the C-terminus. The TPR motifs, important for Rac-GTP binding, are defective in many CGD mutants.^{10,40–42} Interestingly, in the cell-free system, a fragment of p67phox, including the N-terminus and the activation domain, fused with Rac1, is highly active presumably because it is targeted to the membrane.⁴³ Most importantly, the activation domain of p67phox triggers FAD reduction by Nox2.³⁹ The V204A mutant in this region is a competitive inhibitor of wild-type p67phox.⁴⁴ The first SH3 domain increases oxidase activity, but its target is not known.⁴⁵ The PB1 domain allows binding to p40phox, which is abolished by a K355A mutation.^{46,47} Finally, the C-terminal SH3 domain of p67phox is responsible for binding the proline-rich region of p47phox and therefore allows p67phox translocation to the membrane after activation.^{48–50} An additional degree of regulation is provided by phosphorylation, which is both constitutive and stimulated by agonists.^{51–53} Phosphorylation of p67phox, mediated by PKC δ , ERK2 and p38MAPK, increases superoxide production by Nox2.^{54–56} Physiological studies still await the creation of a p67phox knockout model. Although a line of mice with a mutation in the PB1 domain of p67phox and increased susceptibility to infections has been reported, additional work is required to verify that this mutation is the only cause of the phenotype.⁵⁷

Structural and functional studies indicate that p40phox is composed of an N-terminal PX domain,⁴⁰ a central SH3 domain and a C-terminal PB1 domain. Although masked by interaction with the PB1 domain at rest,⁵⁸ the PX domain sustains oxidase activation after stimulation by interaction with lipid products of PI3K.^{59–61} Because p40phox binds




Figure 1. Structure and activation of Nox enzymes. Catalytic Nox subunits, represented in blue, include an N-terminal domain composed of 6 transmembrane helices, numbered I through VI. Four histidine residues in helices III and V coordinate 2 heme iron atoms. A cytosolic C-terminal dehydrogenase domain includes an FAD cofactor and a NADPH substrate binding site. On activation, electrons are transferred from NADPH to FAD and across the membrane, via heme irons, to molecular oxygen, thus producing superoxide anion, which can be dismutated into hydrogen peroxide. **Upper panel.** Both Nox1 and Nox2 (blue) form a complex with p22phox (green), which includes 2 transmembrane domains and a C-terminal proline-rich region (PRR). A cytosolic complex (orange), is composed of an organizer (Noxo1 or p47phox), an activator (Noxa1 or p67phox), and p40phox (only with p67phox). The organizer, stimulated by phosphorylation (red dots) in the case of p47phox, binds the proline-rich region of p22phox and membrane lipids. Likewise, p40phox binds lipids in endosomal membranes. Rac, activated by GTP (brown dot), binds membrane, Nox and Nox activators which trigger FAD reduction. **Middle panel.** Nox4 (blue) also forms a complex with p22phox (green). Its activity, constitutive in the absence of cytosolic subunits, can be increased by binding of Poldip2 to the cytosolic C-terminal of p22phox. **Lower panel.** Whereas Nox5S is composed of a catalytic subunit similar to the other oxidases (blue), Nox5L includes an additional N-terminal segment (red) with 4 EF-hands. Binding of cytosolic calcium to the EF hands triggers Nox5L activation.

inositol headgroups with lower phosphate content than p47phox,⁵⁹ it preferentially interacts with endosomes. Thus, p40phox appears to increase oxidase activity in cooperation with p47phox^{47,62,63} not by inducing translocation to the membrane, but by retaining the oxidase at the phagosome.^{15,61,64} Depletion or mutation of p40phox impairs ROS production in neutrophils^{65,66} and ECs.⁶⁷ Whereas the PX domain is clearly essential to p40phox function, the target of the SH3 domain is unknown and its function is controversial.^{68,69} The C-terminal PB1 domain of p40phox interacts with the PB1 domain of p67phox, as shown using multiple approaches.^{13,47} In addition, phosphorylation of p40phox on serine and threonine by PKC is increased during stimulation,^{67,70,71} particularly at threonine 154, which is usually,⁶⁹ but not always,⁷² required for full oxidase activation. The significance of p40phox is emphasized by a single case of CGD known to date. In this patient a point mutation in the PX domain does not prevent extracellular superoxide production, but impairs oxidase activation following phagocytosis.⁷³

Because Nox2 produces superoxide, a highly reactive and short-lived ROS, its subcellular localization is important for function. Superoxide can be released outside the cell when Nox2 is located at the plasma membrane, thus allowing it, for example, to intercept endothelial-derived nitric oxide (NO) before it reaches the adjacent smooth muscle cell layer in vessels.⁷ However, Nox2 is also frequently found in intracellular compartments, around the nucleus and colocalized with the endoplasmic reticulum (ER) in vascular cells (Table 1).^{2,74} In ischemic cardiomyocytes, Nox2 is upregulated in the cytosol and targeted to the nuclear pore complex.⁷⁵ Production inside cells is consistent with a direct signaling role of superoxide or its metabolites, as will be seen later.

Nox1

Nox1 is most highly expressed in colon epithelium, where it is thought to play a role in host defense.⁷⁶ It is also found in many other tissues and in all layers of the vascular wall, particularly after induction by agonists and in pathological conditions (Table 1). Nox1, like Nox2, associates with p22phox to form a membrane-bound cytochrome.^{77–79} However, in colon the cytosolic subunits p47phox and p67phox are not expressed and are replaced by Noxo1 and Noxa1.^{80–82} We now understand that Nox organizers include both p47phox and its homologue, Noxo1, whereas Nox activators comprise p67phox and the structurally similar Noxa1. Because Nox2 and Nox1 are closely related, both enzymes can be activated in transfected cells by various organizer and activator pairs.⁸⁰ Examples are also known in vivo, as in VSMCs from mouse conduit arteries, where p47phox and Noxa1 activate Nox1.^{83,84} To understand the significance of these substitutions, a brief description of Noxo1 and Noxa1 is required.

Similar to p47phox, Noxo1 facilitates oxidase assembly by binding both an activator subunit and p22phox. The proline-rich region of Noxo1 binds to an SH3 domain of the activator, whereas the tandem SH3 domains of Noxo1 bind to the proline-rich region of p22phox.^{80,87,88} Noxo1 also binds to the dehydrogenase domain of Nox1.⁸⁹ Furthermore, the PX domain of Noxo1 provides an essential affinity for membrane

phosphoinositides.⁹⁰ However, unlike p47phox, because Nox1 lacks an autoinhibitory domain, it is thought to constitutively bind the cytochrome.^{80,81,90} Surprisingly, in spite of coexpression with Nox1 in colon and some vascular cells (Table 1), mice lacking Nox1 apparently only present a deficiency in Nox3 activity during development of the inner ear,⁹¹ perhaps because other organizers may compensate for its loss in other tissues. Besides p47phox, other possible organizers include Tks4 and Tks5, two Src substrates with a PX domain and multiple SH3 domains capable of binding p22phox and Noxa1, but not p67phox.⁹²

Several important features of p67phox are conserved in Noxa1, allowing it to serve as an activator subunit. These include four Rac-binding TPR motifs, a Nox activation domain and an SH3 domain that interacts with the proline-rich region of an organizer subunit. However, the p40phox-binding PB1 domain is not well conserved and the SH3 domain in the middle of the molecule is missing.^{80,81} In contrast to its Nox1 partner, Noxa1 activity appears to be tightly regulated. Phosphorylation of serine 461 by protein kinase A (PKA), serine 282 by ERK1/2 or p38 MAPK, or serine 172 by PKA or PKC, decreases ROS production by Nox1.^{93–95} Phosphorylation of Noxa1 by PKA favors binding to 14–3–3 and dissociation from Nox1,⁹³ whereas other kinases appear to decrease Noxa1 affinity for Rac1 and Nox1.⁹⁵ In contrast, phosphorylation of Noxa1 by Src on tyrosine 110 increases Nox1 activity.⁹⁶ These regulatory mechanisms may be important in vessels that express Noxa1 (Table 1). The recent creation of a knockout model will likely allow future detailed *in vivo* studies of Noxa1 function.⁹⁷

In addition to cytosolic organizers and activators, Nox1 also requires Rac1 for activity, as might be expected from its resemblance to Nox2. Nox1 is stimulated by constitutively active Rac1 and inhibited by Rac1 knockdown.^{87,98} In addition to binding an activator subunit, as mentioned above, Rac1 also has affinity for the plasma membrane due to prenylation⁹⁹ and interacts directly with the carboxy terminus of Nox1, even in the absence of Noxa1.^{12,98,100} Another related small monomeric GTPase, Cdc42 cannot activate Nox1,⁹⁸ but could be a competitive inhibitor, as in the case of Nox2.¹⁰¹ Interestingly, β Pix, a Rac1 guanine nucleotide exchange factor, appears to be constitutively bound to Nox1 and essential for its activity.^{89,100} These results suggest that Rac1 provides a crucial mechanism for activation by agonists, particularly in cells that exclusively express Nox1/Noxa1/Noxa1.¹⁰²

Although the structure of Nox1 is not known, it is expected to be very close to other Nox enzymes (Figure 1),^{1,5} with a dehydrogenase domain binding FAD and NADPH, as well as 4 conserved histidine residues in transmembrane helices III and V, thought to coordinate 2 heme molecules. Similar to Nox2, mutation of these histidines abolishes binding to p22phox.⁷⁸ Family resemblance is further substantiated by the creation of a functional chimera comprised of the transmembrane domain of Nox1 and the cytosolic moiety of Nox4 (abbreviated as Nox1-Nox4 below).^{103,104} However, the signal peptide at the N-terminus of Nox1 is important for localization at the plasma membrane, as it is prevented by replacement with the N-terminus of Nox4.¹⁰³ In VSMCs,

Nox1 can be found at the plasma membrane, in caveolae¹⁰⁵ and in endosomes (Table 1).¹⁰⁶ Interestingly, agonists appear to stimulate Nox1 in specific locations, thus determining where superoxide is produced: extracellularly by muscarinic agonists and thrombin; in endosomes by IL-1 β and TNF α ; both inside and outside cells by Ang II.^{106–108} Of importance, Nox1 stimulation in endosomes is dependent on CIC-3, where this ion exchanger is required to balance the electrogenic activity of the enzyme.^{106,109} Interestingly, Nox1 activity is also dependent on chaperones Hsp90¹⁰⁴ and PDI,^{110,111} which appear to be necessary not only for protein folding after synthesis, but also to maintain enzyme stability.

The primary biochemical function of vascular Nox1 is superoxide production, which is then rapidly converted to hydrogen peroxide. The moderate physiological activity of Nox1, compared with the phagocytic Nox2, can be attributed to its low expression as well as specific regulatory subunits and signaling cascades. However, in pathological conditions, upregulation of Nox1 can lead to oxidative stress in the cardiovascular system.

Nox4

Abundant in kidney cortex where it was originally discovered,^{112,113} Nox4 appears to be expressed almost ubiquitously, including throughout the cardiovascular system and in macrophages important in atherosclerosis.¹¹⁴ Nox4 is usually coexpressed with other homologues such as Nox1 and Nox2, but at significantly higher levels. Upregulation by prolonged exposure to agonists such as TGF- β appears to be a major route of Nox4 regulation (Table 1).

Similar to Nox1 and Nox2, coprecipitation and colocalization studies showed that Nox4 binds to p22phox (Figure 1).^{74,78,115} Furthermore, overexpression and siRNA experiments indicate that association with p22phox is required for Nox4 activity and that the 2 proteins stabilize each other.^{78,79,115–117} As in Nox1 and Nox2, mutation of heme-binding histidine 115 also abolishes Nox4 interaction with p22phox.⁷⁸ In contrast, transfection of organizer and activator subunits does not increase ROS production, suggesting that Nox4 activity is constitutive.^{78,79,115} This interpretation is supported by other observations in transfected cells: addition of cytosol to membrane fractions does not increase Nox4 activity¹¹⁸; mutation of the proline-rich domain of p22phox required for docking organizers does not affect Nox4 activity⁷⁹; Nox1-Nox4 and Nox2-Nox4 chimeras are active without transfection of cytosolic subunits,^{103,104,119} whereas the opposite Nox4-Nox2 chimera requires activation.^{117,119} However, ROS production is enhanced by the multifunctional Poldip2, which also interacts with p22phox,¹²⁰ presumably at the beginning of the cytosolic C-terminus, upstream of the region dispensable for Nox4 activity.¹²¹ Unlike Nox1 and Nox2, Rac1 does not activate Nox4 in transfected cells,^{115,122} in spite of evidence compatible with an indirect effect.^{123–126} A number of reports indicate that agonists can acutely increase Nox4 activity or signaling,^{122,123,127–134} but only a few instances of possible coupling mechanisms are known: very rapid protein upregulation^{134,135}; control of NADPH availability^{136,137}; or direct interaction of the TLR4 receptor with the oxidase.^{127,131,138} These results suggest that Nox4 can

respond to agonist stimulation in addition to its widely accepted role as a constitutively active enzyme mostly regulated by transcription.

In the absence of a complete model, structural information about Nox4 can be inferred from comparison of its primary sequence to other oxidases. Similar to Nox1 and Nox2, the dehydrogenase domain of Nox4 requires FAD and NADPH.¹¹⁹ A P437H mutation in the canonical NADPH binding motif, analogous to the Nox2 mutation of a CGD patient, abolishes activity,^{117,139} and deletion of the NADPH binding domain produces a dominant-negative Nox4.^{128,140} Similar to Nox2, Nox4 is inhibited by an R96E mutation in the cytosolic B loop, a region of the amino-terminal domain that interacts with the NADPH binding site.^{117,141} Despite their sequence homology, Nox4 can produce a higher hydrogen peroxide to superoxide ratio than Nox1 and Nox2.^{104,115,117,118,142} However, the structural determinants allowing hydrogen peroxide production by Nox4 are not well understood. The Nox4 carboxy-terminal dehydrogenase domain would seem responsible, considering that Nox1-Nox4 or Nox2-Nox4 chimeras produce only peroxide in some studies.^{104,119} However, in other reports, the Nox1-Nox4 chimera generates superoxide,¹⁰³ and Nox4-Nox2 peroxide.^{117,119} In addition, replacing the first transmembrane domain of Nox4 by that of Nox1,¹⁰³ or altering the last extracellular loop of Nox4, makes it produce superoxide, rather than peroxide.¹⁴³ Additional studies will be required to resolve these discrepancies and determine if Nox4 can catalyze superoxide dismutation (presumably via the extracellular region), or even directly produce hydrogen peroxide (possibly at the C-terminus).

Besides the type of ROS produced by Nox4 and equally important for signaling is the subcellular location in which they are released. The distribution of Nox4 is surprisingly wide, including endoplasmic reticulum, plasma membrane, nucleus and mitochondria (Table 1), perhaps because its location varies according to cell type. It is also possible that each antibody preferentially labels the enzyme in a specific compartment.¹⁴⁴ Alternatively, different Nox4 isoforms may be present in specific subcellular locations. Five splice variants, named Nox4A through E, are found in lung epithelial cells. The most intriguing isoform, also reported in ECs,¹⁴⁵ is Nox4D because it appears to be fully active in a DCFH-DA assay, although it lacks most of the transmembrane domain.¹⁴⁰ One could speculate, by analogy with dehydrogenase domain assays,¹¹⁹ that Nox4D retains activity by coupling to electron acceptors, such as cytochrome c in mitochondria, which might also be an alternative route of hydrogen peroxide formation by full-length Nox4. Regardless of its activity, because Nox4D is expected to be hydrophilic, it could be present in compartments devoid of membranes. However, the prevalence and functional significance of Nox4 isoforms remain to be explored.

Nox5

Compared with that of its homologues described above, Nox5 expression is restricted to fewer tissues. Although found in human VSMCs, ECs, and whole vessels (Table 1), Nox5 is unexpectedly absent in rodents.⁵ The most striking structural

difference with other Nox enzymes is the presence of an additional cytosolic N-terminal segment, containing 4 calcium binding EF-hands (Figure 1).^{146,147} However, this segment is missing in Nox5S, a short calcium-insensitive variant, which is the dominant isoform in carcinoma cells,¹⁴⁸ and expressed together with the long Nox5L in ECs.^{149,150} Nox5S may be constitutively active¹⁵⁰ or could be a competitive inhibitor of calcium-dependent activation when present in the same tetrameric complex as Nox5L.¹⁵¹ Except for the N-terminus of the long isoform, Nox5 is similar to other Nox enzymes, with 6 transmembrane helices expected to bind 2 hemes and a cytosolic dehydrogenase domain including FAD and NADPH binding sites.¹⁴⁶ However, neither Nox5 isoform appears to require cytosolic subunits or p22phox. Thus, Nox5 is active in a cell-free system without addition of cytosol¹⁴⁷ and is unaffected by expression or knockdown p22phox in transfected cells.^{79,150} Furthermore, Rac1 may participate in Nox5 activation in some systems,¹⁵² but not others.¹⁵³ On stimulation with agonists, such as PDGF, Ang II or endothelin-1 in VSMCs or ECs,^{153,154} or after addition of a calcium ionophore to transfected cells, an increase in cytosolic calcium concentration triggers high superoxide production by Nox5.^{146,155} Calcium induces binding of the N-terminal domain of Nox5 to the dehydrogenase domain, thus relieving autoinhibition.^{147,156} Moreover, 2 mechanisms may increase Nox5 sensitivity, allowing activation by resting calcium concentrations: (1) calcium-dependent binding of calmodulin to another site in the dehydrogenase domain¹⁵⁷ and (2) phosphorylation of serine and threonine residues by PKC and calcium/calmodulin-dependent kinase II.^{155,158,159} In addition, as Nox5 can be upregulated and activated by minute concentrations of hydrogen peroxide, ROS production may be sustained by a positive feed-back loop involving the tyrosine kinase c-Abl.^{160,161} Finally, binding of heat shock protein 90 to the C-terminus of Nox5 appears to stabilize the protein and enhance expression and activity.¹⁰⁴ Because Nox5 is found in intracellular compartments and at the plasma membrane, superoxide is expected to be produced both inside^{118,146} and outside¹⁵⁵ cells. Furthermore, a polybasic region in Nox5 with affinity for phosphoinositides may favor translocation to the plasma membrane and extracellular superoxide release.¹⁶²

p22phox

The foregoing sections highlight the catalytic subunits of NADPH oxidases and their cytosolic regulators. Because p22phox is an integral part of the Nox1, 2, and 4 complexes (as well as Nox3, which is not a vascular oxidase), it is worth a separate discussion. Among all NADPH oxidase subunits, p22phox is the most ubiquitous. It is highly expressed throughout the cardiovascular system, including heart and vessels and further upregulated by many stimuli (Table 1). Although the complete crystal structure of p22phox is not known, it is clearly an integral membrane protein and colocalizes with Nox proteins in numerous cell types and subcellular compartments. Current molecular models of p22phox indicate that two transmembrane helices are located in the middle of the molecule, leaving both N and C termini in the cytosol (Figure 1).^{163,164} The suggestion that histidine

94 in the first transmembrane region might coordinate a heme molecule in conjunction with a Nox subunit appears unlikely because this residue can be replaced without affecting the cytochrome.¹⁶⁵ How p22phox interacts with Nox catalytic subunits is still elusive. Maturation of Nox requires the presence of p22phox,^{121,164,166} and conversely, p22phox stability is increased by the presence of a Nox subunit.⁷⁸ Interestingly, a point mutation in p22phox at tyrosine 121, abolishes interaction with Nox2, but does not affect Nox4.^{117,121,167} Many studies have demonstrated that the proline-rich region in the C-terminal tail of p22phox is a docking site for organizer subunits^{80,87,88} and therefore required for activity of Nox1⁷⁹ and Nox2,¹⁶⁴ but not Nox4.¹²¹ Point mutations at residues 152, 156, or 158 in this region of p22phox abolish cytosolic subunit translocation and activity. Interaction of p22phox with p47phox also appears to require phosphorylation on threonine 147 in the cytosolic tail.¹⁶⁸ Thus, p22phox is not only required for stability of the Nox complex, but also serves to regulate interaction with cytosolic regulators.

Cell Physiology and Pathophysiology

At present, evidence suggests that under physiological conditions Nox proteins and their products superoxide and hydrogen peroxide participate in cardiovascular homeostasis by contributing to cell differentiation, repair of damaged tissue and vascular tone. They achieve all these functions through their roles as structural and signaling molecules. On the other hand, it is also well established that their dysregulation promotes and maintains pathological conditions, as outlined in the next section. Here, we will consider how NADPH oxidases signal distinct functional outcomes.

Nox1

Much effort has been directed toward understanding the specific roles of Nox1 in specific functions of cardiovascular cells. Animal studies show that neointimal formation following femoral artery injury as well as hypertensive remodeling in conduit arteries is significantly reduced in Nox1 knockout mice,^{169,170} suggesting a role of this oxidase in the proliferative and migratory functions of phenotypically modulated VSMCs. Other studies implicate Nox1 in vascular inflammation, as oxidase activity and Nox1 expression are increased in VSMCs exposed to advanced glycation end-products (AGEs),¹⁷¹ as well as in aortas of rats and mice with experimental diabetes.^{172,173} It should be noted that the involvement of Nox1 in these processes is not necessarily disease promoting: outward remodeling induced by high shear stress in a model of arteriovenous fistula, a positive adaptation, is probably mediated by Nox1,¹⁷⁴ and Nox1 has been implicated in angiogenesis, a necessary step in wound healing.¹⁷⁵

A role for Nox1 in proliferation has been established on many fronts (Figure 2). First, Nox1 is activated by agonists that stimulate growth in cell culture, such as Ang II,¹⁷⁶ thyroid hormone,¹⁷⁷ PDGF,¹⁷⁶ urokinase plasminogen activator,¹⁷⁸ prostaglandin F_{2α},¹⁷⁹ thrombin,⁸³ or oxLDL (Table 1).¹⁸⁰ Serum- and PDGF-induced proliferation are increased in VSMCs from Nox1 transgenic mice¹⁶⁹; conversely, target-




Figure 2. Physiological and pathophysiological responses mediated by Nox homologues in cardiovascular cells. This summary of the roles of Nox enzymes in different cell types of the cardiovascular system illustrates the paradox of their mediating both specialized and redundant functions. VSMC indicates vascular smooth muscle cells; EC, endothelial cells; FB, fibroblasts; CM, cardiomyocytes; MΦ, macrophages; and EPC, endothelial progenitor cells.

ing Nox1 with antisense or siRNA or genetic deletion of Nox1 in VSMCs inhibits proliferation induced by serum,¹⁸¹ thyroid hormone¹⁷⁷ and PDGF.¹⁶⁹ In an interesting study, Stanic et al¹⁸² showed that a more oxidized extracellular environment increases Nox1 expression and induces Nox1-dependent proliferation of VSMCs by activation of EGF-receptor signaling. Conversely, in human abdominal aortic ECs, endothelin-1, via the ETB1 receptor, attenuates NADPH oxidase activity by inhibiting the Pyk2-Rac1-Nox1 pathway and thus inhibits Ang II-induced proliferation.¹⁸³

The mechanisms by which Nox1 mediates proliferation are varied. In VSMCs stimulated with Ang II, p38MAPK, and Akt are downstream of Nox1 and are required for the hypertrophic response.^{184,185} Similarly, shRNA against the Nox1 activator Noxal inhibits the activation of Janus kinase 2, Akt, and p38MAPK by thrombin.⁸³ In actively cycling mouse lung epithelial cells, Nox1 stimulates cell proliferation

by reducing the requirement for growth factors to maintain expression of cyclin D1 and stimulates transcriptional activation of Fos family genes required for induction of cell cycle reentry.¹⁸⁶

However, Nox1 may have a dual role in the regulation of the balance between cell proliferation and death. Although in cancer cells downregulation of Nox1 increases caspase-3 activity and cell death,¹⁸⁷ necrosis has been positively linked to Nox1 in mouse fibroblasts exposed to TNF- α .¹⁸⁸ Moreover, Nox1, together with Nox4, appears to mediate arrest in S phase and senescence, a hallmark of atherosclerosis and cause of vascular dysfunction, in ECs exposed to resveratrol.¹⁸⁹

Nox1 has also been implicated in migration in different cell types. In VSMCs stimulated with thrombin, Nox1-derived ROS mediate migration in part through regulation of calcium influx via L-type Ca²⁺ channels.¹⁹⁰ Similarly, Nox1 mediates PDGF- and bFGF-induced migration.^{169,191} In the case of PDGF-induced migration, one role of Nox1 is to mediate the activation of the cofilin phosphatase, Slingshot,¹⁹² an activity shown to be necessary for migration after PDGF stimulation in VSMC.¹⁹³ When bFGF is the stimulus, Nox1 is required for activation of JNK and subsequent phosphorylation of the focal adhesion protein paxillin.¹⁹¹ Another mechanism by which Nox1 may affect migration is via regulation of matrix metalloproteinases. In fibroblasts, it has been demonstrated that Nox1-derived ROS induce the promoter activity of matrix metalloproteinase-9 by an NF κ B-mediated mechanism.¹⁹⁴ Finally, in ECs, Nox1 mediates migration through PPAR- α -dependent pathways.¹⁷⁵

A causal role for Nox1 in vascular inflammation has also been investigated. Exposure of VSMCs to thrombin increases IL-6 secretion, Nox1 expression, and NF κ B nuclear translocation.¹⁹⁵ These effects may be mediated by Nox1, as they were inhibited by atorvastatin, which is known to downregulate this oxidase.¹⁹⁵ Furthermore, stimulation of VSMCs with TNF α , IL-1 β , or AGEs activates NF κ B via Nox1 stimulation, which, at least in the case of AGEs, leads to upregulation of inducible NO synthase (iNOS).^{171,196} Nox1 is also expressed in macrophages and is required for foam cell formation,¹⁹⁷; thus, Nox1 mediates the vascular inflammatory response via several independent mechanisms.

Nox 2

Because of its expression pattern in the cardiovascular system, Nox2 functions mainly to regulate endothelial and cardiac function. It is upregulated and/or activated by a number of vasoactive stimuli, including Ang II, atrial natriuretic peptide, shear stress, activation of adenosine A(2A) receptors, deletion of α 1-AMP kinase, and nonselective nonsteroidal anti-inflammatory drugs (Table 1).^{122,198–202} Nox2 has been implicated in the control of vessel tone, inflammation, EC proliferation and migration (and therefore angiogenesis), as well as cardiac hypertrophy and remodeling.

Because superoxide produced in the endothelium can rapidly neutralize NO, Nox2 is expected to impair NO-mediated relaxation and lead to vessel contraction. Indeed, a negative correlation is observed between Nox2 expression

and endothelium-dependent relaxation in isolated aorta.^{203,204} In CGD patients who have a specific mutation in Nox2, flow-mediated vasodilation (which is largely dependent on NO) is significantly higher than in normal controls, corresponding with higher levels of serum nitrite and nitrate and suggesting that these patients have more bioactive NO.²⁰⁵ However, Nox2 also appears to affect vasodilatation via other mechanisms as well. For example, activation of Nox2 by high glucose leads to eNOS downregulation in human ECs.²⁰⁶ Conversely, activation of NO production can downregulate Nox2 expression,¹⁹⁸ suggesting tight regulation between these two important molecules.

Nox2 has also been closely linked to inflammation. Hwang et al²⁰⁷ showed that in ECs exposed to oscillatory shear stress (a stimulus known to promote inflammation and lesion formation), nox2 mRNA expression correlates with superoxide production and monocyte binding. This suggests that Nox2 activation may be proinflammatory, a conclusion supported by its role in macrophage activation,²⁰⁸ and by the observation that in VSMCs treated with IL-17, p38MAPK-mediated Nox2 activation is required for release of the proinflammatory cytokines IL-6, G-CSF, GM-CSF and MCP-1.²⁰⁹ Moreover, shear stress-dependent activation of p47phox-dependent NADPH oxidases, presumably Nox2 in ECs, causes eNOS uncoupling and xanthine/xanthine oxidase activation, one of many examples of ROS-induced ROS release.^{210,211}

In contrast to Nox1, the role of Nox2 in proliferation and cell cycle progression is more controversial. In human ECs, it has been demonstrated that alone or in coordination with Nox4, Nox2 contributes to cell proliferation by a mechanism that involves p38MAPK and Akt.¹²² Similarly, in adventitial fibroblasts, Nox2-ds-tat (formerly gp91phox-ds-tat), a peptide that prevents the interaction of Nox2 and p47phox, inhibits serum-induced proliferation.²¹² Along the same lines, it was suggested that Nox2 prevents apoptosis by inhibition of caspase 3/7 activity.²¹³ On the other hand, in human microvascular ECs, Nox2 mediates serum starvation-induced cell cycle arrest by a mechanism that targets the expression of p21^{cip1} and p53.²¹⁴ Similarly, Ang II/AT1R-induced senescence in endothelial progenitor cells (EPCs) was shown to require Nox2.²¹⁵ These disparate observations suggest that Nox2 regulation of the cell cycle is cell and context specific.

Like Nox1, Nox2 has been implicated in migration, but in this case almost exclusively in ECs and monocytes/macrophages.^{216–218} It also plays a role in EPC mobilization.²¹⁹ In ECs, Nox2 accumulation at the leading edge of migrating cells is dependent on an intact actin cytoskeleton and associates with both actin and the scaffold protein IQGAP1.²¹⁶ Depletion of IQGAP1 prevents Nox2 localization at the leading edge, reduces ROS production in migrating cells, and impairs migration. During EPC mobilization by erythropoietin, Nox2-derived ROS inactivate the protein-tyrosine phosphatase SHP-2 that activates STAT5.²¹⁹ In macrophages, Nox2 is required for LPS-induced, ERK1/2-mediated MMP-9, MMP-10, and MMP-12 expression and cell migration.²¹⁸

In the heart, Nox2 has been shown to participate in hypertrophy and remodeling after injury or aging.^{220,221} In-

deed, it has been demonstrated that after Ang II/AT1R activation in cardiomyocytes, Nox2-derived superoxide induces hypertrophy by a mechanism that requires Akt activation²²² and Wnt-dependent pathways.²²³ The profibrotic effects of Ang II are also mediated by Nox2. In rats overexpressing renin, septal wall thickness, and perivascular fibrosis are increased. These responses are attenuated when animals are treated with the mineralocorticoid receptor blocker spironolactone, in part due to reduction in Nox2-mediated ROS production.^{224,225} The increase in Nox2 that accompanies mineralocorticoid receptor activation depends on apoptosis signal-regulating kinase 1 (ASK1) activation by a mechanism yet to be determined.²²⁶

Nox 4

Nox4 is the most highly expressed Nox family member in all cells of the cardiovascular system, but its role is perhaps the most controversial. Its expression is marginally higher in arteries than in veins,²²⁷ and significantly higher in the cerebral than systemic vasculature.²²⁸ Of relevance, in different cell types, Nox4 is thought to regulate basal ROS production,^{229,230} suggesting that it may function to control the redox set-point and thus influence cellular metabolism.

Nox4 is upregulated by a wide variety of agonists and cellular stresses (Table 1). In ECs, Nox4 is sensitive to mechanical forces: it has been reported to be downregulated by cyclic strain or pulsatile flow and upregulated by oscillatory shear stress.^{231,232} The proinflammatory mediators TNF- α and oxidized 1-palmitoyl-2-arachidonoyl-sn-glycerol-3-phosphocholine (Ox-PAPC) increase the expression or activity of Nox4.^{125,145} In both endothelial and VSMCs, serum withdrawal also increases Nox4 expression.^{229,233} Although growth-promoting agonists and drugs such as Ang II, PDGF, interleukin-1 β , thrombin, and phorbol myristate acetate downregulate Nox4 in systemic VSMCs and adventitial fibroblasts,^{176,230,234} the proliferative hormone urotensin upregulates Nox4 in pulmonary artery SMCs.¹¹⁶ Only TGF- β has been consistently shown to upregulate Nox4 in all cell types tested and to activate it acutely.^{235–237} In fact, TGF- β is responsible for activation of Nox4 in response to other stimuli, such as hypoxia.²³⁸ Nox4 expression is also increased in pulmonary fibroblasts from patients with idiopathic pulmonary fibrosis and mediates TGF- β 1-induced fibroblast differentiation into myofibroblasts.²³⁹ The ability of Nox4 to respond to such diverse agonists suggests that it mediates many different cellular responses. Indeed, the variety of reported functions of Nox4 in the vasculature is truly remarkable. Nox4 has been linked to differentiation,^{229,236,240,241} migration,^{120,234,242} growth,^{74,178,235} apoptosis,²⁴² senescence,^{112,113} proinflammatory responses,^{125,131,243} and oxygen sensing.^{242,244,245}

Evidence linking Nox4 to differentiation is abundant. Correlative studies show that upregulation of Nox4 in the redifferentiating VSMCs in the neointima coincides with induction of SMC differentiation markers,²⁴⁶ whereas Nox4 in human atherosclerotic lesions correlates with smooth muscle α -actin expression.²⁴⁷ Moreover, electron microscopic analysis of SMCs in human atherosclerotic lesions indicate that only those VSMCs that maintain the contractile phenotype continue to express Nox4, whereas those that

become dedifferentiated lose Nox4.²⁴⁸ Treatment of vascular adventitial fibroblasts with TGF- β , which leads to differentiation to myofibroblasts, upregulates Nox4.²⁴⁹ Conversely, Nox4 expression is high in preadipocytes and goes down as these cells differentiate into adipocytes.²⁵⁰

A mechanistic link between Nox4 and differentiation was established in a number of studies using strategies to attenuate Nox4 expression or transfection to overexpress it. Nox4 depletion leads to a loss of differentiation marker gene expression in adult VSMCs and a reduction in the conversion of cardiac fibroblasts to myofibroblasts.^{229,236} Similar results were found in mouse embryonic stem cells, in which Nox4 overexpression increased SMC differentiation markers and Nox4 depletion decreased the ability of the cells to differentiate to VSMCs.²⁵¹ In both adult and embryonic cells, Nox4 appears to exert its effects by regulating SMC-specific transcription factors serum response factor (SRF) and/or myocardin, in some cases via a p38MAPK-dependent mechanism.^{229,237,251} Differentiation of mouse embryonic stem cells into cardiomyocytes has also been shown to require Nox4.²⁴¹ During differentiation into embryoid bodies, Nox4 expression initially decreases (day 1), and then gradually increases by day 8. Depletion of Nox4 using ribozymes reduces cardiac commitment (as assessed by a decrease in Nkx2.5 and MEF2C transcripts and a consequent reduction in ventricular myosin regulatory light chain 2) and dramatically reduces the percentage of spontaneously beating cells.²⁴¹

Although agonists such as TGF- β do not generally induce migration, clear evidence has accumulated implicating Nox4 in this biologically critical function. Inhibition of Nox4 using siRNA decreases PDGF-induced VSMC migration and wounding-activated EC migration.^{120,252} Paradoxically, overexpression of Nox4 also decreases migration in VSMCs stimulated with PDGF and in adventitial fibroblasts exposed to Ang II.^{120,234} One potential explanation for these apparently opposing observations lies in the fact that the Nox4/Poldip2 complex activates RhoA in SMCs, leading to focal adhesion turnover and stress fiber formation.¹²⁰ Thus, too little Nox4 prevents focal adhesion formation, whereas too much Nox4 prevents focal adhesion dissolution, both of which are required for cell motility. However, Nox4 can also inhibit motility by oxidizing the sarco-/ER Ca²⁺ ATPase (SERCA) and blocking the antimigratory action of NO, as was shown in VSMCs from Zucker obese rats.²⁵³ The opposite is true in ECs, where overexpression of Nox4 upregulates eNOS, leading to an enhanced migratory response.²⁵⁴

Nox4 has also been implicated in progression through the cell cycle. It has been shown that both Nox2 and Nox4 are required for basal ROS production and EC proliferation.⁷⁴ Moreover, knockdown of Nox4 impairs EGF- and HIV-tat induced EC proliferation, whereas overexpression of Nox4 enhances EC growth.^{252,255} Mechanistically, overexpression of Nox4 in HEK293 cells has been shown to activate ERK1/2 and JNK and to enhance insulin-induced phosphorylation of p38MAPK, Akt, and GSK3 β , any of which might contribute to proliferation.¹²² In pulmonary artery SMCs, urotensin-induced proliferation, TGF β -mediated hypoxia-induced proliferation,²³⁸ and TGF β -induced proliferation²³⁵ all require

Nox4. Nox4 also mediates hypoxia-induced proliferation in pulmonary artery adventitial fibroblasts.²⁵⁶ Growth-related signaling pathways downstream of Nox4 are incompletely understood, but may involve ERK1/2, p38MAPK, JNK, and Akt.^{116,213,235,252} In addition, Nox4 inhibits protein tyrosine phosphatase 1B (PTP1B) in the ER, leading to a constant activation of Akt and ultimately increased EC proliferation.^{128,254,255}

With regard to apoptosis, reports of the role of Nox4 are again mixed. TNF- α -induced apoptosis in microvascular ECs is dependent on Nox4,²⁵⁷ but apoptosis in response to serum deprivation is inhibited by Nox4 overexpression.²⁵² Knockdown of Nox4 inhibits BMP4-induced ROS production, phosphorylation of p38MAPK and JNK, and ultimately apoptosis in ECs.²⁶³ In the heart, transgenic overexpression of Nox4 leads to increased apoptosis by 13 to 14 months of age.¹³⁹ In adventitial fibroblasts, however, Nox4 inhibits hypoxia-induced apoptosis.²⁵⁶

Nox4 has also been implicated in senescence in a number of cell types and tissues,^{259,260} and this function of Nox4 may be an important contributor to cardiovascular diseases as well. Correlative data show that Nox4 expression increases with aging, particularly in the aorta, renal cortex and medulla.^{261–263} Functionally, Nox4 has been linked to senescence in ECs and polyploidy in VSMCs.²⁶³ In human umbilical vein ECs, knockdown of Nox4 reduces lifespan-associated increases in DNA damage and extends replicative lifespan independent of telomere shortening.²⁶⁴ Ang II-induced senescence of EPCs is also dependent on Nox4,²⁶⁵ and resveratrol, an antioxidant compound found in red wine, causes eventual endothelial senescence in a Nox4-dependent manner.¹⁸⁹ Of interest, overexpression of miR-146a, an miRNA whose expression increases with age, inhibits Nox4 protein expression and decreases senescence, although a causal link between these two endpoints was not established in this study.²⁶⁶ In VSMCs, overexpression of Nox4 increases polyploidy and decreases expression of the chromosome passenger protein, survivin, that has previously been implicated in polyploidization, suggesting that excess Nox4 impairs passage through the cell cycle.²⁶³ In the heart, NADPH oxidase inhibition normalizes age-dependent deceleration of shortening/relengthening of ventricular myocytes, although the responsible NADPH oxidase homologue has not been identified.²⁶⁷

Another potential role of Nox4 is in regulating proinflammatory/profibrotic gene expression in the vasculature. Urotensin-mediated induction of the profibrotic gene PAI-1 in pulmonary artery SMCs is dependent on Nox4, and downregulation of Nox4 blocks IL-8, MCP-1, LDLR, and ROS production in ECs stimulated with Ox-PAPC.^{116,125} Basal production of PAI-1 in human umbilical vein ECs is also dependent on Nox4 via p38MAPK and NF κ B.²⁶⁸ Another lipid agonist, palmitate, activates NF κ B in ECs via a TLR-4, Nox4-dependent mechanism.²⁶⁹ A similar relationship between Nox4 and inflammatory gene expression was found in cardiomyocytes, where both Nox2 and Nox4 siRNA significantly inhibit TNF- α -induced upregulation of IL-1 β and IL-6.²⁷⁰ Similarly, activation of the Fc γ RIIa by CRP leads to Nox4-dependent ROS production in VSMCs, and depletion of Nox4 attenuates AP-1 and NF κ B activation, as

well as MCP-1, IL-6, and ET-1 production.²⁷¹ These types of responses would support a proatherosclerotic response in intact vessels.

Finally, Nox4 has been proposed to be an oxygen sensor. The most convincing data in this regard were obtained in skeletal muscle, where Nox4-dependent ROS production is proportional to Po₂ and leads to oxidation of ryanodine receptor cysteine thiols to increase Ca²⁺ release in isolated sarcoplasmic reticulum.²⁴⁵ Another molecule proposed to be downstream of oxygen-dependent Nox4 activation is TASK-1, a pH-sensitive K⁺-channel. Hypoxia inhibits TASK-1 activity, a response abolished by Nox4 siRNA.²⁴⁴ Further studies are needed to determine if Nox4 has similar functions in the cardiovascular system.

The apparent ability of Nox4 to participate in such disparate cellular functions suggests that Nox4 may regulate fundamental cellular processes that contribute to each of these responses. Candidate pathways include, for example, the actin cytoskeleton, microtubules or cellular metabolism. Although some data exist linking Nox4 to the cytoskeleton, further investigation is needed to understand Nox4's fundamental role in cell biology.

Nox5

The role of Nox5 in the cardiovascular system has not been well studied, in part due to the lack of suitable animal models. Our knowledge of the potential functions of Nox5 is thus derived mainly from cultured cells and isolated tissue, as well as *Drosophila*, which express an ortholog of Nox5. In the latter system, proctolin, a neuropeptide that causes smooth muscle contraction to move eggs out of the ovary, utilizes Nox5 to stimulate calcium flux and subsequent contraction.²⁷² In cultured human VSMCs, Nox5 is required for PDGF-induced proliferation.¹⁵⁴ Knockdown of Nox5 using siRNA impairs the ability of PDGF to activate the JAK/STAT pathway, thus providing a potential mechanism for its growth-modulating effects. In microvascular ECs, overexpression of Nox5S or Nox5L leads to a modest increase in proliferation and the formation of capillary-like structures, whereas depletion of Nox5 partially reduces thrombin-stimulated growth and tube formation.¹⁵⁰ In these cells, Nox5 expression is increased by endothelin-1 and Ang II,¹⁵³ and mediates activation of ERK1/2, but not p38MAPK or SAPK/JNK.¹⁵³ In intact vessels, overexpression of Nox5 via adenovirus paradoxically increases eNOS activity, but as expected, reduces bioavailable NO via inactivation by superoxide.²⁷³ As a result, Nox5 overexpression in the endothelium impairs acetylcholine-induced endothelium-dependent relaxation and potentiates the contractile responses to phenylephrine.

Physiology and Pathophysiology In Vivo

Hypertension

Twenty years ago, the concept that ROS may play a role in hypertension was established by studies in which administration of heparin-bound SOD to spontaneously hypertensive rats reduced blood pressure.²⁷⁴ Numerous reports since then have verified these initial observations and implicate NADPH oxidases as important sources of ROS in blood pressure regulation. Many investigators found upregulation of NADPH oxidase

subunits in hypertensive animals and a reversal of blood pressure and vascular remodeling with agents such as the superoxide dismutase mimetic tempol or the putative NADPH oxidase inhibitor apocynin.^{275,276} Human studies are limited mostly to showing increased activity of NADPH oxidases in hypertensive subjects and association studies linking p22phox polymorphisms to hypertension.²⁷⁷ However, it was not until the development of animals with specific genetic modifications of Nox subunits that the function of each homologue in hypertension could be rigorously interrogated (Table 2).

The possibility that Nox1 might be implicated in hypertension is supported by descriptive data and correlative evidence. Indeed, in the kidney Nox1 is upregulated in various models of hypertension, including renin transgenic rats,²⁷⁸ 2-kidney 2-clip rats,²⁷⁹ and Dahl salt sensitive rats on a high salt diet.²⁸⁰ In rats infused with Ang II for 7 days, increased aortic Nox1 expression and hypertension are both blunted by administration of the PKC antagonist chelerythrine, which may also block PKC-dependent phosphorylation of p47phox.²⁸¹ Although a causal role for Nox1 in the acute response to hypertension is supported by a number of studies, its importance in chronic hypertension is controversial. Mice in which Nox1 is genetically deleted initially respond to Ang II with an increase in blood pressure, but sustained blood pressure elevation is reduced in these mice compared with their wild-type controls.^{170,282} Endothelium-dependent relaxation to acetylcholine is impaired in wild-type mice exposed to Ang II, but is preserved in Nox1 knockout mice, suggesting that it is Nox1-derived superoxide that inhibits endothelial function in Ang II-induced hypertension.²⁸² The effect of Nox1 deletion on vascular hypertrophy is less clear: it has been reported to have no effect after a moderate dose of Ang II²⁸² or to reduce hypertrophy by reducing matrix deposition after a high dose of Ang II.¹⁷⁰ This discrepancy may result from the extent of hypertension induced in these animals, as overexpression of Nox1 specifically in smooth muscle cells exacerbates the blood pressure response and enhances vascular hypertrophy to the lower dose of Ang II.²⁸³ Mechanistically, Nox1 overexpressing animals exhibit impaired endothelium-dependent relaxation that is a consequence of NO inactivation, implying cross-talk between cell types.²⁸⁴ In contrast to these results, when Nox1 knockout animals are crossed with animals that express human renin and so have a chronically activated renin-angiotensin system, no effect on blood pressure is observed even though ROS levels are reduced in plasma.²⁸⁵ These results are difficult to reconcile with the acute studies, although as in all work involving genetically modified animals, possible compensation must be considered. It should be noted, however, that deletion of Nox1 in this chronic model significantly reduced expression of the proinflammatory molecule VCAM-1.

Studies using recently designed Nox inhibitors or other therapeutic agents indirectly support the genetic studies implicating Nox1 in hypertension. Treatment of aged spontaneously hypertensive rats (which have increased Nox1 and Nox2 but not Nox4 compared with WKY animals) with VAS2870, a Nox1/Nox4 inhibitor, improved endothelium-dependent relaxation.²⁸⁶

Nox2 has also been studied extensively with regard to its potential role in blood pressure control, and is implicated in hypertension by virtue of its ability to inactivate NO and affect vessel wall remodeling, as well as its ability to influence central and renal control of blood pressure. It is upregulated in the vessel wall, the kidney and brain in many forms of hypertension.^{275,287–290} In Nox2 knockout mice, basal blood pressure is reduced, but the response to Ang II is intact.²⁹⁰ However, infusion of Nox2ds-tat, an inhibitor of Nox2 interaction with p47phox, or genetic deletion of p47phox reduces the hypertensive effect of angiotensin II.^{291,292} In Dahl salt-sensitive rats treated with Nox2ds-tat, ROS and endothelial-dependent relaxation are normalized, but blood pressure is not affected,²⁹³ whereas in mice with smooth muscle-specific overexpression of p22phox, the blood pressure response to Ang II is exacerbated.²⁹⁴ It should be noted that interfering with p47phox or p22phox likely affects other Nox homologues in addition to Nox2, but the predominance of Nox2 in resistance arteries, the site of blood pressure control, supports the concept that Nox2 may play a role in hypertension. More recent work has shown that endothelial-specific overexpression of Nox2 does not affect basal blood pressure, but exacerbates the hypertensive response to Ang II.²⁹⁵ Because this mimics the upregulation of Nox2 that occurs in hypertension, these observations suggest that vascular Nox2 may not be involved in the initiation of hypertension, but that once upregulated, it can exacerbate the disease.

Another potential contribution of Nox2 to hypertension lies in its ability to regulate central and renal function. Work from the Davisson group showed that intracellular superoxide production in the subfornical organ of the brain mediates the blood pressure response to Ang II.²⁹⁶ Experiments involving administration of small interfering RNAs to Nox2 or Nox4 to this region showed that both enzymes are required for the full vasopressor effects of Ang II, whereas only Nox2 regulates the dipsogenic response.²⁹⁷ Similarly, administration of p22phox antisense systemically abolishes the contribution of renal ROS production to the slow pressor response to Ang II.²⁹⁸ One mechanism that has been proposed to be the signal that integrates brain, vascular and renal control of blood pressure is the T-cell. Guzik et al²⁹⁹ showed that the blood pressure response to Ang II is reduced in mice lacking T- and B-cells (Rag1^{-/-} mice), and that adoptive transfer of T-cells lacking p47phox only partially restores Ang II-dependent hypertension, whereas wild-type T-cells completely reverse the response. Adoptive transfer of T-cells derived from either gp91phox^{-/-} or wild-type mice into Rag-1^{-/-} mice fully restores the prothrombotic phenotype induced by Ang II, suggesting that T-cell-derived ROS are not involved in the thrombotic response that accompanies hypertension.³⁰⁰

A role for Nox4 in hypertension is quite controversial. Nox4 has been reported to be elevated in the vasculature in several models of hypertension,^{278,301} but not in others.²⁸¹ This increase in Nox4 may be compensatory, however, since Ray et al³⁰² found that endothelial-specific Nox4 overexpression enhances agonist-mediated vasodilation by inducing hyperpolarization and decreases blood pressure by ≈ 10 mm Hg. In agreement with a potential provasodilatory

Table 2. Phenotypes of Mice in Which Nox Enzymes Are Genetically Modified

Genotype	Challenge	Phenotype	Reference
Nox1 ^{-/-}	Ang II	Acute increase of blood pressure and chronic reduction of blood pressure	170, 282
Nox1 ^{-/-}	Ang II	Preserved endothelium-dependent relaxation to acetylcholine	282
Nox1 ^{-/-}	High-dose Ang II	Reduction in hypertrophy	170
Nox1 overexpression in VSMC	Ang II	Increased blood pressure and increased hypertrophy	288
Nox1 overexpression in VSMC	Basal	Impaired endothelium-dependent relaxation	289
Nox1 ^{-/-}	30 min Brain ischemia	Unchanged neurological score, cerebral infarct volume and edema; higher cortical infarct volume	339
Nox1 ^{-/-}	1 h Brain ischemia	Improved neurological outcome, preservation of blood-brain barrier integrity and reduced cerebral edema	335
Nox2 ^{-/-}	Ischemia/reperfusion	Smaller cerebral infarct volume	336, 337
Nox2 ^{-/-}	Ischemia/reperfusion	Loss of protection with ischemic preconditioning in the heart and brain	338
Nox2 ^{-/-}	Basal	Reduction in blood pressure	290
Nox2 ^{-/-}	Ang II	No change in blood pressure	290
p47phox ^{-/-}	Ang II	Reduced hypertensive effect	291
p22phox overexpression in VSMC	Ang II	Exacerbated hypertensive response	294
Nox2 overexpression in EC	Ang II	Exacerbated hypertensive response	295
p47phox deletion in T cells	Ang II	Reduction in Ang II-dependent hypertension	299
Nox4 overexpression in EC	Acetylcholine and histamine	Enhanced agonist-mediated relaxation	302
Nox4 ^{-/-}	Basal	No effect on blood pressure	306
Nox1 ^{-/-}	Ang II	Protected from aortic dissection	307
Nox1 ^{-/-}	Wire injury	Attenuated neointimal response	169
Nox1 ^{-/-}	ApoE ^{-/-}	Reduced lesion area and macrophage infiltration in the ascending and thoracic aorta	308
Nox2 ^{-/-}	ApoE ^{-/-} and high-fat diet	No difference in lesion formation in the ascending aorta	314
Nox2 ^{-/-}	ApoE ^{-/-} and high-fat diet	Reduced atherosclerosis in the descending aorta	315
Nox2 ^{-/-}	Femoral artery ligation	Reduced neovascularization	328
Nox2 ^{-/-} plus wt bone marrow	Femoral artery ligation	Restoration of collateral formation	329
Nox2 ^{-/-}	Arterial injury	Reduced lesion formation	318
Nox2 ^{-/-}	Suppressor and pressor dose of Ang II	Attenuation of the hypertrophic response	344, 346
Nox2 ^{-/-}	Left coronary ligation	Less post-MI remodeling and unchanged infarct size	220, 348
Nox2 ^{-/-}	Aortic constriction	No difference in myocardial hypertrophy	349, 350
Nox2 ^{-/-}	Aortic constriction	Less interstitial hypertrophy and contractile dysfunction	351
Nox2 ^{-/-}	Hypercholesterolemia or diabetes	Protection against the loss of neovascularization	330, 331
p47phox ^{-/-}	ApoE ^{-/-}	Unchanged lesions in the ascending aorta	316
p47phox ^{-/-}	ApoE ^{-/-}	Reduced lesions in the descending aorta	322
p47phox ^{-/-}	ApoE ^{-/-}	No effect on lesion formation in the aortic sinus	321, 322
p47phox ^{-/-}	Myocardial infarction	Attenuation of cardiomyocyte hypertrophy, apoptosis, and interstitial fibrosis	352
Nox4 ^{-/-}	bFGF	Unchanged neovascularization	175
Nox4 overexpression in cardiomyocytes	Pressure overload	Impaired angiogenesis	332
Nox4 overexpression in EC	Hind limb ischemia	eNOS-dependent accelerated neovascularization	254
Nox4 ^{-/-}	Transient occlusion of middle cerebral artery	Less oxidative stress, blood-brain barrier leakage and neuronal apoptosis	342
Nox4 ^{-/-}	Aortic constriction	Greater cardiac hypertrophy, contractile dysfunction and ventricular dilatation	332
Nox4 overexpression in cardiomyocytes	Aortic constriction	Preservation of capillary density, protection against hypertrophy	332
Cardiac-specific Nox4 deletion	Aortic constriction	Reduced hypertrophy and contractile dysfunction	352

effect of Nox4, Paravicini et al³⁰³ showed that Nox4 expression in basilar arteries (which is 10-fold higher than in systemic arteries²²⁸ and is even greater in SHR) is associated with enhanced vasodilation in response to NADPH due to H₂O₂-mediated activation of BK(Ca) channels. In the brain, however, Nox4 in the subformal organ clearly plays a role in the pressor response to Ang II.²⁹⁷ Nox4 in the kidney may also mediate hypertension or the renal injury that accompanies hypertension, as it is upregulated in the macula densa and distal nephron in the SHR model, in the kidney of transgenic hypertensive rats overexpressing the Ren2 gene and in the cortex of rats treated with aldosterone.^{278,304,305} However, Nox4 is unlikely to be a major regulator of blood pressure, since knockout of Nox4 has no effect on basal blood pressure.³⁰⁶

Taken together, data from multiple genetically modified animal models confirm distinct roles for Nox homologues in hypertension. Depending on the cause of hypertension (genetic, diet-induced, hormone-regulated), its duration (acute versus chronic), the contributing organ (brain, kidney, vasculature), and the end point (blood pressure per se versus end organ damage), different homologues may play a predominant role. Translating these observations to human hypertension is obviously the next step but may have to await the development of safe, homologue-specific Nox inhibitors.

Atherosclerosis

Even more so than with hypertension, a significant body of literature implicates ROS in the development of atherosclerosis, at least in animal models. Although early work focused on oxidation of LDL, more recent studies tested the notion that NADPH oxidases in the vessel wall or in infiltrating macrophages and T-cells can affect atherogenesis on multiple levels. Conflicting reports in the literature leave the question of the relative importance of NADPH oxidases in this disease unsettled at this time.

Direct and indirect evidence for a role of Nox1 in atherogenesis is just now emerging. Gavazzi et al³⁰⁷ showed that Nox1 deficient mice infused with the proatherogenic peptide Ang II were protected from aortic dissection, potentially because they express higher levels of tissue inhibitor of metalloproteinase 1 than their wild-type controls, suggesting that Nox1 may regulate turnover of extracellular matrix. Mice in which Nox1 is deleted also exhibit an attenuated neointimal response to arterial injury, due in part to reduced VSMC proliferation and migration.¹⁶⁹ As all of these cellular processes contribute to atherosclerosis, one would predict that atherosclerotic lesions formation would be inhibited in the absence of Nox1. In fact, in the only study to directly assess the role of Nox1 in atherosclerosis, in which Nox1 knockout mice were crossed into the ApoE^{-/-} background and fed a high fat diet for 18 weeks, just such an outcome was observed.³⁰⁸ Compared with ApoE^{-/-} mice, lesion area in the ascending and thoracic aorta was reduced by 20% to 30% in Nox1^{-/-}ApoE^{-/-} mice, and macrophage content was reduced by about 50%. Following carotid injury in mice fed a chow diet, cell proliferation was decreased but collagen content was increased in the double knockout mice compared with ApoE^{-/-} controls. Moreover, Niu et al⁸³ showed that

adenovirus-mediated overexpression of the Nox1 activator Nox1 enhanced neointimal hyperplasia in a model of mouse carotid injury. It is worth noting that Nox1 is increased in atherosclerotic lesions in ApoE^{-/-} mice and in humans.⁸³ Additional evidence comes from a study showing that GKT136901, a newly-developed inhibitor of Nox1 and Nox4 oxidases, attenuates ROS generation and expression of the adhesion molecule CD44 and its principal ligand, hyaluronan, in atherosclerotic lesions.³⁰⁹ Finally, a study using the Janus kinase-2 inhibitor tyrphostin AG490 found that in ApoE^{-/-} mice fed a high fat diet, Nox1, 2, and 4 were elevated in the aorta, and that treatment with the inhibitor reduced expression and activity of all three homologues and reduced lesion area.³¹⁰

The contribution of Nox2 to atherosclerosis is controversial. Because mononuclear cells express high levels of Nox2 and macrophages are critical to lesion development, it was originally assumed that Nox2-derived ROS in macrophages would contribute to atherogenesis. This notion was supported by the observation that p22phox and Nox2 expression is high in atherosclerotic lesions, especially in the shoulder region of the plaque,^{247,311–313} and the findings that risk factors for atherosclerosis such as hypertension, oscillatory shear stress, and diabetes increase Nox2 expression in the vasculature.³¹² However, Kirk et al³¹⁴ called this assumption into question when they observed no difference in lesion formation in the ascending aortas of WT and Nox2 knockout mice fed a high fat diet or crossed with ApoE^{-/-} mice. In contrast, a more recent study found evidence for reduced atherosclerotic burden in the descending aortas of Nox2^{-/-}ApoE^{-/-} mice compared with ApoE^{-/-} mice fed a high fat diet.³¹⁵ These observations are similar to those in p47phox^{-/-}ApoE^{-/-} mice, in which lesion area in the ascending aorta is identical between p47phox^{-/-}ApoE^{-/-} mice and ApoE^{-/-} mice,³¹⁶ but lesions in the descending aorta are reduced.³¹⁷ Lesion formation is also reduced in Nox2^{-/-} mice subjected to arterial injury due to reduced cellular proliferation and reduced leukocyte accumulation.³¹⁸ Further experiments are necessary to define the contributions of cell-specific Nox2 to atherogenesis.

The mechanism for increased Nox expression and activity in atherosclerotic arteries is unclear. There is some evidence for ROS-dependent upregulation of NADPH oxidase subunits, suggesting a potential feed-forward mechanism that could exacerbate disease development.³¹⁹ Moreover, proatherogenic cytokines that activate NFκB, such as TNF-α, upregulate Nox2 and its cytosolic activators.³²⁰ Ox-PAPC has been shown to increase Nox4 activity in ECs.¹²⁵ Because the role of the various Nox isoforms in atherosclerosis is complex, their regulation is likely to be complicated as well, and measurements of different homologues in identical experimental conditions is needed.

Compared with Nox2 and Nox4, much less is known about the role of Nox4 in lesion formation because mice with genetically modified Nox4 alleles have only recently become available. As is the case for Nox2, risk factors such as diabetes^{173,321,322} and hypertension (see above) increase Nox4 expression in the vasculature. ApoE^{-/-} mice treated with a high fat diet also exhibit enhanced expression of Nox4,³¹⁰ and

Nox4 expression is high in human atherosclerotic lesions.²⁴⁷ Interestingly, treatment with statins, which has been shown to be beneficial in atherosclerosis, decreases Nox4 expression.^{322,323} Functionally, Nox4 has been implicated in neointimal formation after vascular injury. Adenoviral-mediated knockdown of Nox4 at the time of carotid injury in Zucker rats reduced the extent of oxidation of SERCA and inhibited the development of the neointima.²⁵³

Even less is known about the role of Nox5 in atherosclerosis, largely because it is not expressed in rodents, making genetic manipulation much more difficult. However, Guzik et al³²⁴ showed that Nox5 expression is higher in human coronary arteries that exhibit coronary artery disease than in those that are disease free. This increased expression was accompanied by a 7-fold increase in calcium-dependent NADPH oxidase activity. Nox5 was observed in ECs in early lesions and in VSMCs in advanced lesions. This is one of the few human studies linking any Nox homologue to atherosclerosis.

Angiogenesis and Collateral Formation

Growth of new blood vessels and expansion of existing ones is important not only during development, but can also occur in response to ischemic injury or during tumor development. Thus, depending on the context, these processes can either be beneficial or detrimental to the health of the organism. ROS both promote and inhibit blood vessel formation, in part depending on the amount of ROS produced. Yun et al³²⁵ proposed the “redox window” hypothesis, which states that although collateral formation requires ROS, either excess or insufficient ROS can inhibit vessel growth. A series of recent studies support a role for several of the Nox homologues in vessel formation and expansion.

Direct evidence for a role of Nox1 in angiogenesis comes from Garrido-Urbani et al,¹⁷⁵ who showed that Nox1 is upregulated by proangiogenic factors, that mice deficient in Nox1 exhibit impaired angiogenesis in the matrigel assay, and that tumor vascularization is inhibited either by genetic deletion of Nox1 in the host or by treatment with the Nox inhibitor GKT136901. They further demonstrated that Nox1 appears to promote tumor angiogenesis by inhibiting the antiangiogenic factor PPAR α . This supports earlier work by Arbiser et al,³²⁶ who found that Nox1-expressing NIH 3T3 cells injected into athymic mice cause highly vascularized, aggressive tumors, due to upregulation of VEGF and its receptors as well as increased MMP-9 activity. Komatsu et al³²⁷ found that Nox1 is necessary but not sufficient for induction of VEGF, indicating other factors are also involved.

Strong data exist both supporting and refuting a role for Nox2 in revascularization following ischemia. Ushio-Fukai's laboratory showed that Nox2 expression and superoxide production are increased in wild-type mice up to 7 days following femoral artery ligation, and that the neovascularization response is reduced in Nox2 knockout mice.³²⁸ Subsequent work suggests that it is Nox2 in bone marrow-derived cells that mediates the neovascularization response. Transplantation of wild-type bone marrow into Nox2-deficient mice restores blood flow recovery.³²⁹ In contrast, Nox2 knockout mice are protected against the loss of neovascular-

ization induced by hypercholesterolemia or diabetes, suggesting that the increase in Nox2 in response to these conditions is responsible for the observed impaired blood flow recovery in wild-type mice.^{330,331} Moreover, in the tumor angiogenesis matrigel plug model, loss of Nox2 has no effect on vasculogenesis.¹⁷⁵ Some of these apparent inconsistencies may relate to the level of ROS present basally in these models, per the redox window hypothesis, or to mechanistic differences between different types of vascular neogenesis.

The role of Nox4 in angiogenesis also seems to be context-specific. Subcutaneous implantation of a matrigel plug loaded with bFGF into Nox4^{-/-} mice leads to vascularization indistinguishable from that observed in wild-type mice.¹⁷⁵ However, the angiogenesis (defined as an increase in capillary density) that accompanies pressure overload-induced cardiac hypertrophy is impaired in Nox4 null mice and enhanced in cardiomyocyte-specific Nox4 overexpressing mice compared with wild-type controls.³³² This appears to be a consequence of Nox4-mediated activation of hypoxia inducible factor-1 and the release of vascular endothelial growth factor from cardiomyocytes, and is supported by *in vitro* studies in which overexpression of Nox4 enhanced, whereas dominant negative Nox4 or knockdown of Nox4 by siRNA impaired, tube formation in ECs.²⁵² Similarly, Craige et al²⁵⁴ showed that endothelial-specific overexpression of Nox4 leads to eNOS-dependent accelerated recovery from hind limb ischemia and enhanced aortic capillary sprouting. Finally, Bhandarkar et al³³³ found that formation of hemangiomas (endothelial-derived neoplasias) induced by grafting polyoma middle T-transformed brain ECs from angiopoietin 2 heterozygotes into nude mice is greatly reduced by lentiviral expression of siNox4. Thus, the bulk of the evidence suggests that Nox4 is proangiogenic, although its role in tumor formation and angiogenesis remains unclear.

Ischemia/Reperfusion in the Brain, Heart, and Lungs

Transient or sustained ischemia in the heart, brain or lung can lead to infarct, stroke, or pulmonary hypertension, respectively. NADPH oxidases have been implicated in the pathophysiology of all three diseases in animal models, with similar mechanisms.

In Nox1 knockout mice, the effects of ischemia have only been studied in the brain. Transient occlusion (30 minutes) of the middle cerebral artery results in similar neurological score, cerebral infarct volume and edema volume in wild-type and Nox1^{-/-} mice, but cortical infarct volume is much higher in brains of Nox1^{-/-} mice compared with wild-type mice after 24 hours.³³⁴ This suggests that Nox1 may actually limit infarct development after cerebral ischemia. However, in another study, a 1-hour but not a 2-hour occlusion of the middle cerebral artery in Nox1^{-/-} resulted in attenuation of lesion size at 24 hours after induction of ischemia, accompanied by a significant improvement of neurological outcome, preservation of blood-brain barrier integrity and reduced cerebral edema. Clearly, more studies are needed to understand the role of Nox1 in stroke.³³⁵

More is known about the role of Nox2 in ischemic injury. Using a similar experimental procedure to that described for Nox1, transient middle cerebral artery occlusion followed by

22 hours of reperfusion, 2 groups showed that male Nox2^{-/-} mice had a significantly smaller infarct volume compared with controls.^{336,337} Bone marrow transplants of Nox2^{-/-} cells into wild-type mice fail to induce this protection, and if wild-type cells are injected into Nox2^{-/-} mice, infarct volumes are not reduced, suggesting that both circulating Nox2 and neural or endothelial Nox2 contribute to ischemic injury in this model. Similarly, in a Langendorff-perfused, isolated mouse heart model of ischemia/reperfusion injury, ischemic preconditioning confers protection against necrotic injury in wild-type, but not Nox2^{-/-} hearts, indicating that in the heart as well as the brain, Nox2 is activated during either ischemia or reperfusion.³³⁸ Finally, administration of intermittent hypoxia to mice induces a phenotype of pulmonary hypertension, including elevated right ventricular systolic pressure, right ventricular hypertrophy, and increased muscularization of the distal pulmonary vessels.³³⁹ These changes are accompanied by an increase in p22phox and Nox4 expression but interestingly are attenuated in Nox2^{-/-} mice, suggesting that Nox2 may in fact regulate expression of other NADPH oxidases, and raise the possibility of a feed-forward mechanism that exacerbates pulmonary hypertension.

Emerging evidence strongly suggests a link between Nox4 and ischemic injury. In pulmonary hypertension induced by chronic intermittent hypoxia, Nox4 is upregulated as a consequence of Nox2 activation.³³⁹ Cortical Nox4 is upregulated in neurons early (24 hours) after middle cerebral artery occlusion, as well as 7 to 14 days later in newly formed capillaries in the peri-infarct region.³⁴⁰ Transient upregulation of Nox4 in the cortex is also observed after endothelin-induced stroke.³⁴¹ A functional role for Nox4 in ischemia-induced neurodegeneration was proposed based on the observation that compared with wild-type mice, Nox4 knockout mice have less oxidative stress, less blood-brain barrier leakage and less neuronal apoptosis after either transient occlusion of the middle cerebral artery or permanent stroke induced by cortical photothrombosis.³⁴² Importantly, post-stroke treatment with the putative Nox1/Nox4 inhibitor VAS2870 improves function, suggesting that Nox4 may be a viable therapeutic target for stroke.

Nox4 has also been implicated in cardiac performance following myocardial infarction (MI). Infanger et al³⁴³ found that surgical ligation of the left anterior descending coronary artery induces a 2-fold increase in Nox4 expression in the paraventricular nucleus (PVN) of the brain compared with sham animals, correlating with the increased sympathoexcitation and cardiac function that occurs following MI. Adenoviral-mediated silencing of Nox in the PVN improves both ejection fraction and fractional shortening two weeks post-MI, apparently by reducing sympathetic outflow. Surprisingly, these outcomes appeared to be a result of superoxide, rather than hydrogen peroxide, signaling in the PVN.

Cardiac Hypertrophy, Post-Myocardial Infarction Remodeling, and Heart Failure

As noted, Nox2 and Nox4 are the most highly expressed Nox homologues in the heart. Although both are important for normal physiology, they seem to regulate distinct aspects of cardiac function and thus have discrete roles in heart disease.

Early work by Ajay Shah's group implicated Nox2 in cardiac hypertrophy and fibrosis. Administration of a subpressor dose of Ang II to wild-type mice increases heart/body weight ratio, myocyte area, and cardiac collagen content in wild-type mice, but these responses are attenuated in Nox2^{-/-} mice.³⁴⁴ A similar reduction in cardiac hypertrophy is observed in mice with a cardiomyocyte-specific deletion of Rac1, which as expected, exhibit reduced ROS formation and reduced NADPH oxidase assembly after Ang II infusion compared with wild-type mice.³⁴⁵ Johar et al³⁴⁶ showed that in response to a pressor dose of Ang II, interstitial fibrosis, as well as upregulation of matrix proteins and matrix metalloproteinase 2 activity, is inhibited in Nox2^{-/-} animals compared with controls. This Nox2-dependent profibrotic effect can be reversed by the aldosterone antagonist spironolactone. Cardiac remodeling also occurs after MI, and Doerries et al³⁴⁷ found that cardiomyocyte hypertrophy and apoptosis, as well as interstitial fibrosis, are attenuated in mice lacking p47phox. Similarly, Nox2 knockout mice subjected to permanent left coronary ligation have less post-MI remodeling (as reflected by these same parameters) at 4 weeks despite no difference in infarct size.^{220,348} Thus, in multiple murine models, there is a clear link between Nox2, cardiac remodeling and fibrosis.

This link is not incontrovertible, however, and in some situations, Nox4 appears to be involved in cardiac remodeling. In contrast to its role in Ang II-induced cardiac hypertrophy, Nox2 does not appear to contribute to hypertrophy induced by pressure overload. Two separate groups reported no difference in myocardial hypertrophy, cardiac mass or atrial natriuretic factor expression in Nox2^{-/-} and wild-type mice subjected to aortic constriction.^{349,350} However, Nox2 does appear to contribute to interstitial hypertrophy and contractile dysfunction in this model.³⁵¹ Several studies have reported an increase in Nox4 expression in the heart in response to pressure overload,^{332,349,352} but only 2 groups have directly addressed the role of Nox4 in cardiac hypertrophy in this model. Using Nox4 knockout mice and cardiomyocyte-specific Nox4 overexpressors, Zhang et al³³² showed that there were no basal differences in cardiac function, but after pressure overload induced by aortic constriction, mice in which Nox4 is deleted exhibited greater cardiac hypertrophy, contractile dysfunction, and ventricular dilatation, whereas Nox4 transgenic mice were protected from these pathologies due to Nox4-dependent preservation of capillary density. However, Kuroda et al³⁵² found different results using mice with cardiac-specific Nox4 deletion or overexpression. In their hands, pressure overload-induced cardiac hypertrophy and contractile dysfunction were reduced in Nox4 knockout mice in parallel with improved mitochondrial function, but exacerbated in Nox4 overexpressing mice. With the exception of the Nox4 transgenic mice generated by Kuroda et al, mice used in both studies were on the C57/B16 background. However, the extent of pressure overload induced in the latter study was greater than that seen by Zhang et al, raising the possibility that the role of Nox4 may change with the severity of disease.

Interactions Among Nox Proteins

From the above summaries, it is evident that cells and tissues express multiple Nox homologues (Figure 2). Because their ultimate products (superoxide and hydrogen peroxide) are the same, the question arises as to the need for multiple oxidases in the same system. In some cases, their function appears to be redundant, whereas in others, each oxidase appears to regulate a distinct function. Moreover, there is cross-talk among oxidase systems, so that activation of Nox enzymes leads to subsequent activation of other sources of ROS, creating a positive feedback loop that can lead to vascular pathology.

In ECs, which express primarily Nox2 and Nox4, agonists often activate both enzymes and each Nox contributes similarly to the resultant physiological response. As an example, Nox2 and Nox4 are both required for the angiogenic response, and both participate in serum-induced proliferation.^{74,252,353} Both are upregulated by oscillatory shear stress and downregulated by pulsatile stimuli.²³² Interestingly, in these cells, Nox2 and Nox4 are perinuclear and colocalize with ER markers, perhaps explaining their redundant functions.⁷⁴ Moreover, knockdown of either of these genes using siRNA leads to an increase in the other, further supporting the notion that their functions are overlapping.³⁵⁴ One report, however, implicates both Nox2 and Nox4 in the regulation of cell proliferation, but provides evidence that only Nox2 prevents apoptosis and modifies the cytoskeleton in ECs.²¹³ In VSMCs from large arteries, where Nox1 and Nox4 are expressed in distinct subcellular compartments (caveolae and the nucleus and focal adhesions, respectively),¹⁰⁵ they mediate completely different physiological functions. Nox1 is required for hypertrophy and proliferation,^{176,181} whereas Nox4 mediates cell differentiation.^{229,251} Both participate in cell migration, although via different mechanisms.^{120,169} In this cell type, Nox1 and Nox4 are also differentially responsive to agonists,^{176,229} most likely a consequence of the differences in their subcellular localization. This hypothesis is supported by observations in HEK293 cells transfected with Nox2 or Nox4, in which each enzyme exhibits a specific subcellular localization (plasma membrane and ER, respectively) and each enzyme responds to specific agonists (Ang II and TNF- α versus no stimulation and insulin) and activates distinct signaling pathways.¹²² Ultimately, the subcellular localization of different Nox enzymes and their activation by specific external stimuli may dictate the integrated response to Nox activation.

It is also likely that Nox enzymes contribute to cardiovascular pathophysiology in a coordinated fashion. This has been best established in hypertension, where, as noted, T-cells coordinate the rise in blood pressure. In Ang II–induced hypertension, brain, vascular and kidney Nox enzymes contribute to blood pressure regulation.²⁹⁹ A similar coordinated response may occur in the vessel wall in atherosclerosis, where macrophages release cytokines in a Nox2-dependent manner, which then activate Nox1 and Nox2 in VSMCs and ECs, respectively.

It should also be noted that ROS production by Nox enzymes often activates other oxidase systems, thus amplifying the response. For example, in ECs exposed to oscillatory

shear stress, xanthine oxidase activation depends on prior activation of NADPH oxidases.²¹⁰ In addition, Ang II stimulation of these cells increases mitochondrial ROS production in a Nox-dependent manner.³⁵⁵ In many forms of hypertension, ROS production by Nox enzymes leads to the formation of peroxynitrite, which uncouples eNOS, greatly increasing superoxide production.²¹¹ The converse is true as well. Hydrogen peroxide from these sources, or even from initial Nox activation, can feedback to further activate Nox enzymes and amplify ROS production. Thus, Nox enzymes appear to act as both initiators and integrators of redox signaling via cross-talk with other ROS producing systems. The notion of cross-talk leads to an important concept: inappropriate feed forward activation of NADPH oxidases can lead to self-reinforcing loops that maintain and worsen pathology.

Conclusion

Tremendous progress since the discovery of the Nox family of enzymes has firmly established their essential role in cardiovascular physiology and pathophysiology in animal models. The field has progressed to a point where the next major challenge is to assess their role in human disease, and then translate these findings into treatments, while continuing to dissect the molecular mechanisms and pathways specific to individual Nox homologues. Several research groups and companies have developed homologue-specific inhibitors that are only now beginning to undergo clinical testing. Because of the far-reaching effects of Nox enzyme activation, the therapeutic potential of these compounds is enormous compared with previously used, nonspecific antioxidants. It is likely that these new agents will enable, for the first time, a true test of the redox hypothesis of human disease.

Sources of Funding

This work was supported by National Institutes of Health grants HL38206, HL095070, HL058863, and HL093115.

Disclosures

Emory University is the owner of a patent on Nox1 with B.L. and K.K.G. cited in the list of inventors.

References

1. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev*. 2007;87:245–313.
2. Bayraktutan U, Blayney L, Shah AM. Molecular characterization and localization of the NAD(P)H oxidase components gp91-phox and p22-phox in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2000;20:1903–1911.
3. Hohler B, Holzapfel B, Kummer W. NADPH oxidase subunits and superoxide production in porcine pulmonary artery endothelial cells. *Histochem Cell Biol*. 2000;114:29–37.
4. Lassegue B, Clempus RE. Vascular NAD(P)H oxidases: specific features, expression, and regulation. *Am J Physiol*. 2003;285:R277–R297.
5. Sumimoto H. Structure, regulation and evolution of nox-family NADPH oxidases that produce reactive oxygen species. *FEBS J*. 2008;275:3249–3277.
6. Biberstine-Kinkade KJ, DeLeo FR, Epstein RI, LeRoy BA, Nauseef WM, Dinauer MC. Heme-ligating histidines in flavocytochrome b(558): identification of specific histidines in gp91(phox). *J Biol Chem*. 2001;276:31105–31112.
7. Lassegue B, Griendling KK. Reactive oxygen species in hypertension: an update. *Am J Hypertens*. 2004;17:852–860.

8. DeCoursey TE. Voltage-gated proton channels find their dream job managing the respiratory burst in phagocytes. *Physiology*. 2010;25:27–40.
9. Moreland JG, Davis AP, Bailey G, Nauseef WM, Lamb FS. Anion channels, including clc-3, are required for normal neutrophil oxidative function, phagocytosis, and transendothelial migration. *J Biol Chem*. 2006;281:12277–12288.
10. Roos D, Kuhns DB, Maddalena A, Bustamante J, Kannengiesser C, de Boer M, van Leeuwen K, Koker MY, Wolach B, Roesler J, Malech HL, Holland SM, Gallin JI, Stasia MJ. Hematologically important mutations: the autosomal recessive forms of chronic granulomatous disease (second update). *Blood Cells Mol Dis*. 2010;44:291–299.
11. Roos D, Kuhns DB, Maddalena A, et al. Hematologically important mutations: X-linked chronic granulomatous disease (third update). *Blood Cells Mol Dis*. 2010;45:246–265.
12. Kao YY, Gianni D, Bohl B, Taylor RM, Bokoch GM. Identification of a conserved rac-binding site on NADPH oxidases supports a direct gtpase regulatory mechanism. *J Biol Chem*. 2008;283:12736–12746.
13. Lapouge K, Smith SJ, Groemping Y, Rittinger K. Architecture of the p40–p47–p67phox complex in the resting state of the NADPH oxidase: a central role for p67phox. *J Biol Chem*. 2002;277:10121–10128.
14. Raad H, Paclet MH, Boussetta T, Krowiarski Y, Morel F, Quinn MT, Gougerot-Pocidallo MA, Dang PM, El-Benna J. Regulation of the phagocyte NADPH oxidase activity: phosphorylation of gp91phox/nox2 by protein kinase C enhances its diaphorase activity and binding to rac2, p67phox, and p47phox. *FASEB J*. 2009;23:1011–1022.
15. Ueyama T, Kusakabe T, Karasawa S, Kawasaki T, Shimizu A, Son J, Leto TL, Miyawaki A, Saito N. Sequential binding of cytosolic phox complex to phagosomes through regulated adaptor proteins: evaluation using the novel monomeric kusabira-green system and live imaging of phagocytosis. *J Immunol*. 2008;181:629–640.
16. Ago T, Nunoi H, Ito T, Sumimoto H. Mechanism for phosphorylation-induced activation of the phagocyte NADPH oxidase protein p47(phox): triple replacement of serines 303, 304, and 328 with aspartates disrupts the sh3 domain-mediated intramolecular interaction in p47(phox), thereby activating the oxidase. *J Biol Chem*. 1999;274:33644–33653.
17. Reeves EP, Dekker LV, Forbes LV, Wientjes FB, Grogan A, Pappin DJ, Segal AW. Direct interaction between p47phox and protein kinase C: evidence for targeting of protein kinase c by p47phox in neutrophils. *Biochem J*. 1999;344:859–866.
18. Dang PM, Fontayne A, Hakim J, El Benna J, Perianin A. Protein kinase c zeta phosphorylates a subset of selective sites of the NADPH oxidase component p47phox and participates in formyl peptide-mediated neutrophil respiratory burst. *J Immunol*. 2001;166:1206–1213.
19. Fontayne A, Dang PM, Gougerot-Pocidallo MA, El-Benna J. Phosphorylation of p47phox sites by PKC alpha, beta II, delta, and zeta: effect on binding to p22phox and on NADPH oxidase activation. *Biochemistry*. 2002;41:7743–7750.
20. Ago T, Kuribayashi F, Hiroaki H, Takeya R, Ito T, Kohda D, Sumimoto H. Phosphorylation of p47phox directs phox homology domain from sh3 domain toward phosphoinositides, leading to phagocyte NADPH oxidase activation. *Proc Natl Acad Sci U S A*. 2003;100:4474–4479.
21. Hoyal CR, Gutierrez A, Young BM, Catz SD, Lin JH, Tschlis PN, Babior BM. Modulation of p47phox activity by site-specific phosphorylation: Akt-dependent activation of the NADPH oxidase. *Proc Natl Acad Sci U S A*. 2003;100:5130–5135.
22. Chen Q, Powell DW, Rane MJ, Singh S, Butt W, Klein JB, McLeish KR. Akt phosphorylates p47phox and mediates respiratory burst activity in human neutrophils. *J Immunol*. 2003;170:5302–5308.
23. Martyn KD, Kim MJ, Quinn MT, Dinauer MC, Knaus UG. P21-activated kinase (PAK) regulates NADPH oxidase activation in human neutrophils. *Blood*. 2005;106:3962–3969.
24. Frey RS, Rahman A, Kefer JC, Minshall RD, Malik AB. Pkc zeta regulates TNF-alpha-induced activation of NADPH oxidase in endothelial cells. *Circ Res*. 2002;90:1012–1019.
25. Touyz RM, Chen X, Tabet F, Yao G, He G, Quinn MT, Pagano PJ, Schiffrin EL. Expression of a functionally active gp91phox-containing neutrophil-type NAD(P)H oxidase in smooth muscle cells from human resistance arteries: regulation by angiotensin II. *Circ Res*. 2002;90:1205–1213.
26. Li JM, Shah AM. Mechanism of endothelial cell NADPH oxidase activation by angiotensin II: role of the p47phox subunit. *J Biol Chem*. 2003;278:12094–12100.
27. Touyz RM, Yao G, Schiffrin EL. C-Src induces phosphorylation and translocation of p47phox: role in superoxide generation by angiotensin II in human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2003;23:981–987.
28. Chowdhury AK, Watkins T, Parinandi NL, Saatian B, Kleinberg ME, Usatyuk PV, Natarajan V. Src-mediated tyrosine phosphorylation of p47phox in hyperoxia-induced activation of NADPH oxidase and generation of reactive oxygen species in lung endothelial cells. *J Biol Chem*. 2005;280:20700–20711.
29. Kanai F, Liu H, Field SJ, Akbary H, Matsuo T, Brown GE, Cantley LC, Yaffe MB. The px domains of p47phox and p40phox bind to lipid products of pi(3)k. *Nat Cell Biol*. 2001;3:675–678.
30. Zhan Y, He D, Newburger PE, Zhou GW. P47(phox) px domain of NADPH oxidase targets cell membrane via moesin-mediated association with the actin cytoskeleton. *J Cell Biochem*. 2004;92:795–809.
31. Touyz RM, Yao G, Quinn MT, Pagano PJ, Schiffrin EL. P47phox associates with the cytoskeleton through cortactin in human vascular smooth muscle cells: role in NAD(P)H oxidase regulation by angiotensin II. *Arterioscler Thromb Vasc Biol*. 2005;25:512–518.
32. Van Buul JD, Fernandez-Borja M, Anthony EC, Hordijk PL. Expression and localization of nox2 and nox4 in primary human endothelial cells. *Antioxidant Redox Signal*. 2005;7:308–317.
33. Groemping Y, Lapouge K, Smerdon SJ, Rittinger K. Molecular basis of phosphorylation-induced activation of the NADPH oxidase. *Cell*. 2003;113:343–355.
34. Ogura K, Nobuhisa I, Yuzawa S, Takeya R, Torikai S, Saikawa K, Sumimoto H, Inagaki F. NMR solution structure of the tandem Src homology 3 domains of p47phox complexed with a p22phox-derived proline-rich peptide. *J Biol Chem*. 2006;281:3660–3668.
35. Huang CK, Zhan L, Hannigan MO, Ai Y, Leto TL. P47(phox)-deficient NADPH oxidase defect in neutrophils of diabetic mouse strains, c57bl/6j-m db/db and db/+. *J Leukoc Biol*. 2000;67:210–215.
36. Jackson SH, Gallin JI, Holland SM. The p47phox mouse knock-out model of chronic granulomatous disease. *J Exp Med*. 1995;182:751–758.
37. Harbord M, Novelli M, Canas B, Power D, Davis C, Godovac-Zimmermann J, Roes J, Segal AW. Ym1 is a neutrophil granule protein that crystallizes in p47phox-deficient mice. *J Biol Chem*. 2002;277:5468–5475.
38. Freeman JL, Lambeth JD. NADPH oxidase activity is independent of p47phox in vitro. *J Biol Chem*. 1996;271:22578–22582.
39. Nisimoto Y, Motalebi S, Han CH, Lambeth JD. The p67(phox) activation domain regulates electron flow from NADPH to flavin in flavocytochrome b(558). *J Biol Chem*. 1999;274:22999–23005.
40. Ponting CP. Novel domains in NADPH oxidase subunits, sorting nexins, and ptdins 3-kinases: binding partners of sh3 domains? *Prot Sci*. 1996;5:2353–2357.
41. Leusen JH, de Klein A, Hilarius PM, Ahlin A, Palmblad J, Smith CI, Diekmann D, Hall A, Verhoeven AJ, Roos D. Disturbed interaction of p21-rac with mutated p67-phox causes chronic granulomatous disease. *J Exp Med*. 1996;184:1243–1249.
42. Koga H, Terasawa H, Nunoi H, Takeshige K, Inagaki F, Sumimoto H. Tetratricopeptide repeat (tpr) motifs of p67(phox) participate in interaction with the small gtpase rac and activation of the phagocyte NADPH oxidase. *J Biol Chem*. 1999;274:25051–25060.
43. Alloul N, Gorzalczy Y, Itan M, Sigal N, Pick E. Activation of the superoxide-generating NADPH oxidase by chimeric proteins consisting of segments of the cytosolic component p67(phox) and the small gtpase rac1. *Biochemistry*. 2001;40:14557–14566.
44. Han CH, Freeman JL, Lee T, Motalebi SA, Lambeth JD. Regulation of the neutrophil respiratory burst oxidase: identification of an activation domain in p67(phox). *J Biol Chem*. 1998;273:16663–16668.
45. Yuzawa S, Miyano K, Honbou K, Inagaki F, Sumimoto H. The domain organization of p67 phox, a protein required for activation of the superoxide-producing NADPH oxidase in phagocytes. *J Innate Immunity*. 2009;1:543–555.
46. Nakamura R, Sumimoto H, Mizuki K, Hata K, Ago T, Kitajima S, Takeshige K, Sakaki Y, Ito T. The PC motif: a novel and evolutionarily conserved sequence involved in interaction between p40phox and p67phox, sh3 domain-containing cytosolic factors of the phagocyte NADPH oxidase. *Eur J Biochem/FEBS*. 1998;251:583–589.
47. Kuribayashi F, Nunoi H, Wakamatsu K, Tsunawaki S, Sato K, Ito T, Sumimoto H. The adaptor protein p40(phox) as a positive regulator of the superoxide-producing phagocyte oxidase. *EMBO J*. 2002;21:6312–6320.

48. Leto TL, Adams AG, de Mendez I. Assembly of the phagocyte NADPH oxidase: binding of Src homology 3 domains to proline-rich targets. *Proc Natl Acad Sci U S A*. 1994;91:10650–10654.
49. Finan P, Shimizu Y, Gout I, Hsuan J, Truong O, Butcher C, Bennett P, Waterfield MD, Kellie S. An sh3 domain and proline-rich sequence mediate an interaction between two components of the phagocyte NADPH oxidase complex. *J Biol Chem*. 1994;269:13752–13755.
50. Mizuki K, Takeya R, Kuribayashi F, Nobuhisa I, Kohda D, Nunoi H, Takeshige K, Sumimoto H. A region c-terminal to the proline-rich core of p47phox regulates activation of the phagocyte NADPH oxidase by interacting with the c-terminal sh3 domain of p67phox. *Arch Biochem Biophys*. 2005;444:185–194.
51. Dusi S, Rossi F. Activation of NADPH oxidase of human neutrophils involves the phosphorylation and the translocation of cytosolic p67phox. *Biochem J*. 1993;296:367–371.
52. Benna JE, Dang PM, Gaudry M, Fay M, Morel F, Hakim J, Gougerot-Pocidalo MA. Phosphorylation of the respiratory burst oxidase subunit p67(phox) during human neutrophil activation: regulation by protein kinase c-dependent and independent pathways. *J Biol Chem*. 1997;272:17204–17208.
53. Forbes LV, Truong O, Wientjes FB, Moss SJ, Segal AW. The major phosphorylation site of the NADPH oxidase component p67phox is thr233. *Biochem J*. 1999;338:99–105.
54. Dang PM, Morel F, Gougerot-Pocidalo MA, El Benna J. Phosphorylation of the NADPH oxidase component p67(phox) by erk2 and p38mapk: selectivity of phosphorylated sites and existence of an intramolecular regulatory domain in the tetratricopeptide-rich region. *Biochemistry*. 2003;42:4520–4526.
55. Zhao X, Xu B, Bhattacharjee A, Oldfield CM, Wientjes FB, Feldman GM, Cathcart MK. Protein kinase cdelta regulates p67phox phosphorylation in human monocytes. *J Leukoc Biol*. 2005;77:414–420.
56. Dang PM, Raad H, Derkawi RA, Boussetta T, Pacllet MH, Belambri SA, Makni-Maalej K, Kroviarski Y, Morel F, Gougerot-Pocidalo MA, El-Benna J. The NADPH oxidase cytosolic component p67phox is constitutively phosphorylated in human neutrophils: regulation by a protein tyrosine kinase, mek1/2 and phosphatases 1/2a. *Biochem Pharmacol*. 2011;82:1145–1152.
57. Sancho-Shimizu V, Malo D. Sequencing, expression, and functional analyses support the candidacy of nc2 in susceptibility to salmonella typhimurium infection in wild-derived mice. *J Immunol*. 2006;176:6954–6961.
58. Ueyama T, Tatsuno T, Kawasaki T, Tsujibe S, Shirai Y, Sumimoto H, Leto TL, Saito N. A regulated adaptor function of p40phox: Distinct p67phox membrane targeting by p40phox and by p47phox. *Mol Biol Cell*. 2007;18:441–454.
59. Ago T, Takeya R, Hiroaki H, Kuribayashi F, Ito T, Kohda D, Sumimoto H. The px domain as a novel phosphoinositide-binding module. *Biochem Biophys Res Commun*. 2001;287:733–738.
60. Zhan Y, Virbasiu JV, Song X, Pomerleau DP, Zhou GW. The p40phox and p47phox px domains of NADPH oxidase target cell membranes via direct and indirect recruitment by phosphoinositides. *J Biol Chem*. 2002;277:4512–4518.
61. Tian W, Li XJ, Stull ND, Ming W, Suh CI, Bissonnette SA, Yaffe MB, Grinstein S, Atkinson SJ, Dinuer MC. Fc gamma r-stimulated activation of the NADPH oxidase: Phosphoinositide-binding protein p40phox regulates NADPH oxidase activity after enzyme assembly on the phagosome. *Blood*. 2008;112:3867–3877.
62. Cross AR. P40(phox) participates in the activation of NADPH oxidase by increasing the affinity of p47(phox) for flavocytochrome b(558). *Biochem J*. 2000;349:113–117.
63. Ueyama T, Nakakita J, Nakamura T, Kobayashi T, Kobayashi T, Son J, Sakuma M, Sakaguchi H, Leto TL, Saito N. Cooperation of p40phox with p47phox for nox2 activation during fcγr-mediated phagocytosis: mechanism for acquisition of p40phox pi(3)p binding. *J Biol Chem*. 2011;286:40693–40705.
64. Dusi S, Donini M, Rossi F. Mechanisms of NADPH oxidase activation: translocation of p40phox, rac1 and rac2 from the cytosol to the membranes in human neutrophils lacking p47phox or p67phox. *Biochem J*. 1996;314:409–412.
65. Ellson C, Davidson K, Anderson K, Stephens LR, Hawkins PT. Ptdins3p binding to the px domain of p40phox is a physiological signal in NADPH oxidase activation. *EMBO J*. 2006;25:4468–4478.
66. Bissonnette SA, Glazier CM, Stewart MQ, Brown GE, Ellson CD, Yaffe MB. Phosphatidylinositol 3-phosphate-dependent and -independent functions of p40phox in activation of the neutrophil NADPH oxidase. *J Biol Chem*. 2008;283:2108–2119.
67. Fan LM, Teng L, Li JM. Knockout of p47 phox uncovers a critical role of p40 phox in reactive oxygen species production in microvascular endothelial cells. *Arterioscler Thromb Vasc Biol*. 2009;29:1651–1656.
68. Suh CI, Stull ND, Li XJ, Tian W, Price MO, Grinstein S, Yaffe MB, Atkinson S, Dinuer MC. The phosphoinositide-binding protein p40phox activates the NADPH oxidase during Fcγm1a receptor-induced phagocytosis. *J Exp Med*. 2006;203:1915–1925.
69. Chessa TA, Anderson KE, Hu Y, Xu Q, Rausch O, Stephens LR, Hawkins PT. Phosphorylation of threonine 154 in p40phox is an important physiological signal for activation of the neutrophil NADPH oxidase. *Blood*. 2010;116:6027–6036.
70. Bouin AP, Grandvaux N, Vignais PV, Fuchs A. P40(phox) is phosphorylated on threonine 154 and serine 315 during activation of the phagocyte NADPH oxidase: implication of a protein kinase c-type kinase in the phosphorylation process. *J Biol Chem*. 1998;273:30097–30103.
71. Someya A, Nunoi H, Hasebe T, Nagaoka I. Phosphorylation of p40-phox during activation of neutrophil NADPH oxidase. *J Leukoc Biol*. 1999;66:851–857.
72. Lopes LR, Dagher MC, Gutierrez A, Young B, Bouin AP, Fuchs A, Babor BM. Phosphorylated p40phox as a negative regulator of NADPH oxidase. *Biochemistry*. 2004;43:3723–3730.
73. Matute JD, Arias AA, Wright NA, Wrobel I, Waterhouse CC, Li XJ, Marchal CC, Stull ND, Lewis DB, Steele M, Kellner JD, Yu W, Meroueh SO, Nauseef WM, Dinuer MC. A new genetic subgroup of chronic granulomatous disease with autosomal recessive mutations in p40 phox and selective defects in neutrophil NADPH oxidase activity. *Blood*. 2009;114:3309–3315.
74. Petry A, Djordjevic T, Weitnauer M, Kietzmann T, Hess J, Gorch A. Nox2 and nox4 mediate proliferative response in endothelial cells. *Antioxidant Redox Signal*. 2006;8:1473–1484.
75. Hahn NE, Meischl C, Wijnker PJ, Musters RJ, Fornerod M, Janssen HW, Paulus WJ, van Rossum AC, Niessen HW, Krijnen PA. Nox2, p22phox and p47phox are targeted to the nuclear pore complex in ischemic cardiomyocytes colocalizing with local reactive oxygen species. *Cell Physiol Biochem*. 2011;27:471–478.
76. Szanto I, Rubbia-Brandt L, Kiss P, Steger K, Banfi B, Kovari E, Herrmann F, Hadengue A, Krause KH. Expression of nox1, a superoxide-generating NADPH oxidase, in colon cancer and inflammatory bowel disease. *J Pathol*. 2005;207:164–176.
77. Hanna IR, Hilenski LL, Dikalova A, Taniyama Y, Dikalov S, Lyle A, Quinn MT, Lassegue B, Griendling KK. Functional association of nox1 with p22phox in vascular smooth muscle cells. *Free Radic Biol Med*. 2004;37:1542–1549.
78. Ambasta RK, Kumar P, Griendling KK, Schmidt HH, Busse R, Brandes RP. Direct interaction of the novel nox proteins with p22phox is required for the formation of a functionally active NADPH oxidase. *J Biol Chem*. 2004;279:45935–45941.
79. Kawahara T, Ritsick D, Cheng G, Lambeth JD. Point mutations in the proline-rich region of p22phox are dominant inhibitors of nox1- and nox2-dependent reactive oxygen generation. *J Biol Chem*. 2005;280:31859–31869.
80. Takeya R, Ueno N, Kami K, Taura M, Kohjima M, Izaki T, Nunoi H, Sumimoto H. Novel human homologues of p47phox and p67phox participate in activation of superoxide-producing NADPH oxidases. *J Biol Chem*. 2003;278:25234–25246.
81. Banfi B, Clark RA, Steger K, Krause KH. Two novel proteins activate superoxide generation by the NADPH oxidase nox1. *J Biol Chem*. 2003;278:3510–3513.
82. Geiszt M, Lekstrom K, Witt J, Leto TL. Proteins homologous to p47phox and p67phox support superoxide production by NAD(P)H oxidase 1 in colon epithelial cells. *J Biol Chem*. 2003;278:20006–20012.
83. Niu XL, Madamanchi NR, Vendrov AE, Tchivilev I, Rojas M, Madamanchi C, Brandes RP, Krause KH, Humphries J, Smith A, Burnand KG, Runge MS. Nox activator 1: A potential target for modulation of vascular reactive oxygen species in atherosclerotic arteries. *Circulation*. 2010;121:549–559.
84. Ambasta RK, Schreiber JG, Janiszewski M, Busse R, Brandes RP. Nox1 is a central component of the smooth muscle NADPH oxidase in mice. *Free Radic Biol Med*. 2006;41:193–201.
85. Ali MI, Ketsawatsomkron P, Belin de Chantemele EJ, Mintz JD, Muta K, Salet C, Black SM, Tremblay ML, Fulton DJ, Marrero MB, Stepp

- DW. Deletion of protein tyrosine phosphatase 1b improves peripheral insulin resistance and vascular function in obese, leptin-resistant mice via reduced oxidant tone. *Circ Res*. 2009;105:1013–1022.
86. Ago T, Kitazono T, Kuroda J, Kumai Y, Kamouchi M, Ooboshi H, Wakisaka M, Kawahara T, Rokutan K, Ibayashi S, Iida M. NAD(P)H oxidases in rat basilar arterial endothelial cells. *Stroke*. 2005;36:1040–1046.
 87. Ueyama T, Geiszt M, Leto TL. Involvement of rac1 in activation of multicomponent nox1- and nox3-based NADPH oxidases. *Mol Cell Biol*. 2006;26:2160–2174.
 88. Dutta S, Rittinger K. Regulation of nox1 activity through reversible interactions with p22 and noxal. *PLoS One*. 2010;5:e10478.
 89. Park HS, Park D, Bae YS. Molecular interaction of NADPH oxidase 1 with betapix and nox organizer 1. *Biochem Biophys Res Commun*. 2006;339:985–990.
 90. Cheng G, Lambeth JD. Nox1, regulation of lipid binding, localization, and activation of nox1 by the phox homology (px) domain. *J Biol Chem*. 2004;279:4737–4742.
 91. Kiss PJ, Knisz J, Zhang Y, Baltrusaitis J, Sigmund CD, Thalmann R, Smith RJ, Verpy E, Banfi B. Inactivation of NADPH oxidase organizer 1 results in severe imbalance. *Curr Biol*. 2006;16:208–213.
 92. Gianni D, DerMardirossian C, Bokoch GM. Direct interaction between tks proteins and the n-terminal proline-rich region (pr) of nox1 mediates nox1-dependent ROS generation. *Eur J Cell Biol*. 2011;90:164–171.
 93. Kim JS, Diebold BA, Babior BM, Knaus UG, Bokoch GM. Regulation of nox1 activity via protein kinase a-mediated phosphorylation of noxal and 14–3-3 binding. *J Biol Chem*. 2007;282:34787–34800.
 94. Oh H, Jung HY, Kim J, Bae YS. Phosphorylation of serine282 in NADPH oxidase activator 1 by erk desensitizes egf-induced ROS generation. *Biochem Biophys Res Commun*. 2010;394:691–696.
 95. Krovjarski Y, Debbabi M, Bachoual R, Perianin A, Gougerot-Pocidalo MA, El-Benna J, Dang PM. Phosphorylation of NADPH oxidase activator 1 (nox1) on serine 282 by map kinases and on serine 172 by protein kinase c and protein kinase a prevents nox1 hyperactivation. *FASEB J*. 2010;24:2077–2092.
 96. Gianni D, Taulat N, DerMardirossian C, Bokoch GM. C-Src-mediated phosphorylation of noxal and tks4 induces the reactive oxygen species (ROS)-dependent formation of functional invadopodia in human colon cancer cells. *Mol Biol Cell*. 2010;21:4287–4298.
 97. Flaherty JP, Spruce CA, Fairfield HE, Bergstrom DE. Generation of a conditional null allele of NADPH oxidase activator 1 (nox1). *Genesis*. 2010;48:568–575.
 98. Cheng G, Diebold BA, Hughes Y, Lambeth JD. Nox1-dependent reactive oxygen generation is regulated by rac1. *J Biol Chem*. 2006;281:17718–17726.
 99. Bokoch GM, Zhao T. Regulation of the phagocyte NADPH oxidase by rac gtpase. *Antioxidant Redox Signal*. 2006;8:1533–1548.
 100. Park HS, Lee SH, Park D, Lee JS, Ryu SH, Lee WJ, Rhee SG, Bae YS. Sequential activation of phosphatidylinositol 3-kinase, beta pix, rac1, and nox1 in growth factor-induced production of h2o2. *Mol Cell Biol*. 2004;24:4384–4394.
 101. Diebold BA, Fowler B, Lu J, Dinauer MC, Bokoch GM. Antagonistic cross-talk between rac and cdc42 gtpases regulates generation of reactive oxygen species. *J Biol Chem*. 2004;279:28136–28142.
 102. Gianni D, Bohl B, Courtneidge SA, Bokoch GM. The involvement of the tyrosine kinase c-src in the regulation of reactive oxygen species generation mediated by NADPH oxidase-1. *Mol Biol Cell*. 2008;19:2984–2994.
 103. Helmcke I, Heumuller S, Tikkanen R, Schroder K, Brandes RP. Identification of structural elements in nox1 and nox4 controlling localization and activity. *Antioxidant Redox Signal*. 2009;11:1279–1287.
 104. Chen F, Pandey D, Chadli A, Catravas JD, Chen T, Fulton DJ. Hsp90 regulates NADPH oxidase activity and is necessary for superoxide but not hydrogen peroxide production. *Antioxidant Redox Signal*. 2011;14:2107–2119.
 105. Hilenski LL, Clemus RE, Quinn MT, Lambeth JD, Griendling KK. Distinct subcellular localizations of nox1 and nox4 in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2004;24:677–683.
 106. Miller FJ Jr, Filali M, Huss GJ, Stanic B, Chamseddine A, Barna TJ, Lamb FS. Cytokine activation of nuclear factor kappa b in vascular smooth muscle cells requires signaling endosomes containing nox1 and clc-3. *Circ Res*. 2007;101:663–671.
 107. Zhang G, Zhang F, Muh R, Yi F, Chalupsky K, Cai H, Li PL. Autocrine/paracrine pattern of superoxide production through NAD(P)H oxidase in coronary arterial myocytes. *Am J Physiol Heart Circ Physiol*. 2007;292:H483–H495.
 108. Miller FJ Jr, Chu X, Stanic B, Tian X, Sharma RV, Davisson RL, Lamb FS. A differential role for endocytosis in receptor-mediated activation of nox1. *Antioxidant Redox Signal*. 2010;12:583–593.
 109. Chu X, Filali M, Stanic B, Takapoo M, Sheehan A, Bhalla R, Lamb FS, Miller FJ Jr. A critical role for chloride channel-3 (cl-3) in smooth muscle cell activation and neointima formation. *Arterioscler Thromb Vasc Biol*. 2011;31:345–351.
 110. Janiszewski M, Lopes LR, Carmo AO, Pedro MA, Brandes RP, Santos CX, Laurindo FR. Regulation of NAD(P)H oxidase by associated protein disulfide isomerase in vascular smooth muscle cells. *J Biol Chem*. 2005;280:40813–40819.
 111. Fernandes DC, Manoel AH, Wosniak J Jr, Laurindo FR. Protein disulfide isomerase overexpression in vascular smooth muscle cells induces spontaneous preemptive NADPH oxidase activation and nox1 mRNA expression: effects of nitrosothiol exposure. *Arch Biochem Biophys*. 2009;484:197–204.
 112. Geiszt M, Kopp JB, Varnai P, Leto TL. Identification of renox, an NAD(P)H oxidase in kidney. *Proc Natl Acad Sci U S A*. 2000;97:8010–8014.
 113. Shiose A, Kuroda J, Tsuruya K, Hirai M, Hirakata H, Naito S, Hattori M, Sakaki Y, Sumimoto H. A novel superoxide-producing NAD(P)H oxidase in kidney. *J Biol Chem*. 2001;276:1417–1423.
 114. Lee CF, Qiao M, Schroder K, Zhao Q, Asmis R. Nox4 is a novel inducible source of reactive oxygen species in monocytes and macrophages and mediates oxidized low density lipoprotein-induced macrophage death. *Circ Res*. 2010;106:1489–1497.
 115. Martyn KD, Frederick LM, von Lohneysen K, Dinauer MC, Knaus UG. Functional analysis of nox4 reveals unique characteristics compared to other NADPH oxidases. *Cell Signal*. 2006;18:69–82.
 116. Djordjevic T, BelAiba RS, Bonello S, Pfeilschifter J, Hess J, Gorlach A. Human urotensin II is a novel activator of NADPH oxidase in human pulmonary artery smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2005;25:519–525.
 117. von Lohneysen K, Noack D, Wood MR, Friedman JS, Knaus UG. Structural insights into nox4 and nox2: motifs involved in function and cellular localization. *Mol Cell Biol*. 2010;30:961–975.
 118. Serrander L, Cartier L, Bedard K, Banfi B, Lardy B, Plastre O, Sienkiewicz A, Forro L, Schlegel W, Krause KH. Nox4 activity is determined by mRNA levels and reveals a unique pattern of ROS generation. *Biochem J*. 2007;406:105–114.
 119. Nisimoto Y, Jackson HM, Ogawa H, Kawahara T, Lambeth JD. Constitutive NADPH-dependent electron transferase activity of the nox4 dehydrogenase domain. *Biochemistry*. 2010;49:2433–2442.
 120. Lyle AN, Deshpande NN, Taniyama Y, Seidel-Rogol B, Pounkova L, Du P, Papaharalambus C, Lassegue B, Griendling KK, Poldip2, a novel regulator of nox4 and cytoskeletal integrity in vascular smooth muscle cells. *Circ Res*. 2009;105:249–259.
 121. von Lohneysen K, Noack D, Jesaitis AJ, Dinauer MC, Knaus UG. Mutational analysis reveals distinct features of the nox4-p22 phox complex. *J Biol Chem*. 2008;283:35273–35282.
 122. Anilkumar N, Weber R, Zhang M, Brewer A, Shah AM. Nox4 and nox2 NADPH oxidases mediate distinct cellular redox signaling responses to agonist stimulation. *Arterioscler Thromb Vasc Biol*. 2008;28:1347–1354.
 123. Gorin Y, Ricono JM, Kim NH, Bhandari B, Choudhury GG, Abboud HE. Nox4 mediates angiotensin II-induced activation of akt/protein kinase B in mesangial cells. *Am J Physiol Renal Physiol*. 2003;285:F219–F229.
 124. Wu RF, Ma Z, Myers DP, Terada LS. HIV-1 tat activates dual nox pathways leading to independent activation of erk and jnk map kinases. *J Biol Chem*. 2007;282:37412–37419.
 125. Lee S, Gharavi NM, Honda H, Chang I, Kim B, Jen N, Li R, Zimman A, Berliner JA. A role for NADPH oxidase 4 in the activation of vascular endothelial cells by oxidized phospholipids. *Free Radic Biol Med*. 2009;47:145–151.
 126. Zhuang J, Jiang T, Lu D, Luo Y, Zheng C, Feng J, Yang D, Chen C, Yan X. NADPH oxidase 4 mediates reactive oxygen species induction of cd146 dimerization in VEGF signal transduction. *Free Radic Biol Med*. 2010;49:227–236.
 127. Park HS, Jung HY, Park EY, Kim J, Lee WJ, Bae YS. Cutting edge: direct interaction of tlr4 with NAD(P)H oxidase 4 isozyme is essential for lipopolysaccharide-induced production of reactive oxygen species and activation of NF-kappa B. *J Immunol*. 2004;173:3589–3593.

128. Mahadev K, Motoshima H, Wu X, Ruddy JM, Arnold RS, Cheng G, Lambeth JD, Goldstein BJ. The NAD(P)H oxidase homolog nox4 modulates insulin-stimulated generation of h₂O₂ and plays an integral role in insulin signal transduction. *Mol Cell Biol*. 2004;24:1844–1854.
129. Kuroda J, Nakagawa K, Yamasaki T, Nakamura K, Takeya R, Kuribayashi F, Imajoh-Ohmi S, Igarashi K, Shibata Y, Sueishi K, Sumimoto H. The superoxide-producing NAD(P)H oxidase nox4 in the nucleus of human vascular endothelial cells. *Genes Cells*. 2005;10:1139–1151.
130. Hu T, Ramachandrarao SP, Siva S, Valancius C, Zhu Y, Mahadev K, Toh I, Goldstein BJ, Woolkalis M, Sharma K. Reactive oxygen species production via NADPH oxidase mediates TGF-beta-induced cytoskeletal alterations in endothelial cells. *Am J Physiol Renal Physiol*. 2005;289:F816–F825.
131. Park HS, Chun JN, Jung HY, Choi C, Bae YS. Role of NADPH oxidase 4 in lipopolysaccharide-induced proinflammatory responses by human aortic endothelial cells. *Cardiovasc Res*. 2006;72:447–455.
132. Grange L, Nguyen MV, Lardy B, Derouazi M, Campion Y, Trocme C, Paclot MH, Gaudin P, Morel F. NAD(P)H oxidase activity of nox4 in chondrocytes is both inducible and involved in collagenase expression. *Antioxidant Redox Signal*. 2006;8:1485–1496.
133. Basuroy S, Tcheranova D, Bhattacharya S, Leffler CW, Parfenova H. Nox4 NADPH oxidase-derived reactive oxygen species, via endogenous carbon monoxide, promote survival of brain endothelial cells during TNF-alpha-induced apoptosis. *Am J Physiol Cell Physiol*. 2011;300:C256–C265.
134. New D, Block K, Bhandari B, Gorin YC, Abboud HE. Igf-1 increases the expression of fibronectin by nox4-dependent akt phosphorylation in renal tubular epithelial cells. *Am J Physiol Cell Physiol*. 2011;302:C122–C130.
135. Bondi CD, Manickam N, Lee DY, Block K, Gorin Y, Abboud HE, Barnes JL. NAD(P)H oxidase mediates tgf-beta1-induced activation of kidney myofibroblasts. *J Am Soc Nephrol*. 2010;21:93–102.
136. Gupte SA, Kaminski PM, Floyd B, Agarwal R, Ali N, Ahmad M, Edwards J, Wolin MS. Cytosolic NADPH may regulate differences in basal nox oxidase-derived superoxide generation in bovine coronary and pulmonary arteries. *Am J Physiol Heart Circ Physiol*. 2005;288:H13–H21.
137. Spencer NY, Yan Z, Boudreau RL, Zhang Y, Luo M, Li Q, Tian X, Shah AM, Davison RL, Davidson B, Banfi B, Engelhardt JF. Control of hepatic nuclear superoxide production by glucose 6-phosphate dehydrogenase and NADPH oxidase-4. *J Biol Chem*. 2011;286:8977–8987.
138. Ben Mkaddem S, Pedruzzi E, Werts C, Coant N, Bens M, Cluzeaud F, Goujon JM, Ogier-Denis E, Vandewalle A. Heat shock protein gp96 and NAD(P)H oxidase 4 play key roles in toll-like receptor 4-activated apoptosis during renal ischemia/reperfusion injury. *Cell Death Differentiation*. 2010;17:1474–1485.
139. Ago T, Kuroda J, Pain J, Fu C, Li H, Sadoshima J. Upregulation of nox4 by hypertrophic stimuli promotes apoptosis and mitochondrial dysfunction in cardiac myocytes. *Circ Res*. 2010;106:1253–1264.
140. Goyal P, Weissmann N, Rose F, Grimminger F, Schafers HJ, Seeger W, Hanze J. Identification of novel nox4 splice variants with impact on ROS levels in a549 cells. *Biochem Biophys Res Commun*. 2005;329:32–39.
141. Jackson HM, Kawahara T, Nisimoto Y, Smith SM, Lambeth JD. Nox4 b-loop creates an interface between the transmembrane and dehydrogenase domains. *J Biol Chem*. 2010;285:10281–10290.
142. Dikalov SI, Dikalova AE, Bikineyeva AT, Schmidt HH, Harrison DG, Griendling KK. Distinct roles of nox1 and nox4 in basal and angiotensin II-stimulated superoxide and hydrogen peroxide production. *Free Radic Biol Med*. 2008;45:1340–1351.
143. Takac I, Schroder K, Zhang L, Lardy B, Anilkumar N, Lambeth JD, Shah AM, Morel F, Brandes RP. The e-loop is involved in hydrogen peroxide formation by the NADPH oxidase nox4. *J Biol Chem*. 2011;286:13304–13313.
144. Zhang L, Nguyen MV, Lardy B, Jesaitis AJ, Grichine A, Rousset F, Talbot M, Paclot MH, Qian G, Morel F. New insight into the nox4 subcellular localization in hek293 cells: first monoclonal antibodies against nox4. *Biochimie*. 2011;93:457–468.
145. Yoshida LS, Tsunawaki S. Expression of NADPH oxidases and enhanced h(2)o(2)-generating activity in human coronary artery endothelial cells upon induction with tumor necrosis factor-alpha. *Int Immunopharmacol*. 2008;8:1377–1385.
146. Banfi B, Molnar G, Maturana A, Steger K, Hegedus B, Demareux N, Krause KH. A Ca(2+)-activated NADPH oxidase in testis, spleen, and lymph nodes. *J Biol Chem*. 2001;276:37594–37601.
147. Banfi B, Tirone F, Durussel I, Knisz J, Moskwa P, Molnar GZ, Krause KH, Cox JA. Mechanism of Ca²⁺ activation of the NADPH oxidase 5 (nox5). *J Biol Chem*. 2004;279:18583–18591.
148. Fu X, Beer DG, Behar J, Wands J, Lambeth D, Cao W. Camp-response element-binding protein mediates acid-induced NADPH oxidase nox5-s expression in Barrett esophageal adenocarcinoma cells. *J Biol Chem*. 2006;281:20368–20382.
149. Si J, Fu X, Behar J, Wands J, Beer DG, Souza RF, Spechler SJ, Lambeth D, Cao W. NADPH oxidase nox5-s mediates acid-induced cyclooxygenase-2 expression via activation of NF-kappaB in Barrett's esophageal adenocarcinoma cells. *J Biol Chem*. 2007;282:16244–16255.
150. BelAiba RS, Djordjevic T, Petry A, Diemer K, Bonello S, Banfi B, Hess J, Pogrebniak A, Bickel C, Grolach A. Nox5 variants are functionally active in endothelial cells. *Free Radic Biol Med*. 2007;42:446–459.
151. Kawahara T, Jackson HM, Smith SM, Simpson PD, Lambeth JD. Nox5 forms a functional oligomer mediated by self-association of its dehydrogenase domain. *Biochemistry*. 2011;50:2013–2025.
152. Hong J, Resnick M, Behar J, Wands J, DeLellis RA, Cao W. Role of rac1 in regulation of nox5-s function in Barrett's esophageal adenocarcinoma cells. *Am J Physiol Cell Physiol*. 2011;301:C413–C420.
153. Montezano AC, Burger D, Paravicini TM, Chignalia AZ, Yusuf H, Almasri M, He Y, Callera GE, He G, Krause KH, Lambeth D, Quinn MT, Touyz RM. Nicotinamide adenine dinucleotide phosphate reduced oxidase 5 (nox5) regulation by angiotensin II and endothelin-1 is mediated via calcium/calmodulin-dependent, rac-1-independent pathways in human endothelial cells. *Circ Res*. 2010;106:1363–1373.
154. Jay DB, Papaharalambus CA, Seidel-Rogol B, Dikalova AE, Lassegue G, Griendling KK. Nox5 mediates pdgf-induced proliferation in human aortic smooth muscle cells. *Free Radic Biol Med*. 2008;45:329–335.
155. Serrander L, Jaquet V, Bedard K, Plastre O, Hartley O, Arnaudeau S, Demareux N, Schlegel W, Krause KH. Nox5 is expressed at the plasma membrane and generates superoxide in response to protein kinase c activation. *Biochimie*. 2007;89:1159–1167.
156. Tirone F, Radu L, Craescu CT, Cox JA. Identification of the binding site for the regulatory calcium-binding domain in the catalytic domain of nox5. *Biochemistry*. 2010;49:761–771.
157. Tirone F, Cox JA. NADPH oxidase 5 (nox5) interacts with and is regulated by calmodulin. *FEBS Lett*. 2007;581:1202–1208.
158. Jagannand D, Church JE, Banfi B, Stuehr DJ, Marrero MB, Fulton DJ. Novel mechanism of activation of NADPH oxidase 5. Calcium sensitization via phosphorylation. *J Biol Chem*. 2007;282:6494–6507.
159. Pandey D, Gratton JP, Rafikov R, Black SM, Fulton DJ. Calcium/calmodulin-dependent kinase ii mediates the phosphorylation and activation of NADPH oxidase 5. *Mol Pharmacol*. 2011;80:407–415.
160. El Jamali A, Valente AJ, Lechleiter JD, Gamez MJ, Pearson DW, Nauseef WM, Clark RA. Novel redox-dependent regulation of nox5 by the tyrosine kinase c-abl. *Free Radic Biol Med*. 2008;44:868–881.
161. Zhou X, Li D, Resnick MB, Behar J, Wands J, Cao W. Signaling in h₂O₂-induced increase in cell proliferation in Barrett's esophageal adenocarcinoma cells. *J Pharmacol Exp Ther*. 2011;339:218–227.
162. Kawahara T, Lambeth JD. Phosphatidylinositol (4,5)-bisphosphate modulates nox5 localization via an n-terminal polybasic region. *Mol Biol Cell*. 2008;19:4020–4031.
163. Taylor RM, Burritt JB, Baniulis D, Foubert TR, Lord CI, Dinauer MC, Parkos CA, Jesaitis AJ. Site-specific inhibitors of NADPH oxidase activity and structural probes of flavocytochrome B: characterization of six monoclonal antibodies to the p22phox subunit. *J Immunol*. 2004;173:7349–7357.
164. Zhu Y, Marchal CC, Casbon AJ, Stull N, von Lohneysen K, Knaus UG, Jesaitis AJ, McCormick S, Nauseef WM, Dinauer MC. Deletion mutagenesis of p22phox subunit of flavocytochrome b558: identification of regions critical for gp91phox maturation and NADPH oxidase activity. *J Biol Chem*. 2006;281:30336–30346.
165. Biberstine-Kinkade KJ, Yu L, Stull N, LeRoy B, Bennett S, Cross A, Dinauer MC. Mutagenesis of p22(phox) histidine 94 A histidine in this position is not required for flavocytochrome b558 function. *J Biol Chem*. 2002;277:30368–30374.
166. Porter CD, Parkar MH, Verhoeven AJ, Levinsky RJ, Collins MK, Kinnon C. P22-phox-deficient chronic granulomatous disease: reconstitution by retrovirus-mediated expression and identification of a biosynthetic intermediate of gp91-phox. *Blood*. 1994;84:2767–2775.
167. Nakano Y, Longo-Guess CM, Bergstrom DE, Nauseef WM, Jones SM, Banfi B. Mutation of the cyb gene encoding p22phox causes vestibular and immune defects in mice. *J Clin Invest*. 2008;118:1176–1185.

168. Lewis EM, Sergeant S, Ledford B, Stull N, Dinauer MC, McPhail LC. Phosphorylation of p22phox on threonine 147 enhances NADPH oxidase activity by promoting p47phox binding. *J Biol Chem*. 2010;285:2959–2967.
169. Lee MY, San Martin A, Mehta PK, Dikalova AE, Garrido AM, Datla SR, Lyons E, Krause KH, Banfi B, Lambeth JD, Lassegue B, Griendling KK. Mechanisms of vascular smooth muscle NADPH oxidase 1 (nox1) contribution to injury-induced neointimal formation. *Arterioscler Thromb Vasc Biol*. 2009;29:480–487.
170. Gavazzi G, Banfi B, Deffert C, Fiette L, Schappi M, Herrmann F, Krause KH. Decreased blood pressure in nox1-deficient mice. *FEBS Lett*. 2006;580:497–504.
171. San Martin A, Foncea R, Laurindo FR, Ebensperger R, Griendling KK, Leighton F. Nox1-based NADPH oxidase-derived superoxide is required for VSMC activation by advanced glycation end-products. *Free Radic Biol Med*. 2007;42:1671–1679.
172. Wendt MC, Daiber A, Kleschyov AL, Mulsch A, Sydow K, Schulz E, Chen K, Keaney JF Jr, Lassegue B, Walter U, Griendling KK, Munzel T. Differential effects of diabetes on the expression of the gp91phox homologues nox1 and nox4. *Free Radic Biol Med*. 2005;39:381–391.
173. San Martin A, Du P, Dikalova A, Lassegue B, Aleman M, Gongora MC, Brown K, Joseph G, Harrison DG, Taylor WR, Jo H, Griendling KK. Reactive oxygen species-selective regulation of aortic inflammatory gene expression in type 2 diabetes. *Am J Physiol Heart Circ Physiol*. 2007;292:H2073–H2082.
174. Castier Y, Brandes RP, Leseche G, Tedgui A, Lehoux S. P47phox-dependent NADPH oxidase regulates flow-induced vascular remodeling. *Circ Res*. 2005;97:533–540.
175. Garrido-Urbani S, Jemelin S, Deffert C, Carnesecchi S, Basset O, Szyndralewicz C, Heitz F, Page P, Montet X, Michalik L, Arbiser J, Ruegg C, Krause KH, Imhof BA. Targeting vascular NADPH oxidase 1 blocks tumor angiogenesis through a pparalpha mediated mechanism. *PLoS One*. 2011;6:e14665.
176. Lassegue B, Sorescu D, Szocs K, Yin Q, Akers M, Zhang Y, Grant SL, Lambeth JD, Griendling KK. Novel gp91(phox) homologues in vascular smooth muscle cells: Nox1 mediates angiotensin II-induced superoxide formation and redox-sensitive signaling pathways. *Circ Res*. 2001;88:888–894.
177. Wang X, Sun Z. Thyroid hormone induces artery smooth muscle cell proliferation: discovery of a new tralpa1-nox1 pathway. *J Cell Mol Med*. 2010;14:368–380.
178. Menshikov M, Plekhanova O, Cai H, Chalupsky K, Parfyonova Y, Bashtrikov P, Tkachuk V, Berk BC. Urokinase plasminogen activator stimulates vascular smooth muscle cell proliferation via redox-dependent pathways. *Arterioscler Thromb Vasc Biol*. 2006;26:801–807.
179. Katsuyama M, Fan C, Yabe-Nishimura C. NADPH oxidase is involved in prostaglandin f2alpha-induced hypertrophy of vascular smooth muscle cells: Induction of nox1 by pgf2alpha. *J Biol Chem*. 2002;277:13438–13442.
180. Yin CC, Huang KT. H2O2 but not O2- elevated by oxidized LDL enhances human aortic smooth muscle cell proliferation. *J Biomed Sci*. 2007;14:245–254.
181. Suh YA, Arnold RS, Lassegue B, Shi J, Xu X, Sorescu D, Chung AB, Griendling KK, Lambeth JD. Cell transformation by the superoxide-generating oxidase mox1. *Nature*. 1999;401:79–82.
182. Stanic B, Katsuyama M, Miller FJ Jr. An oxidized extracellular oxidation-reduction state increases nox1 expression and proliferation in vascular smooth muscle cells via epidermal growth factor receptor activation. *Arterioscler Thromb Vasc Biol*. 2010;30:2234–2241.
183. Dammanahalli JK, Sun Z. Endothelin (ET)-1 inhibits nicotinamide adenine dinucleotide phosphate oxidase activity in human abdominal aortic endothelial cells: a novel function of etb1 receptors. *Endocrinology*. 2008;149:4979–4987.
184. Ushio-Fukai M, Alexander RW, Akers M, Griendling KK. P38map kinase is a critical component of the redox-sensitive signaling pathways by angiotensin II: role in vascular smooth muscle cell hypertrophy. *J Biol Chem*. 1998;273:15022–15029.
185. Ushio-Fukai M, Alexander RW, Akers M, Yin Q, Fujio Y, Walsh K, Griendling KK. Reactive oxygen species mediate the activation of akt/protein kinase B by angiotensin II in vascular smooth muscle cells. *J Biol Chem*. 1999;274:22699–22704.
186. Ranjan P, Anathy V, Burch PM, Weirather K, Lambeth JD, Heintz NH. Redox-dependent expression of cyclin d1 and cell proliferation by nox1 in mouse lung epithelial cells. *Antioxidant Redox Signal*. 2006;8:1447–1459.
187. Sancho P, Bertran E, Caja L, Carmona-Cuenca I, Murillo MM, Fabregat I. The inhibition of the epidermal growth factor (EGF) pathway enhances TGF-beta-induced apoptosis in rat hepatoma cells through inducing oxidative stress coincident with a change in the expression pattern of the NADPH oxidases (NOX) isoforms. *Biochim Biophys Acta*. 2009;1793:253–263.
188. Kim YS, Morgan MJ, Choksi S, Liu ZG. Tnf-induced activation of the nox1 NADPH oxidase and its role in the induction of necrotic cell death. *Mol Cell*. 2007;26:675–687.
189. Schilder YD, Heiss EH, Schachner D, Ziegler J, Reznicek G, Sorescu D, Dirsch VM. NADPH oxidases 1 and 4 mediate cellular senescence induced by resveratrol in human endothelial cells. *Free Radic Biol Med*. 2009;46:1598–1606.
190. Zimmerman MC, Takapoo M, Jagadeesha DK, Stanic B, Banfi B, Bhalla RC, Miller FJ Jr. Activation of NADPH oxidase 1 increases intracellular calcium and migration of smooth muscle cells. *Hypertension*. 2011;58:446–453.
191. Schroder K, Helmcke I, Palfi K, Krause KH, Busse R, Brandes RP. Nox1 mediates basic fibroblast growth factor-induced migration of vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2007;27:1736–1743.
192. Maheswaranathan M, Gole HK, Fernandez I, Lassegue B, Griendling KK, San Martin A. Platelet-derived growth factor (PDGF) regulates slingshot phosphatase activity via nox1-dependent auto-dephosphorylation of serine 834 in vascular smooth muscle cells. *J Biol Chem*. 2011;286:35430–35437.
193. San Martin A, Lee MY, Williams HC, Mizuno K, Lassegue B, Griendling KK. Dual regulation of cofilin activity by lim kinase and slingshot-11 phosphatase controls platelet-derived growth factor-induced migration of human aortic smooth muscle cells. *Circ Res*. 2008;102:432–438.
194. Shinohara M, Adachi Y, Mitsushita J, Kuwabara M, Nagasawa A, Harada S, Furuta S, Zhang Y, Seheli K, Miyazaki H, Kamata T. Reactive oxygen generated by NADPH oxidase 1 (nox1) contributes to cell invasion by regulating matrix metalloproteinase-9 production and cell migration. *J Biol Chem*. 2010;285:4481–4488.
195. Haloui M, Meilhac O, Jandrot-Perrus M, Michel JB. Atorvastatin limits the pro-inflammatory response of rat aortic smooth muscle cells to thrombin. *Eur J Pharmacol*. 2003;474:175–184.
196. Ungvari Z, Csiszar A, Edwards JG, Kaminski PM, Wolin MS, Kaley G, Koller A. Increased superoxide production in coronary arteries in hyperhomocysteinemia: role of tumor necrosis factor-alpha, NAD(P)H oxidase, and inducible nitric oxide synthase. *Arterioscler Thromb Vasc Biol*. 2003;23:418–424.
197. Lee JG, Lim EJ, Park DW, Lee SH, Kim JR, Baek SH. A combination of lox-1 and nox1 regulates tlr9-mediated foam cell formation. *Cell Signal*. 2008;20:2266–2275.
198. Duerschmidt N, Stielow C, Muller G, Pagano PJ, Morawietz H. NO-mediated regulation of NAD(P)H oxidase by laminar shear stress in human endothelial cells. *J Physiol*. 2006;576:557–567.
199. Furst R, Brueckl C, Kuebler WM, Zahler S, Krotz F, Gorchach A, Vollmar AM, Kiemer AK. Atrial natriuretic peptide induces mitogen-activated protein kinase phosphatase-1 in human endothelial cells via rac1 and NAD(P)H oxidase/nox2-activation. *Circ Res*. 2005;96:43–53.
200. Li H, Hortmann M, Daiber A, Oelze M, Ostad MA, Schwarz PM, Xu H, Xia N, Kleschyov AL, Mang C, Warnholtz A, Munzel T, Forstermann U. Cyclooxygenase 2-selective and nonselective nonsteroidal anti-inflammatory drugs induce oxidative stress by up-regulating vascular NADPH oxidases. *J Pharmacol Exp Ther*. 2008;326:745–753.
201. Schuhmacher S, Foretz M, Knorr M, Jansen T, Hortmann M, Wenzel P, Oelze M, Kleschyov AL, Daiber A, Keaney JF Jr, Wegener G, Lackner K, Munzel T, Viollot B, Schulz E. Alpha1amp-activated protein kinase preserves endothelial function during chronic angiotensin II treatment by limiting nox2 upregulation. *Arterioscler Thromb Vasc Biol*. 2011;31:560–566.
202. Thakur S, Du J, Hourani S, Ledent C, Li JM. Inactivation of adenosine a2a receptor attenuates basal and angiotensin II-induced ros production by nox2 in endothelial cells. *J Biol Chem*. 2010;285:40104–40113.
203. Takenouchi Y, Kobayashi T, Matsumoto T, Kamata K. Gender differences in age-related endothelial function in the murine aorta. *Atherosclerosis*. 2009;206:397–404.
204. Zemse SM, Hilgers RH, Webb RC. Interleukin-10 counteracts impaired endothelium-dependent relaxation induced by ang II in murine aortic rings. *Am J Physiol Heart Circ Physiol*. 2007;292:H3103–H3108.

205. Violi F, Sanguigni V, Carnevale R, et al. Hereditary deficiency of gp91(phox) is associated with enhanced arterial dilatation: results of a multicenter study. *Circulation*. 2009;120:1616–1622.
206. Taye A, Saad AH, Kumar AH, Morawietz H. Effect of apocynin on NADPH oxidase-mediated oxidative stress-lox-1-eNOS pathway in human endothelial cells exposed to high glucose. *Eur J Pharmacol*. 2010;627:42–48.
207. Hwang J, Saha A, Boo YC, Sorescu GP, McNally JS, Holland SM, Dikalov S, Giddens DP, Griendling KK, Harrison DG, Jo H. Oscillatory shear stress stimulates endothelial production of O₂⁻ from p47phox-dependent NAD(P)H oxidases, leading to monocyte adhesion. *J Biol Chem*. 2003;278:47291–47298.
208. Cathcart MK. Regulation of superoxide anion production by NADPH oxidase in monocytes/macrophages: contributions to atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2004;24:23–28.
209. Pietrowski E, Bender B, Huppert J, White R, Luhmann HJ, Kuhlmann CR. Pro-inflammatory effects of interleukin-17a on vascular smooth muscle cells involve NAD(P)H-oxidase derived reactive oxygen species. *J Vasc Res*. 2011;48:52–58.
210. McNally JS, Davis ME, Giddens DP, Saha A, Hwang J, Dikalov S, Jo H, Harrison DG. Role of xanthine oxidoreductase and NAD(P)H oxidase in endothelial superoxide production in response to oscillatory shear stress. *Am J Physiol Heart Circ Physiol*. 2003;285:H2290–H2297.
211. Landmesser U, Dikalov S, Price SR, McCann L, Fukai T, Holland SM, Mitch WE, Harrison DG. Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J Clin Invest*. 2003;111:1201–1209.
212. Weaver M, Liu J, Pimentel D, Reddy DJ, Harding P, Peterson EL, Pagano PJ. Adventitial delivery of dominant-negative p67phox attenuates neointimal hyperplasia of the rat carotid artery. *Am J Physiol Heart Circ Physiol*. 2006;290:H1933–H1941.
213. Peshavariya H, Dusting GJ, Jiang F, Halmos LR, Sobey CG, Drummond GR, Selemidis S. NADPH oxidase isoform selective regulation of endothelial cell proliferation and survival. *Naunyn Schmiedebergs Arch Pharmacol*. 2009;380:193–204.
214. Li JM, Fan LM, George VT, Brooks G. Nox2 regulates endothelial cell cycle arrest and apoptosis via p21cip1 and p53. *Free Radic Biol Med*. 2007;43:976–986.
215. Imanishi T, Kobayashi K, Kuroi A, Ikejima H, Akasaka T. Pioglitazone inhibits angiotensin II-induced senescence of endothelial progenitor cell. *Hypertens Res*. 2008;31:757–765.
216. Ikeda S, Yamaoka-Tojo M, Hilenski L, Patrushev NA, Anwar GM, Quinn MT, Ushio-Fukai M. Iqgap1 regulates reactive oxygen species-dependent endothelial cell migration through interacting with nox2. *Arterioscler Thromb Vasc Biol*. 2005;25:2295–2300.
217. Pendyala S, Gorshkova IA, Usatyuk PV, He D, Pennathur A, Lambeth JD, Thannickal VJ, Natarajan V. Role of nox4 and nox2 in hyperoxia-induced reactive oxygen species generation and migration of human lung endothelial cells. *Antioxidant Redox Signal*. 2009;11:747–764.
218. Kim SY, Lee JG, Cho WS, Cho KH, Sakong J, Kim JR, Chin BR, Baek SH. Role of NADPH oxidase-2 in lipopolysaccharide-induced matrix metalloproteinase expression and cell migration. *Immunol Cell Biol*. 2010;88:197–204.
219. Schroder K, Kohnen A, Aicher A, Liehn EA, Buchse T, Stein S, Weber C, Dimmeler S, Brandes RP. NADPH oxidase nox2 is required for hypoxia-induced mobilization of endothelial progenitor cells. *Circ Res*. 2009;105:537–544.
220. Looi YH, Grieve DJ, Siva A, Walker SJ, Anilkumar N, Cave AC, Marber M, Monaghan MJ, Shah AM. Involvement of nox2 NADPH oxidase in adverse cardiac remodeling after myocardial infarction. *Hypertension*. 2008;51:319–325.
221. Wang M, Zhang J, Walker SJ, Dworakowski R, Lakatta EG, Shah AM. Involvement of NADPH oxidase in age-associated cardiac remodeling. *J Mol Cell Cardiol*. 2010;48:765–772.
222. Hingtgen SD, Tian X, Yang J, Dunlay SM, Peek AS, Wu Y, Sharma RV, Engelhardt JF, Davison RL. Nox2-containing NADPH oxidase and akt activation play a key role in angiotensin II-induced cardiomyocyte hypertrophy. *Physiol Genome*. 2006;26:180–191.
223. Shanmugam P, Valente AJ, Prabhu SD, Venkatesan B, Yoshida T, Delafontaine P, Chandrasekar B. Angiotensin-II type 1 receptor and nox2 mediate tcf/lef and creb dependent wisp1 induction and cardiomyocyte hypertrophy. *J Mol Cell Cardiol*. 2011;50:928–938.
224. Stas S, Whaley-Connell A, Habibi J, Apeesh L, Hayden MR, Karuparthi PR, Qazi M, Morris EM, Cooper SA, Link CD, Stump C, Hay M, Ferrario C, Sowers JR. Mineralocorticoid receptor blockade attenuates chronic overexpression of the renin-angiotensin-aldosterone system stimulation of reduced nicotinamide adenine dinucleotide phosphate oxidase and cardiac remodeling. *Endocrinology*. 2007;148:3773–3780.
225. Wilson P, Morgan J, Funder JW, Fuller PJ, Young MJ. Mediators of mineralocorticoid receptor-induced profibrotic inflammatory responses in the heart. *Clin Sci*. 2009;116:731–739.
226. Nakamura T, Kataoka K, Fukuda M, Nako H, Tokutomi Y, Dong YF, Ichijo H, Ogawa H, Kim-Mitsuyama S. Critical role of apoptosis signal-regulating kinase 1 in aldosterone/salt-induced cardiac inflammation and fibrosis. *Hypertension*. 2009;54:544–551.
227. Guzik TJ, Sadowski J, Kapelak B, Jopek A, Rudzinski P, Pillai R, Korbut R, Channon KM. Systemic regulation of vascular NAD(P)H oxidase activity and nox isoform expression in human arteries and veins. *Arterioscler Thromb Vasc Biol*. 2004;24:1614–1620.
228. Miller AA, Drummond GR, Schmidt HH, Sobey CG. NADPH oxidase activity and function are profoundly greater in cerebral versus systemic arteries. *Circ Res*. 2005;97:1055–1062.
229. Clempus RE, Sorescu D, Dikalova AE, Pounkova L, Jo P, Sorescu GP, Schmidt HH, Lassegue B, Griendling KK. Nox4 is required for maintenance of the differentiated vascular smooth muscle cell phenotype. *Arterioscler Thromb Vasc Biol*. 2007;27:42–48.
230. Ellmark SH, Dusting GJ, Fui MN, Guzzo-Pernell N, Drummond GR. The contribution of nox4 to NADPH oxidase activity in mouse vascular smooth muscle. *Cardiovasc Res*. 2005;65:495–504.
231. Goettsch C, Goettsch W, Arsov A, Hofbauer LC, Bornstein SR, Morawietz H. Long-term cyclic strain downregulates endothelial nox4. *Antioxid Redox Signal*. 2009;11:2385–2397.
232. Hwang J, Ing MH, Salazar A, Lassegue B, Griendling K, Navab M, Sevanian A, Hsiai TK. Pulsatile versus oscillatory shear stress regulates NADPH oxidase subunit expression: implication for native LDL oxidation. *Circ Res*. 2003;93:1225–1232.
233. Peshavariya H, Jiang F, Taylor CJ, Selemidis S, Chang CW, Dusting G. Translation-linked mRNA destabilization accompanying serum-induced nox4 expression in human endothelial cells. *Antioxidant Redox Signal*. 2009;11:2399–2408.
234. Haurani MJ, Cifuentes ME, Shepard AD, Pagano PJ. Nox4 oxidase overexpression specifically decreases endogenous nox4 mRNA and inhibits angiotensin II-induced adventitial myofibroblast migration. *Hypertension*. 2008;52:143–149.
235. Sturrock A, Cahill B, Norman K, Huecksteadt TP, Hill K, Sanders K, Karwande SV, Stringham JC, Bull DA, Gleich M, Kennedy TP, Hoidal JR. Transforming growth factor-beta1 induces nox4 NAD(P)H oxidase and reactive oxygen species-dependent proliferation in human pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol*. 2006;290:L661–L673.
236. Cucoranu I, Clempus R, Dikalova A, Phelan PJ, Ariyan S, Dikalov S, Sorescu D. NAD(P)H oxidase 4 mediates transforming growth factor-beta1-induced differentiation of cardiac fibroblasts into myofibroblasts. *Circ Res*. 2005;97:900–907.
237. Martin-Garrido A, Brown DI, Lyle AN, Dikalova A, Seidel-Rogol B, Lassegue B, San Martin A, Griendling KK. NADPH oxidase 4 mediates tgf-beta-induced smooth muscle alpha-actin via p38mapk and serum response factor. *Free Radic Biol Med*. 2011;50:354–362.
238. Ismail S, Sturrock A, Wu P, Cahill B, Norman K, Huecksteadt T, Sanders K, Kennedy T, Hoidal J. Nox4 mediates hypoxia-induced proliferation of human pulmonary artery smooth muscle cells: the role of autocrine production of transforming growth factor- β 1 and insulin-like growth factor binding protein-3. *Am J Physiol Lung Cell Mol Physiol*. 2009;296:L489–L499.
239. Amara N, Goven D, Prost F, Muloway R, Crestani B, Boczkowski J. Nox4/NADPH oxidase expression is increased in pulmonary fibroblasts from patients with idiopathic pulmonary fibrosis and mediates tgf β 1-induced fibroblast differentiation into myofibroblasts. *Thorax*. 2010;65:733–738.
240. Yang S, Zhang Y, Ries W, Key L. Expression of nox4 in osteoclasts. *J Cell Biochem*. 2004;92:238–248.
241. Li J, Stouffs M, Serrander L, Banfi B, Bettiol E, Charnay Y, Steger K, Krause KH, Jaconi ME. The NADPH oxidase nox4 drives cardiac differentiation: role in regulating cardiac transcription factors and map kinase activation. *Mol Biol Cell*. 2006;17:3978–3988.
242. Pedruzzi E, Guichard C, Ollivier V, Driss F, Fay M, Prunet C, Marie JC, Pouzet C, Samadi M, Elbim C, O'Dowd J, Bens M, Vandewalle A, Gougerot-Pocidallo MA, Lizard G, Ogier-Denis E. NAD(P)H oxidase nox-4 mediates 7-ketocholesterol-induced endoplasmic reticulum stress

- and apoptosis in human aortic smooth muscle cells. *Mol Cell Biol*. 2004;24:10703–10717.
243. Gorin Y, Block K, Hernandez J, Bhandari B, Wagner B, Barnes JL, Abboud HE. Nox4 NAD(P)H oxidase mediates hypertrophy and fibronectin expression in the diabetic kidney. *J Biol Chem*. 2005;280:39616–39626.
 244. Lee YM, Kim BJ, Chun YS, So I, Choi H, Kim MS, Park JW. Nox4 as an oxygen sensor to regulate task-1 activity. *Cell Signal*. 2006;18:499–507.
 245. Sun QA, Hess DT, Nogueira L, Yong S, Bowles DE, Eu J, Laurita KR, Meissner G, Stamler JS. Oxygen-coupled redox regulation of the skeletal muscle ryanodine receptor-Ca²⁺ release channel by NADPH oxidase 4. *Proc Natl Acad Sci U S A*. 2011;108:16098–16103.
 246. Szocs K, Lassegue B, Sorescu D, Hlenski LL, Valppu L, Couse TL, Wilcox JN, Quinn MT, Lambeth JD, Griendling KK. Upregulation of nox-based NAD(P)H oxidases in restenosis after carotid injury. *Arterioscler Thromb Vasc Biol*. 2002;22:21–27.
 247. Sorescu D, Weiss D, Lassegue B, Clempus RE, Szocs K, Sorescu GP, Valppu L, Quinn MT, Lambeth JD, Vega JD, Taylor WR, Griendling KK. Superoxide production and expression of nox family proteins in human atherosclerosis. *Circulation*. 2002;105:1429–1435.
 248. Perrotta I, Sciangula A, Perrotta E, Donato G, Cassese M. Ultrastructural analysis and electron microscopic localization of nox4 in healthy and atherosclerotic human aorta. *Ultrastruct Pathol*. 2011;35:1–6.
 249. Guo SJ, Wu LY, Shen WL, Chen WD, Wei J, Gao PJ, Zhu DL. Gene profile for differentiation of vascular adventitial myofibroblasts. *Sheng li xue bao: [Acta physiologica Sinica]*. 2006;58:337–344.
 250. Mouche S, Mkaddem SB, Wang W, Katic M, Tseng YH, Carnesecchi S, Steger K, Foti M, Meier CA, Muzzin P, Kahn CR, Ogier-Denis E, Szanto I. Reduced expression of the NADPH oxidase nox4 is a hallmark of adipocyte differentiation. *Biochim Biophys Acta*. 2007;1773:1015–1027.
 251. Xiao Q, Luo Z, Pepe AE, Margariti A, Zeng L, Xu Q. Embryonic stem cell differentiation into smooth muscle cells is mediated by nox4-produced h2o2. *Am J Physiol Cell Physiol*. 2009;296:C711–C723.
 252. Datla SR, Peshavariya H, Dusting GJ, Mahadev K, Goldstein BJ, Jiang F. Important role of nox4 type NADPH oxidase in angiogenic responses in human microvascular endothelial cells in vitro. *Arterioscler Thromb Vasc Biol*. 2007;27:2319–2324.
 253. Tong X, Hou X, Jourdeuil D, Weisbrod RM, Cohen RA. Upregulation of nox4 by TGFβ1 oxidizes serca and inhibits no in arterial smooth muscle of the prediabetic Zucker rat. *Circ Res*. 2010;107:975–983.
 254. Craig SM, Chen K, Pei Y, Li C, Huang X, Chen C, Shibata R, Sato K, Walsh K, Keane JF Jr. NADPH oxidase 4 promotes endothelial angiogenesis through endothelial nitric oxide synthase activation. *Circulation*. 2011;124:731–740.
 255. Chen K, Kirber MT, Xiao H, Yang Y, Keane JF Jr. Regulation of ros signal transduction by NADPH oxidase 4 localization. *J Cell Biol*. 2008;181:1129–1139.
 256. Li S, Tabar SS, Malec V, Eul BG, Klepetko W, Weissmann N, Grimmering F, Seeger W, Rose F, Hanzel J. Nox4 regulates ros levels under normoxic and hypoxic conditions, triggers proliferation, and inhibits apoptosis in pulmonary artery adventitial fibroblasts. *Antioxidant Redox Signal*. 2008;10:1687–1698.
 257. Basuroy S, Bhattacharya S, Leffler CW, Parfenova H. Nox4 NADPH oxidase mediates oxidative stress and apoptosis caused by tnf-alpha in cerebral vascular endothelial cells. *Am J Physiol Cell Physiol*. 2009;296:C422–C432.
 258. Tian XY, Yung LH, Wong WT, Liu J, Leung FP, Liu L, Chen Y, Kong SK, Kwan KM, Ng SM, Lai PB, Yung LM, Yao X, Huang Y. Bone morphogenic protein-4 induces endothelial cell apoptosis through oxidative stress-dependent p38mapk and jnk pathway. *J Mol Cell Cardiol*. 2012;52:237–244.
 259. Senturk S, Mumcuoglu M, Gursoy-Yuzugullu O, Cingoz B, Akcali KC, Ozturk M. Transforming growth factor-beta induces senescence in hepatocellular carcinoma cells and inhibits tumor growth. *Hepatology*. 2010;52:966–974.
 260. Weyemi U, Lagente-Chevallier O, Boufraquech M, Prenois F, Courtin F, Caillou B, Talbot M, Dardalhon M, Al Ghuzlan A, Bidart JM, Schlumberger M, Dupuy C. ROS-generating NADPH oxidase nox4 is a critical mediator in oncogenic h-ras-induced DNA damage and subsequent senescence. *Oncogene*. 2012;31:1117–1129.
 261. Gomes P, Simao S, Silva E, Pinto V, Amaral JS, Afonso J, Serrao MP, Pinho MJ, Soares-da-Silva P. Aging increases oxidative stress and renal expression of oxidant and antioxidant enzymes that are associated with an increased trend in systolic blood pressure. *Oxid Med Cell Longev*. 2009;2:138–145.
 262. Li L, Smith A, Hagen TM, Frei B. Vascular oxidative stress and inflammation increase with age: ameliorating effects of alpha-lipoic acid supplementation. *Ann N Y Acad Sci*. 2010;1203:151–159.
 263. McCrann DJ, Yang D, Chen H, Carroll S, Ravid K. Upregulation of nox4 in the aging vasculature and its association with smooth muscle cell polyploidy. *Cell Cycle*. 2009;8:902–908.
 264. Lener B, Koziel R, Pircher H, Hutter E, Greussing R, Herndl-Brandstetter D, Hermann M, Unterluggauer H, Jansen-Durr P. The NADPH oxidase nox4 restricts the replicative lifespan of human endothelial cells. *Biochem J*. 2009;423:363–374.
 265. Li H, Liu Q, Wang N, Xu J. Correlation of different NADPH oxidase homologues with late endothelial progenitor cell senescence induced by angiotensin II: effect of telmisartan. *Intern Med*. 2011;50:1631–1642.
 266. Vasa-Nicotera M, Chen H, Tucci P, Yang AL, Saintigny G, Menghini R, Mahe C, Agostini M, Knight RA, Melino G, Federici M. Mir-146a is modulated in human endothelial cell with aging. *Atherosclerosis*. 2011;217:326–330.
 267. Rueckschloss U, Villmow M, Klockner U. NADPH oxidase-derived superoxide impairs calcium transients and contraction in aged murine ventricular myocytes. *Exp Gerontol*. 2010;45:788–796.
 268. Jaulmes A, Sansilvestri-Morel P, Rolland-Valognes G, Bernhardt F, Gaertner R, Lockhart BP, Cordi A, Wierzbicki M, Rupin A, Verbeuren TJ. Nox4 mediates the expression of plasminogen activator inhibitor-1 via p38 mapk pathway in cultured human endothelial cells. *Thromb Res*. 2009;124:439–446.
 269. Maloney E, Sweet IR, Hockenbery DM, Pham M, Rizzo NO, Tateya S, Handa P, Schwartz MW, Kim F. Activation of nf-kappab by palmitate in endothelial cells: a key role for NADPH oxidase-derived superoxide in response to tlr4 activation. *Arterioscler Thromb Vasc Biol*. 2009;29:1370–1375.
 270. Moe KT, Yin NO, Naylynn TM, Khairunnisa K, Wutyi MA, Gu Y, Atan MS, Wong MC, Koh TH, Wong P. Nox2 and nox4 mediate tumour necrosis factor-alpha-induced ventricular remodelling in mice. *J Cell Mol Med*. 2011;15:2601–2613.
 271. Ryu J, Lee CW, Shin JA, Park CS, Kim JJ, Park SJ, Han KH. Fcγmariia mediates c-reactive protein-induced inflammatory responses of human vascular smooth muscle cells by activating NADPH oxidase 4. *Cardiovasc Res*. 2007;75:555–565.
 272. Ritsick DR, Edens WA, Finnerty V, Lambeth JD. Nox regulation of smooth muscle contraction. *Free Radic Biol Med*. 2007;43:31–38.
 273. Zhang Q, Malik P, Pandey D, Gupta S, Jagnandan D, Belin de Chantemele E, Banfi B, Marrero MB, Rudic RD, Stepp DW, Fulton DJ. Paradoxical activation of endothelial nitric oxide synthase by NADPH oxidase. *Arterioscler Thromb Vasc Biol*. 2008;28:1627–1633.
 274. Nakazono K, Watanabe N, Matsuno K, Sasaki J, Sato T, Inoue M. Does superoxide underlie the pathogenesis of hypertension? *Proc Natl Acad Sci U S A*. 1991;88:10045–10048.
 275. Lassegue B, Griendling KK. NADPH oxidases: functions and pathologies in the vasculature. *Arterioscler Thromb Vasc Biol*. 2010;30:653–661.
 276. Sirker A, Zhang M, Shah AM. NADPH oxidases in cardiovascular disease: insights from in vivo models and clinical studies. *Basic Res Cardiol*. 2011;106:735–747.
 277. Zalba G, San Jose G, Moreno MU, Fortuno A, Diez J. NADPH oxidase-mediated oxidative stress: genetic studies of the p22(phox) gene in hypertension. *Antioxidant Redox Signal*. 2005;7:1327–1336.
 278. Winkler K, Wunsch S, Kreutz R, Rothermund L, Paul M, Schmidt HH. Upregulation of the vascular NAD(P)H-oxidase isoforms nox1 and nox4 by the renin-angiotensin system in vitro and in vivo. *Free Radic Biol Med*. 2001;31:1456–1464.
 279. Wang P, Tang F, Li R, Zhang H, Chen S, Liu P, Huang H. Contribution of different nox homologues to cardiac remodeling in two-kidney two-clip renovascular hypertensive rats: effect of valsartan. *Pharmacol Res*. 2007;55:408–417.
 280. Nishiyama A, Yoshizumi M, Rahman M, Kobori H, Seth DM, Miyatake A, Zhang GX, Yao L, Hitomi H, Shokoji T, Kiyomoto H, Kimura S, Tamaki T, Kohno M, Abe Y. Effects of at1 receptor blockade on renal injury and mitogen-activated protein activity in Dahl salt-sensitive rats. *Kidney Int*. 2004;65:972–981.
 281. Mollnau H, Wendt M, Szocs K, et al. Effects of angiotensin II infusion on the expression and function of NAD(P)H oxidase and components of nitric oxide/cgmp signaling. *Circ Res*. 2002;90:E58–E65.

282. Matsuno K, Yamada H, Iwata K, Jin D, Katsuyama M, Matsuki M, Takai S, Yamanishi K, Miyazaki M, Matsubara H, Yabe-Nishimura C. Nox1 is involved in angiotensin II-mediated hypertension: A study in nox1-deficient mice. *Circulation*. 2005;112:2677–2685.
283. Dikalova A, Clempus R, Lassegue B, Cheng G, McCoy J, Dikalov S, San Martin A, Lyle A, Weber DS, Weiss D, Taylor WR, Schmidt HH, Owens GK, Lambeth JD, Griendling KK. Nox1 overexpression potentiates angiotensin II-induced hypertension and vascular smooth muscle hypertrophy in transgenic mice. *Circulation*. 2005;112:2668–2676.
284. Dikalova AE, Gongora MC, Harrison DG, Lambeth JD, Dikalov S, Griendling KK. Upregulation of nox1 in vascular smooth muscle leads to impaired endothelium-dependent relaxation via enos uncoupling. *Am J Physiol Heart Circ Physiol*. 2010;299:H673–H679.
285. Yogi A, Mercure C, Touyz J, Callera GE, Montezano AC, Aranha AB, Tostes RC, Reudelhuber T, Touyz RM. Renal redox-sensitive signaling, but not blood pressure, is attenuated by nox1 knockout in angiotensin II-dependent chronic hypertension. *Hypertension*. 2008;51:500–506.
286. Wind S, Beuerlein K, Armitage ME, Taye A, Kumar AH, Janowitz D, Neff C, Shah AM, Wingle K, Schmidt HH. Oxidative stress and endothelial dysfunction in aortas of aged spontaneously hypertensive rats by nox1/2 is reversed by NADPH oxidase inhibition. *Hypertension*. 2010;56:490–497.
287. Datla SR, Griendling KK. Reactive oxygen species, NADPH oxidases, and hypertension. *Hypertension*. 2010;56:325–330.
288. Cook-Mills JM. Reactive oxygen species regulation of immune function. *Mol Immunol*. 2002;39:497–498.
289. Cifuentes ME, Rey FE, Carretero OA, Pagano PJ. Upregulation of p67(phox) and gp91(phox) in aortas from angiotensin II-infused mice. *Am J Physiol Heart Circ Physiol*. 2000;279:H2234–H2240.
290. Wang HD, Xu S, Johns DG, Du Y, Quinn MT, Cayatte AJ, Cohen RA. Role of NADPH oxidase in the vascular hypertrophic and oxidative stress response to angiotensin II in mice. *Circ Res*. 2001;88:947–953.
291. Landmesser U, Cai H, Dikalov S, McCann L, Hwang J, Jo H, Holland SM, Harrison DG. Role of p47(phox) in vascular oxidative stress and hypertension caused by angiotensin II. *Hypertension*. 2002;40:511–515.
292. Rey FE, Cifuentes ME, Kiarash A, Quinn MT, Pagano PJ. Novel competitive inhibitor of NAD(P)H oxidase assembly attenuates vascular O(2)(-) and systolic blood pressure in mice. *Circ Res*. 2001;89:408–414.
293. Zhou MS, Hernandez Schulman I, Pagano PJ, Jaimes EA, Raj L. Reduced NAD(P)H oxidase in low renin hypertension: link among angiotensin II, atherogenesis, and blood pressure. *Hypertension*. 2006;47:81–86.
294. Weber DS, Rocic P, Mellis AM, Laude K, Lyle AN, Harrison DG, Griendling KK. Angiotensin II-induced hypertrophy is potentiated in mice overexpressing p22phox in vascular smooth muscle. *Am J Physiol Heart Circ Physiol*. 2005;288:H37–H42.
295. Murdoch CE, Alom-Ruiz SP, Wang M, Zhang M, Walker S, Yu B, Brewer A, Shah AM. Role of endothelial nox2 NADPH oxidase in angiotensin II-induced hypertension and vasomotor dysfunction. *Basic Res Cardiol*. 2011;106:527–538.
296. Zimmerman MC, Lazartigues E, Sharma RV, Davisson RL. Hypertension caused by angiotensin II infusion involves increased superoxide production in the central nervous system. *Circ Res*. 2004;95:210–216.
297. Peterson JR, Burmeister MA, Tian X, Zhou Y, Guraju MR, Stupinski JA, Sharma RV, Davisson RL. Genetic silencing of nox2 and nox4 reveals differential roles of these NADPH oxidase homologues in the vasopressor and dipsogenic effects of brain angiotensin II. *Hypertension*. 2009;54:1106–1114.
298. Modlinger P, Chabrashvili T, Gill PS, Mendonca M, Harrison DG, Griendling KK, Li M, Raggio J, Wellstein A, Chen Y, Welch WJ, Wilcox CS. Rna silencing in vivo reveals role of p22phox in rat angiotensin slow pressor response. *Hypertension*. 2006;47:238–244.
299. Guzik TJ, Hoch NE, Brown KA, McCann LA, Rahman A, Dikalov S, Goronzy J, Weyand C, Harrison DG. Role of the T cell in the genesis of angiotensin II induced hypertension and vascular dysfunction. *J Exp Med*. 2007;204:2449–2460.
300. Senchenkova EY, Russell J, Kurmaeva E, Ostanin D, Granger DN. Role of T lymphocytes in angiotensin II-mediated microvascular thrombosis. *Hypertension*. 2011;58:959–965.
301. Akasaki T, Ohya Y, Kuroda J, Eto K, Abe I, Sumimoto H, Iida M. Increased expression of gp91phox homologues of NAD(P)H oxidase in the aortic media during chronic hypertension: involvement of the renin-angiotensin system. *Hypertens Res*. 2006;29:813–820.
302. Ray R, Murdoch CE, Wang M, Santos CX, Zhang M, Alom-Ruiz S, Anilkumar N, Ouattara A, Cave AC, Walker SJ, Grieve DJ, Charles RL, Eaton P, Brewer AC, Shah AM. Endothelial nox4 NADPH oxidase enhances vasodilatation and reduces blood pressure in vivo. *Arterioscler Thromb Vasc Biol*. 2011;31:1368–1376.
303. Paravicini TM, Chrissobolis S, Drummond GR, Sobey CG. Increased NADPH-oxidase activity and nox4 expression during chronic hypertension is associated with enhanced cerebral vasodilatation to NADPH in vivo. *Stroke*. 2004;35:584–589.
304. Chabrashvili T, Tojo A, Onozato ML, Kitiyakara C, Quinn MT, Fujita T, Welch WJ, Wilcox CS. Expression and cellular localization of classic NADPH oxidase subunits in the spontaneously hypertensive rat kidney. *Hypertension*. 2002;39:269–274.
305. Nishiyama A, Yao L, Nagai Y, Miyata K, Yoshizumi M, Kagami S, Kondo S, Kiyomoto H, Shokoji T, Kimura S, Kohno M, Abe Y. Possible contributions of reactive oxygen species and mitogen-activated protein kinase to renal injury in aldosterone/salt-induced hypertensive rats. *Hypertension*. 2004;43:841–848.
306. Schmidt H, Hermans J, Schiffrers P, Kleikers P, Radermacher K, Altenhofer S, Wingle K. NADPH oxidase as a pharmacological target in disease-relevant oxidative stress. *Acta Physiologica*. 2011;203(Suppl 687):O-22.
307. Gavazzi G, Deffert C, Trocme C, Schappi M, Herrmann FR, Krause KH. Nox1 deficiency protects from aortic dissection in response to angiotensin II. *Hypertension*. 2007;50:189–196.
308. Sheehan AL, Carrell S, Johnson B, Stanic B, Banfi B, Miller FJ Jr. Role for nox1 NADPH oxidase in atherosclerosis. *Atherosclerosis*. 2011;216:321–326.
309. Vendrov AE, Madamanchi NR, Niu XL, Molnar KC, Runge M, Szyn-dralewicz C, Page P, Runge MS. NADPH oxidases regulate cd44 and hyaluronic acid expression in thrombin-treated vascular smooth muscle cells and in atherosclerosis. *J Biol Chem*. 2010;285:26545–26557.
310. Fenyó IM, Florea IC, Raicu M, Manea A. Tyrphostin ag490 reduces naphd oxidase activity and expression in the aorta of hypercholesterolemia apolipoprotein e-deficient mice. *Vasc Pharmacol*. 2011;54:100–106.
311. Guzik TJ, Sadowski J, Guzik B, Jopek A, Kapelak B, Przybylowski P, Wierzbicki K, Korbut R, Harrison DG, Channon KM. Coronary artery superoxide production and nox isoform expression in human coronary artery disease. *Arterioscler Thromb Vasc Biol*. 2006;26:333–339.
312. Azumi H, Inoue N, Takeshita S, Rikitake Y, Kawashima S, Hayashi Y, Itoh H, Yokoyama M. Expression of NADH/NADPH oxidase p22(phox) in human coronary arteries. *Circulation*. 1999;100:1494–1498.
313. Kalinina N, Agrotis A, Tararak E, Antropova Y, Kanellakis P, Ilyinskaya O, Quinn MT, Smirnov V, Bobik A. Cytochrome b558-dependent NAD(P)H oxidase-phox units in smooth muscle and macrophages of atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2002;22:2037–2043.
314. Kirk EA, Dinauer MC, Rosen H, Chait A, Heinecke JW, LeBoeuf RC. Impaired superoxide production due to a deficiency in phagocyte NADPH oxidase fails to inhibit atherosclerosis in mice. *Arterioscler Thromb Vasc Biol*. 2000;20:1529–1535.
315. Judkins CP, Diep H, Broughton BR, Mast AE, Hooker EU, Miller AA, Selemidis S, Dusting GJ, Sobey CG, Drummond GR. Direct evidence of a role for nox2 in superoxide production, reduced nitric oxide bioavailability, and early atherosclerotic plaque formation in apoe-/- mice. *Am J Physiol Heart Circ Physiol*. 2010;298:H24–H32.
316. Hsieh E, Segal BH, Pagano PJ, Rey FE, Paigen B, Deleonardis J, Hoyt RF, Holland SM, Finkel T. Vascular effects following homozygous disruption of p47(phox): an essential component of NADPH oxidase. *Circulation*. 2000;101:1234–1236.
317. Barry-Lane PA, Patterson C, van der Merwe M, Hu Z, Holland SM, Yeh ET, Runge MS. P47phox is required for atherosclerotic lesion progression in apoe-/- mice. *J Clin Invest*. 2001;108:1513–1522.
318. Chen Z, Keaney JF Jr, Schulz E, Levison B, Shan L, Sakuma M, Zhang X, Shi C, Hazen SL, Simon DI. Decreased neointimal formation in nox2-deficient mice reveals a direct role for NADPH oxidase in the response to arterial injury. *Proc Natl Acad Sci U S A*. 2004;101:13014–13019.
319. Fukui T, Ishizaka N, Rajagopalan S, Laursen JB, Capers Q IV, Taylor WR, Harrison DG, de Leon H, Wilcox JN, Griendling KK. P22phox mRNA expression and NADPH oxidase activity are increased in aortas from hypertensive rats. *Circ Res*. 1997;80:45–51.

320. Gauss KA, Nelson-Overton LK, Siemsen DW, Gao Y, DeLeo FR, Quinn MT. Role of nf-kappab in transcriptional regulation of the phagocyte NADPH oxidase by tumor necrosis factor-alpha. *J Leukoc Biol.* 2007;82:729–741.
321. Ding H, Hashem M, Triggle C. Increased oxidative stress in the streptozotocin-induced diabetic apoe-deficient mouse: changes in expression of NADPH oxidase subunits and enos. *Eur J Pharmacol.* 2007;561:121–128.
322. Li J, Wang JJ, Yu Q, Chen K, Mahadev K, Zhang SX. Inhibition of reactive oxygen species by lovastatin downregulates vascular endothelial growth factor expression and ameliorates blood-retinal barrier breakdown in db/db mice: role of NADPH oxidase 4. *Diabetes.* 2010;59:1528–1538.
323. Tian XY, Wong WT, Xu A, Chen ZY, Lu Y, Liu LM, Lee VW, Lau CW, Yao X, Huang Y. Rosuvastatin improves endothelial function in db/db mice: role of angiotensin II type 1 receptors and oxidative stress. *Br J Pharmacol.* 2011;164:598–606.
324. Guzik TJ, Chen W, Gongora MC, Guzik B, Lob HE, Mangalat D, Hoch N, Dikalov S, Rudzinski P, Kapelak B, Sadowski J, Harrison DG. Calcium-dependent nox5 nicotinamide adenine dinucleotide phosphate oxidase contributes to vascular oxidative stress in human coronary artery disease. *J Am Coll Cardiol.* 2008;52:1803–1809.
325. Yun J, Rocic P, Pung YF, Belmadani S, Carrao AC, Ohanyan V, Chilian WM. Redox-dependent mechanisms in coronary collateral growth: the “redox window” hypothesis. *Antioxidant Redox Signal.* 2009;11:1961–1974.
326. Arbiser JL, Petros J, Klafter R, Govindajaran B, McLaughlin ER, Brown LF, Cohen C, Moses M, Kilroy S, Arnold RS, Lambeth JD. Reactive oxygen generated by nox1 triggers the angiogenic switch. *Proc Natl Acad Sci U S A.* 2002;99:715–720.
327. Komatsu D, Kato M, Nakayama J, Miyagawa S, Kamata T. NADPH oxidase 1 plays a critical mediating role in oncogenic ras-induced vascular endothelial growth factor expression. *Oncogene.* 2008;27:4724–4732.
328. Tojo T, Ushio-Fukai M, Yamaoka-Tojo M, Ikeda S, Patrushev N, Alexander RW. Role of gp91phox (nox2)-containing NAD(P)H oxidase in angiogenesis in response to hindlimb ischemia. *Circulation.* 2005;111:2347–2355.
329. Urao N, Inomata H, Razvi M, Kim HW, Wary K, McKinney R, Fukai T, Ushio-Fukai M. Role of nox2-based NADPH oxidase in bone marrow and progenitor cell function involved in neovascularization induced by hindlimb ischemia. *Circ Res.* 2008;103:212–220.
330. Haddad P, Dussault S, Groleau J, Turgeon J, Maingrette F, Rivard A. Nox2-derived reactive oxygen species contribute to hypercholesterolemia-induced inhibition of neovascularization: effects on endothelial progenitor cells and mature endothelial cells. *Atherosclerosis.* 2011;217:340–349.
331. Ebrahimi TG, Heymes C, You D, Blanc-Brude O, Mees B, Waeckel L, Duriez M, Vilar J, Brandes RP, Levy BI, Shah AM, Silvestre JS. NADPH oxidase-derived overproduction of reactive oxygen species impairs postischemic neovascularization in mice with type 1 diabetes. *Am J Pathol.* 2006;169:719–728.
332. Zhang M, Brewer AC, Schroder K, Santos CX, Grieve DJ, Wang M, Anilkumar N, Yu B, Dong X, Walker SJ, Brandes RP, Shah AM. NADPH oxidase-4 mediates protection against chronic load-induced stress in mouse hearts by enhancing angiogenesis. *Proc Natl Acad Sci U S A.* 2010;107:18121–18126.
333. Bhandarkar SS, Jaconi M, Fried LE, et al. Fulvene-5 potently inhibits NADPH oxidase 4 and blocks the growth of endothelial tumors in mice. *J Clin Invest.* 2009;119:2359–2365.
334. Jackman KA, Miller AA, Drummond GR, Sobey CG. Importance of nox1 for angiotensin II-induced cerebrovascular superoxide production and cortical infarct volume following ischemic stroke. *Brain Res.* 2009;1286:215–220.
335. Kahles T, Kohlen A, Heumueller S, Rappert A, Bechmann I, Liebner S, Wittko IM, Neumann-Haefelin T, Steinmetz H, Schroeder K, Brandes RP. NADPH oxidase nox1 contributes to ischemic injury in experimental stroke in mice. *Neurobiol Dis.* 2010;40:185–192.
336. Walder CE, Green SP, Darbonne WC, Mathias J, Rae J, Dinauer MC, Curnutte JT, Thomas GR. Ischemic stroke injury is reduced in mice lacking a functional NADPH oxidase. *Stroke.* 1997;28:2252–2258.
337. Brait VH, Jackman KA, Walduck AK, Selemidis S, Diep H, Mast AE, Guida E, Broughton BR, Drummond GR, Sobey CG. Mechanisms contributing to cerebral infarct size after stroke: gender, reperfusion, t lymphocytes, and nox2-derived superoxide. *J Cereb Blood Flow Metab.* 2010;30:1306–1317.
338. Bell RM, Cave AC, Johar S, Hearse DJ, Shah AM, Shattock MJ. Pivotal role of NOX-2-containing NADPH oxidase in early ischemic preconditioning. *FASEB J.* 2005;19:2037–2039.
339. Nisbet RE, Graves AS, Kleinhenz DJ, Rupnow HL, Reed AL, Fan TH, Mitchell PO, Sutliff RL, Hart CM. The role of NADPH oxidase in chronic intermittent hypoxia-induced pulmonary hypertension in mice. *Am J Resp Cell Mol Biol.* 2009;40:601–609.
340. Vallet P, Charnay Y, Steger K, Ogier-Denis E, Kovari E, Herrmann F, Michel JP, Szanto I. Neuronal expression of the NADPH oxidase nox4, and its regulation in mouse experimental brain ischemia. *Neuroscience.* 2005;132:233–238.
341. McCann SK, Dusting GJ, Roulston CL. Early increase of nox4 NADPH oxidase and superoxide generation following endothelin-1-induced stroke in conscious rats. *J Neurosci Res.* 2008;86:2524–2534.
342. Kleinschnitz C, Grund H, Wingle K, et al. Post-stroke inhibition of induced NADPH oxidase type 4 prevents oxidative stress and neurodegeneration. *PLoS Biol.* 2010;8:e1000479.
343. Infanger DW, Cao X, Butler SD, Burmeister MA, Zhou Y, Stupinski JA, Sharma RV, Davisson RL. Silencing nox4 in the paraventricular nucleus improves myocardial infarction-induced cardiac dysfunction by attenuating sympathoexcitation and perinfarct apoptosis. *Circ Res.* 2010;106:1763–1774.
344. Bendall JK, Cave AC, Heymes C, Gall N, Shah AM. Pivotal role of a gp91(phox)-containing NADPH oxidase in angiotensin II-induced cardiac hypertrophy in mice. *Circulation.* 2002;105:293–296.
345. Satoh M, Ogita H, Takeshita K, Mukai Y, Kwiatkowski DJ, Liao JK. Requirement of racl1 in the development of cardiac hypertrophy. *Proc Natl Acad Sci U S A.* 2006;103:7432–7437.
346. Johar S, Cave AC, Narayanapanicker A, Grieve DJ, Shah AM. Aldosterone mediates angiotensin II-induced interstitial cardiac fibrosis via a nox2-containing NADPH oxidase. *FASEB J.* 2006;20:1546–1548.
347. Doerries C, Grote K, Hilfiker-Kleiner D, Luchtefeld M, Schaefer A, Holland SM, Sorrentino S, Manes C, Schieffer B, Drexler H, Landmesser U. Critical role of the NAD(P)H oxidase subunit p47phox for left ventricular remodeling/dysfunction and survival after myocardial infarction. *Circ Res.* 2007;100:894–903.
348. Hoffmeyer MR, Jones SP, Ross CR, Sharp B, Grisham MB, Laroux FS, Stalker TJ, Scalia R, Lefler DJ. Myocardial ischemia/reperfusion injury in NADPH oxidase-deficient mice. *Circ Res.* 2000;87:812–817.
349. Byrne JA, Grieve DJ, Bendall JK, Li JM, Gove C, Lambeth JD, Cave AC, Shah AM. Contrasting roles of NADPH oxidase isoforms in pressure-overload versus angiotensin II-induced cardiac hypertrophy. *Circ Res.* 2003;93:802–805.
350. Maytin M, Siwik DA, Ito M, Xiao L, Sawyer DB, Liao R, Colucci WS. Pressure overload-induced myocardial hypertrophy in mice does not require gp91phox. *Circulation.* 2004;109:1168–1171.
351. Grieve DJ, Byrne JA, Siva A, Layland J, Johar S, Cave AC, Shah AM. Involvement of the nicotinamide adenine dinucleotide phosphate oxidase isoform nox2 in cardiac contractile dysfunction occurring in response to pressure overload. *J Am Coll Cardiol.* 2006;47:817–826.
352. Kuroda J, Ago T, Matsushima S, Zhai P, Schneider MD, Sadoshima J. NADPH oxidase 4 (nox4) is a major source of oxidative stress in the failing heart. *Proc Natl Acad Sci U S A.* 2010;107:15565–15570.
353. Ushio-Fukai M, Tang Y, Fukai T, Dikalov SI, Ma Y, Fujimoto M, Quinn MT, Pagano PJ, Johnson C, Alexander RW. Novel role of gp91(phox)-containing NAD(P)H oxidase in vascular endothelial growth factor-induced signaling and angiogenesis. *Circ Res.* 2002;91:1160–1167.
354. Pendyala S, Usatyuk PV, Gorskova IA, Garcia JG, Natarajan V. Regulation of NADPH oxidase in vascular endothelium: the role of phospholipases, protein kinases, and cytoskeletal proteins. *Antioxidant Redox Signal.* 2009;11:841–860.
355. Doughan AK, Harrison DG, Dikalov SI. Molecular mechanisms of angiotensin II-mediated mitochondrial dysfunction: linking mitochondrial oxidative damage and vascular endothelial dysfunction. *Circ Res.* 2008;102:488–496.
356. Spallarossa P, Altieri P, Garibaldi S, Ghigliotti G, Barisione C, Manca V, Fabbri P, Ballestrero A, Brunelli C, Barsotti A. Matrix metalloproteinase-2 and -9 are induced differently by doxorubicin in h9c2 cells: the role of map kinases and NAD(P)H oxidase. *Cardiovasc Res.* 2006;69:736–745.
357. Sorecucu GP, Song H, Tressel SL, Hwang J, Dikalov S, Smith DA, Boyd NL, Platt MO, Lassègue B, Griendling KK, Jo H. Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress induces

- monocyte adhesion by stimulating reactive oxygen species production from a nox1-based NADPH oxidase. *Circ Res*. 2004;95:773–779.
358. Fan C, Kawai Y, Inaba S, Arakawa K, Katsuyama M, Kajinami K, Yasuda T, Yabe-Nishimura C, Konoshita T, Miyamori I. Synergy of aldosterone and high salt induces vascular smooth muscle hypertrophy through up-regulation of nox1. *J Steroid Biochem Mol Biol*. 2008;111:29–36.
 359. Manea A, Tanase LI, Raicu M, Simionescu M. Jak/stat signaling pathway regulates nox1 and nox4-based NADPH oxidase in human aortic smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 2010;30:105–112.
 360. Briones AM, Tabet F, Callera GE, Montezano AC, Yogi A, He Y, Quinn MT, Salaices M, Touyz RM. Differential regulation of nox1, nox2 and nox4 in vascular smooth muscle cells from wky and shr. *J Am Soc Hypertens*. 2011;5:137–153.
 361. Cevik MO, Katsuyama M, Kanda S, Kaneko T, Iwata K, Ibi M, Matsuno K, Kakehi T, Cui W, Sasaki M, Yabe-Nishimura C. The ap-1 site is essential for the promoter activity of nox1/NADPH oxidase, a vascular superoxide-producing enzyme: possible involvement of the erk1/2-junb pathway. *Biochem Biophys Res Commun*. 2008;374:351–355.
 362. Katsuyama M, Fan C, Arakawa N, Nishinaka T, Miyagishi M, Taira K, Yabe-Nishimura C. Essential role of atf-1 in induction of nox1, a catalytic subunit of NADPH oxidase: involvement of mitochondrial respiratory chain. *Biochem J*. 2005;386:255–261.
 363. Katsuyama M, Ozgur Cevik M, Arakawa N, Kakehi T, Nishinaka T, Iwata K, Ibi M, Matsuno K, Yabe-Nishimura C. Myocyte enhancer factor 2b is involved in the inducible expression of nox1/NADPH oxidase, a vascular superoxide-producing enzyme. *FEBS J*. 2007;274:5128–5136.
 364. Manea A, Tanase LI, Raicu M, Simionescu M. Transcriptional regulation of NADPH oxidase isoforms, nox1 and nox4, by nuclear factor-kappab in human aortic smooth muscle cells. *Biochem Biophys Res Commun*. 2010;396:901–907.
 365. Drummond GR, Selemidis S, Griendling KK, Sobey CG. Combating oxidative stress in vascular disease: NADPH oxidases as therapeutic targets. *Nature Rev Drug Discover*. 2011;10:453–471.
 366. Heymes C, Bendall JK, Ratajczak P, Cave AC, Samuel JL, Hasenfuss G, Shah AM. Increased myocardial NADPH oxidase activity in human heart failure. *J Am Coll Cardiol*. 2003;41:2164–2171.
 367. Park YM, Lim BH, Touyz RM, Park JB. Expression of NAD(P)H oxidase subunits and their contribution to cardiovascular damage in aldosterone/salt-induced hypertensive rat. *J Korean Med Sci*. 2008;23:1039–1045.
 368. Meischl C, Krijnen PA, Sipkens JA, Cillessen SA, Munoz IG, Okroj M, Ramska M, Muller A, Visser CA, Musters RJ, Simonides WS, Hack CE, Roos D, Niessen HW. Ischemia induces nuclear nox2 expression in cardiomyocytes and subsequently activates apoptosis. *Apoptosis*. 2006;11:913–921.
 369. Maejima Y, Kuroda J, Matsushima S, Ago T, Sadoshima J. Regulation of myocardial growth and death by NADPH oxidase. *J Mol Cell Cardiol*. 2011;50:408–416.
 370. Li JM, Shah AM. Intracellular localization and preassembly of the NADPH oxidase complex in cultured endothelial cells. *J Biol Chem*. 2002;277:19952–19960.
 371. Pollock JD, Williams DA, Gifford MA, Li LL, Du X, Fisherman J, Orkin SH, Doerschuk CM, Dinuer MC. Mouse model of x-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide production. *Nat Genet*. 1995;9:202–209.
 372. Muzaffar S, Shukla N, Angelini GD, Jeremy JY. NADPH oxidase 4 mediates upregulation of type 4 phosphodiesterases in human endothelial cells. *J Cell Physiol*. 2012;227:1941–1950.
 373. Zhang L, Sheppard OR, Shah AM, Brewer AC. Positive regulation of the NADPH oxidase nox4 promoter in vascular smooth muscle cells by e2f. *Free Radic Biol Med*. 2008;45:679–685.
 374. Diebold I, Petry A, Hess J, Gorch A. The NADPH oxidase subunit nox4 is a new target gene of the hypoxia-inducible factor-1. *Mol Biol Cell*. 2010;21:2087–2096.
 375. Pendyala S, Moitra J, Kalari S, Kleeberger SR, Zhao Y, Reddy SP, Garcia JG, Natarajan V. Nrf2 regulates hyperoxia-induced nox4 expression in human lung endothelium: identification of functional antioxidant response elements on the nox4 promoter. *Free Radic Biol Med*. 2011;50:1749–1759.
 376. Goettsch C, Goettsch W, Brux M, Haschke C, Brunssen C, Muller G, Bornstein SR, Duerrschmidt N, Wagner AH, Morawietz H. Arterial flow reduces oxidative stress via an antioxidant response element and oct-1 binding site within the NADPH oxidase 4 promoter in endothelial cells. *Basic Res Cardiol*. 2011;106:551–561.
 377. Goettsch C, Goettsch W, Muller G, Seebach J, Schnittler HJ, Morawietz H. Nox4 overexpression activates reactive oxygen species and p38 mapk in human endothelial cells. *Biochem Biophys Res Commun*. 2009;380:355–360.
 378. Carneseccchi S, Deffert C, Donati Y, Basset O, Hinz B, Preynat-Seauve O, Guichard C, Arbiser JL, Banfi B, Pache JC, Barazzzone-Argiroffo C, Krause KH. A key role for nox4 in epithelial cell death during development of lung fibrosis. *Antioxidant Redox Signal*. 2011;15:607–619.
 379. Manea A, Manea SA, Gafencu AV, Raicu M, Simionescu M. Ap-1-dependent transcriptional regulation of NADPH oxidase in human aortic smooth muscle cells: role of p22phox subunit. *Arterioscler Thromb Vasc Biol*. 2008;28:878–885.
 380. Djordjevic T, Pogrebniak A, BelAiba RS, Bonello S, Wotzlaw C, Acker H, Hess J, Gorch A. The expression of the NADPH oxidase subunit p22phox is regulated by a redox-sensitive pathway in endothelial cells. *Free Radic Biol Med*. 2005;38:616–630.
 381. Moriwaki K, Kiyomoto H, Hitomi H, Ihara G, Kaifu K, Matsubara K, Hara T, Kondo N, Ohmori K, Nishiyama A, Fukui T, Kohno M. Interferon-gamma enhances superoxide production in human mesangial cells via the jak-stat pathway. *Kidney Int*. 2006;70:788–793.
 382. Li L, Frei B. Prolonged exposure to lps increases iron, heme, and p22phox levels and NADPH oxidase activity in human aortic endothelial cells: Inhibition by desferrioxamine. *Arterioscler Thromb Vasc Biol*. 2009;29:732–738.
 383. Hitomi H, Fukui T, Moriwaki K, Matsubara K, Sun GP, Rahman M, Nishiyama A, Kiyomoto H, Kimura S, Ohmori K, Abe Y, Kohno M. Synergistic effect of mechanical stretch and angiotensin II on superoxide production via NADPH oxidase in vascular smooth muscle cells. *J Hypertens*. 2006;24:1089–1095.
 384. Manea A, Manea SA, Gafencu AV, Raicu M. Regulation of NADPH oxidase subunit p22(phox) by nf-kb in human aortic smooth muscle cells. *Arch Physiol Biochem*. 2007;113:163–172.
 385. Ni W, Zhan Y, He H, Maynard E, Balschi JA, Oettgen P. Ets-1 is a critical transcriptional regulator of reactive oxygen species and p47(phox) gene expression in response to angiotensin II. *Circ Res*. 2007;101:985–994.
 386. Honjo T, Otsui K, Shiraki R, Kawashima S, Sawamura T, Yokoyama M, Inoue N. Essential role of nox1 in generation of reactive oxygen species induced by oxidized low-density lipoprotein in human vascular endothelial cells. *Endothelium*. 2008;15:137–141.

Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Biochemistry, Physiology, and Pathophysiology of NADPH Oxidases in the Cardiovascular System

Bernard Lassègue, Alejandra San Martín and Kathy K. Griendling

Circ Res. 2012;110:1364-1390

doi: 10.1161/CIRCRESAHA.111.243972

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2012 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circres.ahajournals.org/content/110/10/1364>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>