

Impaired G1-Arrest, Autophagy, and Apoptosis in *Atg7*-Knockout Mice

Shun Kageyama, Masaaki Komatsu

Atg7 Modulates p53 Activity to Regulate Cell Cycle and Survival During Metabolic Stress

Lee et al
Science. 2012;336:225–228.

Exit from cell division cycle, induction of autophagy, and activation of apoptosis in response to metabolic stress occur simultaneously or sequentially. However, the molecular interrelation(s) between these intracellular events remains obscure. Recent work by Finkel et al¹ provides evidence that *Atg7*, an autophagy-related protein, regulates G1-arrest by interacting with p53 and that p53-mediated apoptosis is activated under autophagy-deficient conditions.

Macroautophagy (hereafter referred to as autophagy) is a self-eating system conserved among eukaryotes. In this process, cellular components, including organelles, are entrapped into a double-membrane structure called the autophagosome and then degraded by lysosomal hydrolases (Figure 1).² In addition to its role in supplying amino acids in response to nutrient starvation, autophagy is involved in quality control to maintain cell health. Thus, inactivation of autophagy results in the accumulation of cytoplasmic protein aggregates of misfolded proteins and damaged and degenerated organelles, which compromise cell function and often result in life-threatening diseases.²

The molecular underpinnings of autophagosome formation were uncovered mainly in yeast *Saccharomyces cerevisiae*. To date, genetic studies of *S. cerevisiae* have identified >30 autophagy-related (*ATG*) genes, 18 of which are core *ATG* genes that are essential for autophagosome formation.³ Importantly, core *ATG* genes are well conserved in mammals, where the functions of the encoded proteins markedly overlap with the corresponding proteins in yeast, although a few mammal-specific proteins have been identified. *Atg* proteins are categorized into several subclasses, but they function sequentially and cooperatively to regulate autophagosome formation (Figure 1).⁴ The ubiquitin-like modifiers *Atg12* and *LC3* (mammalian *Atg8* homolog) are activated by the E1-like

enzyme *Atg7* and transferred to 2 different E2-like enzymes: *Atg10* and *Atg3*, respectively. Whereas *Atg12* forms an isopeptide bond with *Atg5*, *LC3* forms an amide bond with phosphatidylethanolamine (PE) in a reaction dependent on the *Atg12-Atg5* conjugation. *Atg16L* forms a high-molecular-weight complex with *Atg12-Atg5*. The *Atg12-Atg5-Atg16L* complex functions as an E3-like enzyme, determining the site of *LC3* lipidation.⁴ Although the *Atg12-Atg5-Atg16L* complex is required for elongation of the isolation membrane, the phosphatidylethanolamine-bound *LC3* (*LC3-II*) is thought to be important for membrane biogenesis and closure of the isolation membrane (Figure 1).⁴

In a recent article published in *Science*, Lee et al¹ reported that in primary mouse embryonic fibroblasts, loss of *Atg7* abrogated G1-arrest in response to metabolic stress and contact inhibition. Surprisingly, depending on the nature of metabolic stress, *Atg7* interacted with p53 directly, and this complex then bound to the promoter region of *Cdkn1a*, a cyclin-dependent kinase inhibitor (Figure 2). These events resulted in modulation of *Cdkn1a* expression and inhibition of entry into the S-phase. Paradoxically, the same group noticed the upregulation of p53-dependent proapoptotic genes, such as *Puma*, *Bax*, and *Noxa*, in primary *Atg7*^{-/-} mouse embryonic fibroblasts. Autophagy-deficient mouse embryonic fibroblasts showed increased production of reactive oxygen species and DNA damage, which activated the DNA damage sensor, ataxia teleangiectasia mutated kinase. Subsequently, the mediator molecule *Chk2* phosphorylated Ser20 of p53, followed by induction of a series of proapoptotic genes (Figure 2). As expected, changes in the expression of these genes were completely blocked by the additional loss of *Chk2*. The phosphorylation of p53 and enhanced cell death observed in *Atg7*^{-/-} mouse embryonic fibroblasts were restored by simultaneous depletion of *Chk2*. Furthermore, the neonatal lethality of *Atg7*-knockout mice was rescued, at least in part, in *Atg7 Chk2*-double knockout mice.

Finkel et al¹ have suggested that the interaction of *Atg7* with p53 in response to metabolic stress and contact inhibition is a rate-limiting step for the induction of *Cdkn1a*, followed by G1-arrest (Figure 2). What are the factors that enhance the *Atg7*-p53 interaction? Evidence suggests that acetylation plays an important role in autophagy.^{5,6} Previous studies indicated that Sirt1 deacetylates several *Atg* proteins, including *Atg7*, which is indispensable for starvation-induced autophagy,⁷ and conversely that p300 acetyltransferase directly interacts with and acetylates *Atg7* to inhibit autophagy.⁸ Beyond autophagic regulation, deacetylation of *Atg7* might promote the interaction with p53 to induce *Cdkn1a* gene expression. Because the structures of both *Atg7* and tetramerization domain of p53 have been analyzed,^{9,10} determination of the structure of the

The opinions in this Commentary are not necessarily those of the editors or of the American Heart Association.

Commentaries serve as a forum in which experts highlight and discuss articles (published here and elsewhere) that the editors of *Circulation Research* feel are of particular significance to cardiovascular medicine.

Commentaries are edited by Aruni Bhatnagar & Ali J. Marian.

From the Protein Metabolism Project, Tokyo Metropolitan Institute of Medical Science, Setagaya-ku, Tokyo, Japan.

Correspondence to Masaaki Komatsu, Protein Metabolism Project, Tokyo Metropolitan Institute of Medical Science, Setagaya-ku, Tokyo 156-8506, Japan. E-mail komatsu-ms@igakuken.or.jp

(*Circ Res*. 2012;111:962-964.)

© 2012 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>
DOI: 10.1161/CIRCRESAHA.112.280768




Figure 1. Autophagy. The initial steps of autophagy include the formation and subsequent elongation of the isolation membrane. The isolation membrane then envelops various cytoplasmic constituents, such as organelles, until its edges fuse with each other to form a double-membrane structure called the autophagosome. Finally, the outer membrane of the autophagosome fuses with the lysosome and endosome. The sequestered cytoplasmic components, together with the inner membrane of the autophagosome, are completely degraded by lysosomal hydrolases (Mizushima et al⁴ for details on the molecular players). The ubiquitin-like modifier, LC3, covalently conjugates with phosphatidylethanolamine (PE) through an enzymatic cascade consisting of Atg7 (E1-like enzyme), Atg3 (E2-like enzyme), and Atg12-Atg5-Atg16L complex (function as an E3-like enzyme). The PE-conjugated LC3, named LC3-II, is localized to the inner and outer membranes of the isolation membrane and is essential for membrane biogenesis and closure of the isolation membrane. The activated ULK1 complex regulates class III PI3K complex and the multimembrane spanning protein Atg9L at pre-autophagosomal structure/phagophore-assembly site (PAS) close to the endoplasmic reticulum (ER), which is involved in the formation of the omegasome. PI3P-binding WIPIs, Atg12-Atg5-Atg16L1 complex, and LC3-PE conjugate play important roles in the elongation and closure of the isolation membrane/phagophore. ER indicates endoplasmic reticulum; PI3P, phosphatidylinositol 3-phosphate.

Atg7-TET domain complex is feasible. Structural analysis should enhance our understanding of the role of *Atg7* in cell cycle regulation.

What is the physiological importance of the p53-*Atg7* pathway in the regulation of cell cycle arrest? Exit from cell division cycle (ie, upregulation of *Cdkn1a*) seems to be dependent on *Atg7* but not other *Atg* proteins, such as *Atg5* and *Beclin 1*, indicating an additive function of *Atg7* beyond autophagy. Hence, loss of *Atg7* in mice should be associated with worse phenotypes than other *Atg*-knockout mice. Nevertheless, to date, we have not observed any profound phenotypes of *Atg7*-knockout mice compared with those of *Atg5*- or *Atg3*-knockout mice.^{11–13} These mutant mice are born in accordance with Mendelian inheritance ratios and survive but have lower amino acid levels in their sera and tissues and die within the first day of life. Further *in vivo* add-back experiment of mutant *Atg7*, which is defective in interaction with p53, but capable of induction of autophagy, will be critical for understanding the physiological significance of the p53-*Atg7* pathway.

Overcoming neonatal death in autophagy-deficient mice by additional loss of *Chk2* is surprising because the autophagy-deficient mice have multiple systemic defects, including neurological abnormalities and impaired trophic dynamics.^{11,14}

Finkel et al¹ have shown the induction of proapoptotic pathways only in livers of *Atg7*^{-/-} neonate mice. The livers of neonate mice contain a large number of hematopoietic stem cells and related cells, such as erythroid cells. Recent studies have shed light on the importance of autophagy in mitochondrial homeostasis (mitophagy)¹⁵ and indicated that defective mitophagy in hematopoietic cells is directly linked to rapid cell death. In fact, impaired autophagy in hematopoietic stem cells, T cells, and B cells causes acute cell death because of the accumulation of damaged mitochondria, followed by production of reactive oxygen species.^{16,17} Therefore, suppression of cell death in these cells by additional loss of *Chk2* is plausible. What about other tissues? The authors asserted that there was neurodegeneration in *Atg7 Chk2*-double knockout mice. However, partial rescue of neurological defect in autophagy-deficient mice is expected because loss of *Chk2* restores the suckling disorder in *Atg7*^{-/-} mice (the double-mutant mouse cannot survive >1 day unless it overcomes this defect).

Imbalance between cell growth and death contributes to tumorigenesis. Mice heterozygous for *Beclin 1* mutation, systemically mosaic for *Atg5* deficiency, or lacking *Atg7* specifically in liver tissue develop liver tumors, probably because of dysregulation of signal transduction pathways, as




Figure 2. Schematic model of cell cycle and cell death in *Atg7*-deficient mouse embryonic fibroblasts (MEFs) under metabolic stress conditions. In response to metabolic stress, the p53-Atg7 complex binds to the promoter region of *Cdkn1a* to induce its expression, leading to G1-arrest. Lack of *Atg7* abrogates the induction of *Cdkn1a* and exit from cell division cycle. Meanwhile, accumulation of DNA damage through impaired mitochondria homeostasis in *Atg7*-knockout mice activates DNA damage sensor, ATM kinase. Subsequently, Chk2 phosphorylates Ser20 of p53, followed by induction of a series of proapoptotic genes. Imbalance between cell division and cell death could lead to hematopoietic cell death, neurodegeneration, and tumorigenesis. ROS indicates reactive oxygen species.

well as impaired organelle quality control.^{18–21} The growth of liver adenoma in mice with liver-specific knockouts of *Atg7* is markedly suppressed when *p62* is also knocked out, because marked accumulation of *p62*, due to loss of autophagy, leads to dysregulation of nuclear factor- κ B signaling, activation of apoptosis, and responses to environmental stress.^{22,23} However, loss of *p62* does not suppress hepatocarcinogenesis itself. The autophagy-related p53 pathways might be engaged in tumorigenesis (Figure 2). Further studies are needed to determine the state of tumorigenesis in *Atg7 Chk2*-double deficient mice.

The report by Finkel et al¹ helps our understanding of the molecular mechanism by which impaired autophagy and Atg-related proteins are involved in apoptosis and cell cycle arrest. However, pathophysiological significance of the regulation remains unclear as described in this commentary. With the recent development of sophisticated research tools, such as Cre-mediated conditional knockout techniques, it has become clear that loss of autophagy is associated with various life-threatening diseases, such as neurodegeneration, cardiomyopathy, liver injury, glomerulosclerosis, myopathy, and cancer.² Further studies are needed to clarify the pathological role of proapoptotic genes in autophagy-deficient tissues in these diseases (Figure 2).

Sources of Funding

M. Komatsu is supported by the Funding Program for Next Generation World-Leading Researchers (Japan).

Disclosures

None.

References

- Lee IH, Kawai Y, Fergusson MM, Rovira II, Bishop AJ, Motoyama N, Cao L, Finkel T. Atg7 modulates p53 activity to regulate cell cycle and survival during metabolic stress. *Science*. 2012;336:225–228.
- Mizushima N, Komatsu M. Autophagy: renovation of cells and tissues. *Cell*. 2011;147:728–741.
- Nakatogawa H, Suzuki K, Kamada Y, Ohsumi Y. Dynamics and diversity in autophagy mechanisms: lessons from yeast. *Nat Rev Mol Cell Biol*. 2009;10:458–467.
- Mizushima N, Yoshimori T, Ohsumi Y. The role of Atg proteins in autophagosome formation. *Annu Rev Cell Dev Biol*. 2011;27:107–132.
- Yi C, Ma M, Ran L, et al. Function and molecular mechanism of acetylation in autophagy regulation. *Science*. 2012;336:474–477.
- Lin SY, Li TY, Liu Q, et al. GSK3-TIP60-ULK1 signaling pathway links growth factor deprivation to autophagy. *Science*. 2012;336:477–481.
- Lee IH, Cao L, Mostoslavsky R, Lombard DB, Liu J, Bruns NE, Tsokos M, Alt FW, Finkel T. A role for the NAD-dependent deacetylase Sirt1 in the regulation of autophagy. *Proc Natl Acad Sci USA*. 2008;105:3374–3379.
- Lee IH, Finkel T. Regulation of autophagy by the p300 acetyltransferase. *J Biol Chem*. 2009;284:6322–6328.
- Cho Y, Gorina S, Jeffrey PD, Pavletich NP. Crystal structure of a p53 tumor suppressor-DNA complex: understanding tumorigenic mutations. *Science*. 1994;265:346–355.
- Noda NN, Satou K, Fujioka Y, Kumeta H, Ogura K, Nakatogawa H, Ohsumi Y, Inagaki F. Structural basis of Atg8 activation by a homodimeric E1, Atg7. *Mol Cell*. 2011;44:462–475.
- Kuma A, Hatano M, Matsui M, Yamamoto A, Nakaya H, Yoshimori T, Ohsumi Y, Tokuhisa T, Mizushima N. The role of autophagy during the early neonatal starvation period. *Nature*. 2004;432:1032–1036.
- Komatsu M, Waguri S, Ueno T, Iwata J, Murata S, Tanida I, Ezaki J, Mizushima N, Ohsumi Y, Uchiyama Y, Kominami E, Tanaka K, Chiba T. Impairment of starvation-induced and constitutive autophagy in *Atg7*-deficient mice. *J Cell Biol*. 2005;169:425–434.
- Sou YS, Waguri S, Iwata J, Ueno T, Fujimura T, Hara T, Sawada N, Yamada A, Mizushima N, Uchiyama Y, Kominami E, Tanaka K, Komatsu M. The Atg8 conjugation system is indispensable for proper development of autophagic isolation membranes in mice. *Mol Biol Cell*. 2008;19:4762–4775.
- Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R, Yokoyama M, Mishima K, Saito I, Okano H, Mizushima N. Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice. *Nature*. 2006;441:885–889.
- Youle RJ, Narendra DP. Mechanisms of mitophagy. *Nat Rev Mol Cell Biol*. 2011;12:9–14.
- Pua HH, Dzhagalov I, Chuck M, Mizushima N, He YW. A critical role for the autophagy gene *Atg5* in T cell survival and proliferation. *J Exp Med*. 2007;204:25–31.
- Mortensen M, Soilleux EJ, Djordjevic G, Tripp R, Lutteropp M, Sadighi-Akha E, Stranks AJ, Glanville J, Knight S, Jacobsen SE, Kranc KR, Simon AK. The autophagy protein *Atg7* is essential for hematopoietic stem cell maintenance. *J Exp Med*. 2011;208:455–467.
- Yue Z, Jin S, Yang C, Levine AJ, Heintz N. Beclin 1, an autophagy gene essential for early embryonic development, is a haploinsufficient tumor suppressor. *Proc Natl Acad Sci USA*. 2003;100:15077–15082.
- Qu X, Yu J, Bhagat G, Furuya N, Hibshoosh H, Troxel A, Rosen J, Eskelinen EL, Mizushima N, Ohsumi Y, Cattoretti G, Levine B. Promotion of tumorigenesis by heterozygous disruption of the beclin 1 autophagy gene. *J Clin Invest*. 2003;112:1809–1820.
- Inami Y, Waguri S, Sakamoto A, Kouno T, Nakada K, Hino O, Watanabe S, Ando J, Iwadate M, Yamamoto M, Lee MS, Tanaka K, Komatsu M. Persistent activation of Nrf2 through p62 in hepatocellular carcinoma cells. *J Cell Biol*. 2011;193:275–284.
- Takamura A, Komatsu M, Hara T, Sakamoto A, Kishi C, Waguri S, Eishi Y, Hino O, Tanaka K, Mizushima N. Autophagy-deficient mice develop multiple liver tumors. *Genes Dev*. 2011;25:795–800.
- Mathew R, Karp CM, Beaudoin B, Vuong N, Chen H, Chen HY, Bray K, Reddy A, Bhanot G, Gelinas C, Dipaola RS, Karantzis-Wadsworth V, White E. Autophagy suppresses tumorigenesis through elimination of p62. *Cell*. 2009;137:1062–1075.
- Komatsu M, Kurokawa H, Waguri S, et al. The selective autophagy substrate p62 activates the stress responsive transcription factor Nrf2 through inactivation of Keap1. *Nat Cell Biol*. 2010;12:213–223.

Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Impaired G1-Arrest, Autophagy, and Apoptosis in *Atg7*-Knockout Mice Shun Kageyama and Masaaki Komatsu

Circ Res. 2012;111:962-964

doi: 10.1161/CIRCRESAHA.112.280768

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2012 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circres.ahajournals.org/content/111/8/962>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>