

This Review is in a thematic series on **Autophagy in Health and Disease**, which includes the following articles:

Molecular Mechanisms of Autophagy in the Cardiovascular System [*Circ Res* 2015;116:456–467]

Autophagy in Vascular Disease [*Circ Res* 2015;116:468–479]

The Role of Autophagy in Vascular Biology [*Circ Res* 2015;116:480–488]

Therapeutic Targeting of Autophagy: Potential and Concerns in Treating Cardiovascular Disease [*Circ Res* 2015;116:489–503]

Untangling Autophagy Measurements: All Fluxed Up [*Circ Res* 2015;116:504–514]

Molecular Mechanisms of Mitochondrial Autophagy/Mitophagy in the Heart

Daniel Klionsky, Guest Editor

Molecular Mechanisms of Mitochondrial Autophagy/Mitophagy in the Heart

Toshiro Saito, Junichi Sadoshima

Abstract: Mitochondrial quality is a crucial determinant of cell viability, and mitochondrial autophagy plays a central role in this control mechanism. Based on studies in yeast, numerous investigations of this process have been conducted, and the framework of mammalian mitochondrial autophagy is progressively appearing. However, many enigmas about the molecular mechanisms involved remain unsolved. Furthermore, the pathological significance of mitochondrial autophagy in the heart remains largely unclear. In this review, we discuss the current understanding of mitochondrial autophagy in mammals with reference to that in yeast. Regarding the process in yeast, some points of uncertainty have arisen. We also summarize recent advances in the research of autophagy and mitochondrial autophagy in the heart. This article is a part of a review series on *Autophagy in Health and Disease*. (*Circ Res*. 2015;116:1477-1490. DOI: 10.1161/CIRCRESAHA.116.303790.)

Key Words: autophagy ■ mitophagy ■ Parkin ■ Pink1

Mitochondria are evolutionarily preserved organelles found in almost all eukaryotic cells. The primary role of mitochondria in mammalian cells is adenosine triphosphate synthesis, which is achieved primarily through oxidative respiratory phosphorylation. Additionally, mitochondrial proteins also regulate innate immunity, apoptosis, necrosis, and autophagy.^{1–3} Adenosine triphosphate production through the respiratory chain is accompanied by production of reactive oxygen species (ROS) as a byproduct through leakage of electrons from the electron transport chain. Low levels of ROS are directly removed by antioxidants within mitochondria or actively stimulate counter-regulatory signaling/transcription pathways to maintain the proper redox balance. However, excessive accumulation of ROS during stress damages mitochondrial components, including mitochondrial DNA (mtDNA), protein, and lipids, which exacerbates ROS production and mitochondrial dysfunction in a feedforward manner. In

response to ROS, mitochondria initiate and orchestrate various signaling mechanisms, thereby controlling the growth and death of various cell types.⁴ Damaged mtDNA that escapes from lysosomal degradation causes Toll-like receptor 9-mediated inflammation and myocarditis as a result of its inflammatory unmethylated CpG motifs.⁵ Mitochondria-derived excessive ROS eventually promote aging, carcinogenesis, neurodegenerative disorders, and cardiovascular diseases.^{6,7} Because damaged mitochondria induce catastrophic consequences, the quality of mitochondria is essential and regulated by multiple mechanisms, including fission and fusion of mitochondria, degradation of mitochondrial proteins mediated by chaperones (Hsp10, Hsp60, and others), proteinases (Lon, AAA proteases), and proteasomes, and mitochondrial biogenesis.⁸ The constant production of ROS necessitates turnover and replacement of mitochondria every 10 to 25 days, even in the quiescent cell or organ.⁹ Increasing lines of evidence

Original Received December 18, 2014; revision received February 24, 2015; accepted February 26, 2015. In January 2015, the average time from submission to first decision for all original research papers submitted to *Circulation Research* was 14.7 days.

From the Department of Cell Biology and Molecular Medicine, New Jersey Medical School, Rutgers University, Newark.

Correspondence to Junichi Sadoshima, MD, PhD, Department of Cell Biology and Molecular Medicine, Rutgers–New Jersey Medical School, 185 S Orange Ave, MSB G609, Newark, NJ 07103. E-mail sadoshju@njms.rutgers.edu

© 2015 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.116.303790

Nonstandard Abbreviations and Acronyms

Bcl-2	b-cell leukemia/lymphoma 2 protein
Bnip3	BCL2/adenovirus E1B 19kDa interacting protein 3
CK2	casein kinase 2
Drp1	dynamitin 1-like
FUNDC1	FUN14 domain containing 1
GFP	green fluorescence protein
IMM	inner mitochondrial membrane
IPC	ischemic preconditioning
I/R	ischemia/reperfusion
LC3	microtubule associated protein 1 light chain 3
LIR	LC3-interacting region
Mfn	mitofusin
MI	myocardial infarction
mtDNA	mitochondrial deoxyribonucleic acid
Nix/Bnip3L	NIP3-like protein X
OMM	outer mitochondrial membrane
Opa1	optic atrophy 1
Parkin	Parkinson juvenile disease protein 2
Pink1	PTEN-inducible putative kinase 1
p53	tumor protein p53
p62	sequestosome 1
RFP	red fluorescent protein
ROS	reactive oxygen species
Ser	serine

suggest that autophagy plays a major role in mediating mitochondrial quality control.^{10,11}

Autophagy is a mechanism of cellular degradation through lysosomes. Autophagy is particularly important for cell survival during energy stress because amino acids and fatty acids recovered through degradation of cellular constituents by autophagy can be recycled to generate adenosine triphosphate. Autophagy is classified into 3 categories: macroautophagy, microautophagy, and chaperone-mediated autophagy.^{12–14} Macroautophagy is characterized by the formation of an autophagosome, a double-membrane structure which engulfs organelles or long-lived proteins and fuses with lysosomes. Microautophagy is defined by the direct incorporation of cytosolic content into a lysosome/vacuole by the invagination of its surface. Chaperone-mediated autophagy involves the direct translocation of proteins into lysosomes in a 70 kDa heat-shock cognate protein-mediated manner. The machinery of macroautophagy is fairly well understood, whereas that of microautophagy has not yet been fully elucidated in mammals. The process of macroautophagy proceeds in multiple sequential steps: initiation and nucleation of isolation membranes, elongation of phagophores and engulfment of organelles or long-lived proteins, docking and fusion of phagophores (autophagosome formation), fusion of autophagosomes with lysosomes, and lysosomal degradation of cargos. Although autophagy has been regarded as a nonselective process, accumulating lines of evidence indicate the existence of selective autophagy, through which damaged organelles, such as mitochondria, endoplasmic reticulum, ribosomes, and

peroxisomes, are selectively degraded. This selective autophagy is collectively called organelle-specific autophagy.^{15,16} Recent investigations have unveiled defined molecular mechanisms and the functions of organelle-specific autophagy. Among them, mitochondria-specific autophagy, termed mitophagy, has been most intensively investigated. Here, we discuss the current understanding of the molecular mechanisms mediating mitophagy and the role of mitophagy in mediating physiological and pathological functions in the heart.

What Is Organelle-Specific Autophagy?

The existence of a selective autophagic process for organelles was first proposed for the endoplasmic reticulum in 1973.^{17,18} Since then, molecular mechanisms mediating the organelle-specific autophagy in the endoplasmic reticulum (endoplasmic reticulum-phagy/reticulophagy), peroxisomes (pexophagy), ribosomes (ribophagy), and mitochondria (mitophagy) have been investigated in yeast.^{19–29} Both macroautophagy and microautophagy play a role in pexophagy and are termed macropexophagy and micropexophagy, respectively.^{19–21,28,30} Because microautophagy uses autophagy-related (Atg) proteins in yeast, it is hard to distinguish the particular contributions of microautophagy and macroautophagy to overall organelle-specific autophagy.³⁰ Although some studies suggest a contribution of microautophagy to mitophagy, its contribution is largely unknown.^{31,32} Therefore, we hereafter use the term mitophagy to reference macro-mitophagy.

Yeast peroxisomes were ideal organelles for the initial studies of selective autophagy.²¹ Macropexophagy eliminates redundant or inactive peroxisomes without acceleration of autophagic degradation of other organelles.²¹ Although peroxisomes can be degraded nonspecifically, the specific process also takes place under certain conditions, and the term pexophagy was generally accepted by researchers.^{18,19} Some labeling proteins localized in the peroxisome were later identified, such as peroxisomal membrane anchor protein PEX14, Atg30, and Atg37, which are essential for macropexophagy.^{20,28,33} The phosphorylation of Atg30 is essential for its interaction with Atg8 and Atg11, which is followed by macropexophagy.^{28,33,34}

The term mitophagy was first introduced by Lemasters in 2005 to describe the selective autophagy of mitochondria and emphasize the nonrandom nature of the process in yeast after the identification of SUN family protein UTH1.⁹ This term had been widely accepted without enough evidence for its selectivity at the beginning.¹⁸ The molecular basis of mitophagy has since been first investigated in yeast. Most importantly, mitochondrial-targeted proteins, such as SUN family protein UTH1, Aup1p, and Atg32, which mark damaged mitochondria and subject them to autophagic removal, have been identified in yeast.^{24,25,35,36} Atg32 accumulates in the outer membrane of mitochondria (OMM) in response to starvation or in the post-log phase.^{24,25} Atg32 interacts with Atg8 (Figure 1),^{24,25} After phosphorylation of Atg32 at serine (Ser) 114 and 119 by casein kinase 2 (CK2), the N-terminus of Atg32 binds to the C-terminus of Atg11 (Figure 1).²⁶ Subsequently, Atg11, which connects to dynamitin-related GTPase DNM1, a mitochondrial fission protein, recruits the fission complex to mitochondria, a step that is essential for mitophagy (Figure 1).³⁷

Although knowledge obtained from yeast has served as a useful resource for elucidating the mechanism of mitophagy in mammalian cells, the metabolic processes during the induction of mitophagy are distinct between yeast and mammalian cells. *Saccharomyces cerevisiae*, a budding yeast, prefers using the glycolytic pathway even when oxygen is abundant.³⁸ When glucose is exhausted, a diauxic shift takes place and yeast start to use mitochondrial oxidative respiratory phosphorylation.³⁸ On the other hand, mammalian cells primarily use mitochondrial oxidative phosphorylation in the presence of oxygen. Thus, induction of mitophagy in yeast and mammalian cells may be distinct in terms of functional significance and mechanism of activation.

In yeast, vacuoles containing mitochondria alone but no other organelles can be observed frequently through electron microscopic analyses. Yeast strains lacking mitophagy still retain bulk autophagy,²⁴ suggesting that mitophagy is clearly separable from general autophagy in yeast. In contrast, the boundary between mitophagy and bulk autophagy of mitochondria seems more obscure in mammalian cells. Thus, when studying mitophagy in mammalian cells, it is crucial to carefully observe whether the autophagosomes contain mitochondria alone or other cargos as well. Because many publications in the field of cardiovascular medicine do not distinguish mitophagy from general autophagy degrading mitochondria in a nonspecific manner, we here use the term mitophagy only when a given work most likely investigates mitochondria-specific autophagy. When the distinction is not clear, we use the term mitochondrial autophagy.

How Can We Detect Mitophagy in Mammalian Cells?

The simplest and most reliable demonstration of mitochondrial autophagy is through observation of mitochondria engulfed in autophagosomes, using an electron microscope. This approach is laborious but provides critical information, such as the content of the autophagosome and the structure of autophagosomes/autolysosomes. Concomitant use of immunogold antibodies can help identify the cargo content and the origin of the autophagosomes. The use of electron tomography, in which tilt series of data for each section are reconstituted into 3-dimensional images (3D), would be useful to prove the sole presence of mitochondria in autophagosomes, which is prerequisite for documenting the presence of mitophagy separate from general autophagy.^{39,40}

Practically, colocalization of microtubule-associated protein 1 light chain 3 (LC3) puncta and mitochondrial proteins is most commonly used to demonstrate the presence of mitochondrial autophagy. LC3 puncta are most conveniently observed with exogenous LC3 fused with fluorescent proteins, such as green fluorescence protein (GFP), mRFP, and mCherry. Time-lapsed imaging in living cells expressing these markers allows continuous observation of the incorporation of mitochondria into autophagosomes with the extension of the isolation membrane.⁴¹ As the fluorescence of GFP disappears under acidic conditions, GFP-LC3 puncta do not label autolysosomes. On the other hand, the fluorescence of mRFP and mCherry is resistant to the acidic environment of

lysosomes. Using the distinct properties of GFP and mRFP, mRFP–GFP tandem-fluorescence-tagged LC3 makes it possible to distinguish autophagosomes and autolysosomes. However, this evaluation may overlook the previously reported mitochondrial autophagy mediated by an LC3-independent mechanism.^{42–44}

Keima is a coral-derived acid-stable fluorescent protein that emits different-colored signals at acidic and neutral pHs.⁴⁵ Mitochondria incorporated into lysosomes are visualized as acidic-colored dots representing fluorescence emitted by mitochondria-targeted Keima.⁴⁵ Although this indicator alone cannot prove that translocation of mitochondrial proteins to lysosomes occurs through mitophagy, if the translocation is inhibited by interventions to suppress autophagy or mitophagy, it is possible to show that the lysosomal translocation of mitochondria-targeted Keima is caused by either autophagy or mitophagy.

Although microscopic morphological studies provide solid qualitative data, it is important to also show degradation of mitochondria using biochemical methods. The quantitative analysis of mitochondrial DNA or protein seems indispensable. It should be noted that, in addition to autophagy, the ubiquitin proteasome system also mediates degradation of mitochondria.⁴⁶ In particular, proteins associated with the OMM undergo degradation through the ubiquitin proteasome system under certain conditions.⁴⁶ In this sense, the molecules in the matrix or inner mitochondrial membrane (IMM), including mtDNA and proteins involved in the electron transport chain, may be helpful for the assessment of mitochondrial autophagy. Mito Timer, a fluorescent protein targeted to mitochondria, is useful for monitoring the turnover of mitochondrial proteins. Mito Timer transitions from green to red fluorescence over a span of 48 h. This property allows one to evaluate the turnover of mitochondria facilitated by mitophagy.⁴⁷ For example, Mito Timer in autophagy-deficient cells with low mitochondrial turnover tends to show more red than green.

The impairment of mitophagy or mitochondrial autophagy is often accompanied by accumulation of dysfunctional mitochondria. In theory, the volume of mitochondria may be quantitated by electron microscopic tomography.⁴⁸ However, estimating the total volume of mitochondria in cells using this method is extremely laborious and may not be practical. Accumulation of damaged mitochondria is often accompanied by increases in mtDNA content. This can be evaluated with short-range and long-range quantitative polymerase chain reaction, which allows for calculation of the amount of damaged mtDNA by subtracting intact mtDNA from total mtDNA.^{49,50} It should be noted that the impairment of mitochondrial dynamics caused by genetic deletion of mitofusin 1 (Mfn1) and Mfn2 induces severe depletion of mtDNA through unknown mechanisms, leading to mtDNA instability in skeletal muscle.⁵¹ Thus, suppression of mitophagy may not necessarily be accompanied by accumulation of mtDNA.⁵¹ Furthermore, mtDNA has been reported as a poor biomarker of mitochondrial content because mitochondrial DNA copy number can vary considerably from animal to animal.⁵² Thus, it is recommended that it be used in combination with other




Figure 1. The mechanism of mitophagy in yeast. Casein kinase 2 (CK2) directly phosphorylates autophagy-related protein (Atg)32 at serine (Ser) 114 and Ser 119 when cells are cultured in lactate medium and then shifted to nitrogen starvation medium supplemented with glucose. This phosphorylation, especially phosphorylation at Ser 114, is critically important for the Atg32–Atg11 interaction, which is required for the delivery of mitochondria to the phagosome assembly site (PAS). Atg11 recruits the mitochondrial fission complex, which consists of dynamin-related GTPase DNM1 (Dnm1), fission 1 (Fis1), Mdv1, and Caf4. Subsequently, mitochondrial fission and mitophagy take place. Mitochondrial division is important for mitophagy in yeast.

biomarkers. For example, accumulation of damaged mitochondria can also be demonstrated by increases in the mitochondrial mass, which can be evaluated by isolating mitochondria and determining their content relative to the starting tissue or cell preparation, or by increases in mitochondrial content, which is estimated by determining the relative ratios of mitochondrial biomarkers, including citrate synthase activity, cardiolipin content, and complex I-IV proteins, to cytosolic proteins.⁵² Mitochondrial swelling assays are used to show increased sensitivity of the mitochondrial permeability transition pore to Ca^{2+} . The mitochondrial membrane potential is often depolarized, and this can be monitored using JC-1, TMRE, or TMRM. Mitochondrial respiration can be evaluated with measurement of the time-dependent oxygen consumption using Clarke electrodes, and the activity of the electron transport chain can be monitored with biochemical analyses of freshly prepared mitochondrial fractions.

At present, there is no single experimental method sufficient for the assessment of mitochondrial autophagy because none of these assays alone can prove the presence of cargo-specific autophagy, such as mitophagy. Caution should be exercised when interpreting the results obtained from these assays. If more information regarding how mitophagy is induced becomes available in the future, the mechanism can be traced as an indicator of mitophagy, and loss-of-function approaches can be taken to show the specific involvement of mitophagy. Currently, Parkinson juvenile disease protein 2 (Parkin) translocation to mitochondria and Parkin-dependent ubiquitination of mitochondrial proteins (See below) are often used to show activation of mitophagy. However, whether this mechanism represents the major mechanism of mitochondrial autophagy in the presence of a given stress in the heart requires further investigation.




Figure 2. Microtubule-associated protein 1 light chain 3 (LC3)-interacting molecules act as receptors for autophagosomes on mitochondria. A, B, Nix and Bnip3, BH3-only proteins, interact with LC3 through the LC3-interacting region (LIR) and regulate mitochondrial autophagy. Oxidative stress induces homodimerization and activation of Bnip3. Phosphorylation of the LIR in Bnip3 promotes the association between Bnip3 and LC3. C, Under hypoxia, dephosphorylation of FUNDC1 at Ser 13 or tyrosine (Tyr) 18 stabilizes the interaction between FUNDC1 and LC3. D, Cardiolipin, a phospholipid of the IMM, translocates to the mitochondrial surface in response to mitochondrial damage. The externalized cardiolipin binds to LC3.

Molecular Mechanisms Mediating Mitophagy

Most of the studies conducted thus far have focused on the molecular mechanism by which damaged mitochondria are recognized and engulfed by autophagosomes. In the case of mitophagy in yeast, Atg32 labels damaged mitochondria and recruits phagophores by acting as a receptor for Atg8 through its Atg8 family interacting motif. In the case of mammalian




Figure 3. The mechanism of mitochondrial autophagy mediated by PTEN-inducible putative kinase 1 (Pink1)-Parkinson juvenile disease protein 2 (Parkin) in mammalian cells. Mitochondrial proteases and peptidases continuously degrade Pink1 in intact mitochondria. However, Pink1 is not imported to the inner membrane and is not cleaved in depolarized mitochondria. Pink1 then accumulates at the outer membrane and recruits Parkin. Mfn2 phosphorylated by Pink1 acts as a receptor for Parkin on mitochondria. Parkin ubiquitinates multiple proteins of the outer membrane. These ubiquitinated proteins are recognized by p62, a ubiquitin- and microtubule-associated protein 1 light chain 3 (LC3)-binding adaptor protein, followed by mitochondrial autophagy.

cells, the process seems more complex, and there are several ways for damaged mitochondria to be recognized by phagophores. As in yeast, one mechanism uses either proteins or lipids on the mitochondrial membrane as receptors for LC3. To date, a mammalian homolog of Atg32 has not been reported. As introduced further, however, NIP3-like protein X (Nix/Bnip3L), BCL2/adenovirus E1B 19kDa interacting protein 3 (Bnip3), FUN14 domain containing 1 (FUNDC1), and Cardiolipin function as receptors for LC3 (Figure 2).^{53–57} Another mechanism uses ubiquitination of mitochondrial proteins followed by interaction with the adaptor proteins connecting ubiquitin with LC3. The representative case is the PTEN-inducible putative kinase 1 (Pink1)–Parkin pathway for the ubiquitination of mitochondrial proteins, with sequestosome 1 (p62) and neighbor of BRCA1 gene 1 acting as adaptor proteins (Figure 3).^{58,59} In addition, CK2, which phosphorylates Atg32 in yeast, has been reported to phosphorylate p62 at Ser 403, and this phosphorylation increases the affinity between its ubiquitin-associated domain and polyubiquitination chains on mitochondrial proteins in mammalian cells.⁶⁰ However, the involvement of this machinery in mitochondrial autophagy has not yet been validated.⁶⁰ The roles of key molecules in mediating mitophagy are discussed below.

Pink1, Parkin, and p62

Pink1- and Parkin-mediated mitochondrial autophagy is currently the most well established mechanism mediating mitophagy in mammalian cells.⁶¹ Both Pink1 and Parkin are linked to the pathogenesis of autosomal recessive juvenile Parkinsonism.⁶² Pink1 is a serine/threonine kinase that has a mitochondria-targeting signal in its N-terminus.⁶³ After import into mitochondria through translocase of outer membrane and translocase of inner membrane complexes, Pink1 is anchored at the IMM.⁶³ In intact mitochondria, matrix processing peptidase and presenilin-associated rhomboid-like continuously degrade Pink1.^{64,65} However, in depolarized mitochondria, the import to the IMM is inhibited and Pink1 accumulates at the OMM. Pink1 then forms a 700 kDa complex with translocase of outer membrane and undergoes autophosphorylation at Ser 228 and Ser 402.^{66,67} Pink1 is also accumulated on the OMM in response to an increase in unfolded proteins in mitochondria, thereby playing an essential role in mediating mitophagic removal of polarized mitochondria.⁶⁸ Activated Pink1 recruits Parkin, a cytosolic E3 ubiquitin ligase, to damaged mitochondria and promotes their degradation through phosphorylation of multiple substrates. Pink1 phosphorylates Mfn2, which in turn acts as a mitochondrial receptor for Parkin in cardiomyocytes.⁶⁹ A defect in Parkin translocation was observed in Mfn2-deficient neurons,⁷⁰ suggesting that Mfn2 plays a critical role in recruiting Parkin to mitochondria in multiple cell types. However, it should be noted that the translocation of Parkin to mitochondria can still be observed in mouse embryonic fibroblasts lacking Mfn1 and Mfn2, suggesting the existence of an alternative or compensatory mechanism for recruiting Parkin.⁷¹ Stable expression of Pink1 anchored to the outer membrane can induce mitochondrial accumulation of Parkin, followed by mitochondrial autophagy in a manner independent of membrane potential.⁶⁴ Ectopic expression of Pink1 in peroxisomes recruits Parkin

to peroxisomes, accompanied by ubiquitination and selective autophagic clearance of peroxisomes.⁶⁶ These results suggest that Pink1 acts downstream or independently of mitochondrial depolarization. Although initial studies suggested that Pink1 phosphorylates Parkin in its ubiquitin-like domain, thereby stimulating the E3 ligase activity of Parkin,^{72,73} mutations of conserved serine/threonine residues in Parkin do not completely suppress its activity, suggesting that Pink1 has additional substrates through which it activates Parkin. In fact, Pink1 phosphorylates ubiquitin conjugated to mitochondrial proteins at Ser 65,^{74–76} which in turn directly activates Parkin and promotes further ubiquitination of mitochondrial proteins.⁷⁷ Multiple mitochondrial proteins, including voltage-dependent anion channel, Mfn1/2, translocase of outer membrane, fission 1, ras homolog family member T1/2, and mitochondrial hexokinase, have been identified as substrates of Parkin.^{46,78–80} It has been shown that Parkin-labeled mitochondria are removed by autophagy bit-by-bit, which occurs where Parkin-labeled mitochondria and endoplasmic reticulum intersect.⁸¹ Polyubiquitination catalyzed by Parkin leads to recruitment of p62, a ubiquitin- and LC3-binding adaptor protein, followed by perinuclear clustering of damaged mitochondria.^{59,78,82,83} However, whether p62 is essential for Parkin-mediated mitochondrial autophagy is controversial.^{59,82,83} Although degradation of the IMM and matrix proteins occurs through mitochondrial autophagy, degradation of OMM proteins can take place also through a proteasome-dependent mechanism mediated by Parkin.⁴⁶ Parkin promotes recruitment of the AAA ATPase valosin containing protein, which extracts ubiquitinated proteins for proteasomal degradation, to damaged mitochondria.⁸⁴ Parkin and Pink1 are also involved in the removal of damaged mitochondria through a vesicular trafficking pathway in which mitochondria-derived vesicles containing mitochondrial proteins are transported to lysosomes for degradation.^{85–87} These findings account for the difficulty in distinguishing typical mitochondrial autophagy from other forms of mitochondrial protein degradation based on Pink1–Parkin dependency alone.

Pink1–Parkin-mediated mitophagy is negatively regulated by USP30, a deubiquitinase localized in the OMM, by removing ubiquitin from substrates ubiquitinated by Parkin⁸⁸ and Clec16a, a membrane-associated endosomal protein which promotes proteasomal degradation of Parkin through Nrdp1, another E3 ubiquitin ligase. Tumor protein p53 (p53) inhibits mitophagy by sequestering Parkin in the cytosol,⁸⁹ whereas b-cell leukemia/lymphoma 2 protein (Bcl-2) family proteins, including Bcl-xL, MCL-1, and Bcl-2, on the OMM, inhibit mitophagy through inhibition of Parkin translocation to depolarized mitochondria.⁹⁰

Besides mitophagy and the mitochondria-derived vesicles pathway, Parkin and Pink1 also affect the quality control of mitochondria through several other mechanisms.⁶² For example, Parkin and Pink1 affect mitochondrial motility through Pink1-mediated phosphorylation and Parkin-mediated proteasomal degradation of Miro, an OMM protein involved in mitochondrion trafficking. Stabilization of Pink1 and Parkin activation immobilize damaged mitochondria through degradation of Miro, which in turn facilitates the removal of damaged mitochondria through mitophagy.⁹¹ Parkin positively

regulates mitochondrial biogenesis through proteasomal degradation of PARIS, a zinc-finger protein, which suppresses transcription of peroxisome proliferator-activated receptor- γ coactivator-1 α , a positive regulator of mitochondrial biogenesis.⁹² Parkin also regulates fat uptake by controlling the stability of CD36 through ubiquitination in hepatocytes.⁹³ These studies suggest that Parkin contributes in diverse ways to mitochondrial quality control and metabolism.

Although many molecules of the autophagic machinery were discovered in yeast, p62 is absent in lower eukaryotes.⁹⁴ Recently, Lu et al discovered CUE (coupling of ubiquitin endoplasmic reticulum degradation)-domain targeting proteins, a new class of ubiquitin-Atg8/LC3 adaptors.⁹⁴ The CUET proteins are Cue5p in yeast and toll interacting protein in human.⁹⁴ Although it differs in its domain arrangement, toll interacting protein fulfills both the biochemical and genetic criteria for being a functional human homolog of yeast Cue5p.⁹⁴ These proteins interact with Atg8 (LC3) through Atg8 family interacting motif (also termed as LC3-interacting region [LIR]) and with ubiquitin through a CUE domain but not through a ubiquitin-associated domain like p62 or neighbor of BRCA1 gene 1.⁹⁴ Toll interacting protein binds to ubiquitin conjugates more tightly than p62 and is more effective at clearing huntington poly-glutamine protein than p62.⁹⁴ Toll interacting protein seems to specifically target highly aggregation-prone proteins for autophagy, but its role in organelle-selective autophagy has not yet been clarified.⁹⁴

Nix/Bnip3L and Bnip3

To date, some mitochondria-labeling proteins, which also bind to LC3 or LC3 homologs through the LIR as a mitochondrial receptor for autophagosomes, have been identified in mammalian cells. Nix/Bnip3L and Bnip3 are included among these OMM proteins.^{53–55} They are BH3-only proteins and proapoptotic members of the Bcl-2 family.^{95–97} The protein level of Nix/Bnip3L is increased in the final stage of erythroid maturation, and genetic ablation of Nix/Bnip3L prevents the loss of mitochondrial membrane potential and sequestration of mitochondria into autophagosomes.⁹⁸ Treatment with carbonyl cyanide *m*-chlorophenyl hydrazine, an uncoupling chemical, or BH3 mimetic induces a loss of membrane potential and restores mitochondrial autophagy in Nix/Bnip3L^{-/-} erythroids, suggesting that Nix/Bnip3L-mediated loss of membrane potential is important for mitochondrial autophagy.⁹⁸ Nix/Bnip3L regulates Parkin translocation to mitochondria.⁹⁹ Mutation of the LIR of Nix/Bnip3L results in a partial reduction of mitochondrial autophagy in reticulocytes.⁵⁴

Kubli et al reported that oxidative stress-induced homodimerization of Bnip3 is observed after ischemia/reperfusion (IR) in the heart and that this dimerization correlates with its activation.¹⁰⁰ Bnip3 induces extensive fragmentation and autophagy of mitochondria in HL-1 cells, and this phenomenon is independent of mitochondrial permeability transition pore opening or of Bcl-2-associated X protein/Bcl-2-antagonist/killer 1.^{101,102} The mitochondrial fragmentation induced by Bnip3 occurs through its direct inhibition of optic atrophy 1 (Opa1), a fusion protein.¹⁰³ Recent evidence showed that phosphorylation of the LIR in Bnip3 promotes the interaction

between Bnip3 and LC3B, thereby inducing mitochondrial autophagy.¹⁰⁴ However, the responsible kinase has not yet been identified. Nix and Bnip3 promote apoptosis of cardiomyocytes through stimulation of OMM permeabilization.^{105,106} Nix also promotes necrosis of cardiomyocytes by stimulating endoplasmic reticulum-mitochondria crosstalk, Ca²⁺ overload, and consequent mitochondrial permeability transition pore opening.¹⁰⁷ Bnip3 also stimulates myocardial necrosis in response to doxorubicin treatment by disrupting formation of mitochondrial protein complexes between the key respiratory proteins.¹⁰⁸ Thus, what determines the role of Bnip3 in promoting cell death or survival through mitochondrial autophagy remains to be elucidated in the heart.

FUNDC1

FUNDC1 is a mitochondrial outer membrane protein that integrates with LC3 through its LIR under hypoxic conditions.⁵⁶ During starvation, mitochondrial clearance does not depend on FUNDC1, suggesting that it is specific for hypoxia-induced mitochondrial autophagy.⁵⁶ FUNDC1 is expressed under normoxia; however, the interaction between FUNDC1 and LC3 is stabilized under hypoxic conditions, which is attributable to dephosphorylation at tyrosine 18 during hypoxia. FUNDC1 also acts as a mitochondrial receptor for unc-51-like autophagy activating kinase 1.¹⁰⁹ unc-51-like autophagy activating kinase 1 translocates to depolarized mitochondria and phosphorylates FUNDC1 at Ser 17, thereby promoting mitochondrial autophagy.¹⁰⁹ Mitochondrially localized phosphoglycerate mutase family member 5 phosphatase directly dephosphorylates FUNDC1 at Ser 13 on hypoxia or carbonyl cyanide-*p*-trifluoromethoxyphenylhydrazine treatment.¹¹⁰ This dephosphorylation of FUNDC1 promotes its interaction with LC3 and mitochondrial autophagy, whereas CK2 phosphorylates the same residue of FUNDC1, thereby negatively regulating mitochondrial autophagy.¹¹⁰ CK2 phosphorylates Atg32 and promotes mitophagy in yeast. However, the effect of phosphorylation of FUNDC1 is opposite to that of phosphorylation of Atg32.

Cardiolipin

Cardiolipin, a phospholipid of the IMM, has been reported to translocate to the OMM in response to mitochondrial injury and interact directly with LC3 in primary cortical neurons and SH-SY5H cells.⁵⁷ The inhibition of cardiolipin synthase or of phospholipid scramblase-3 reduces the delivery of mitochondria to autophagosomes.⁵⁷ Mutation of the residues predicted to be cardiolipin interaction sites in LC3 inhibits mitochondrial autophagy.⁵⁷ Given that oxidized cardiolipin promotes the release of proapoptotic factors into the cytosol, Chu et al speculated that externalized cardiolipin undergoes oxidation and accelerates apoptosis in the event of unsuccessful clearance of damaged mitochondria.⁵⁷

Overexpression of acyl-CoA:monolysocardiolipin acyltransferase, an enzyme that catalyzes pathological remodeling of cardiolipin, stimulates oxidative stress and clearance of mitochondria via autophagy in H9c2 cells. ALCAT knockout mice show upregulation of Pink1, and both oxidative stress and mitochondrial dysfunction in response to L-tyrosine-induced stress are attenuated in these mice. Whether cardiolipin

peroxidation directly affects mitochondrial autophagy or mitochondrial autophagy is regulated secondarily by mitochondrial dysfunction/oxidation remains to be elucidated.¹¹¹

General Control of Amino Acid Synthesis 5-Like 1

Increasing lines of evidence suggest that lysine acetylation controls many cellular functions, including autophagy, in response to metabolic stress.¹¹² Acetyl-CoA, an intermediate product of metabolic pathways whose acetyl moiety is transferred for protein acetylation, is a negative regulator of autophagy.¹¹³ During starvation, the level of acetyl-CoA is thought to be diminished. In addition, general control of amino acid synthesis 5-like 1 (GCN5L1), a component of the mitochondrial acetyltransferase machinery, is downregulated, whereas sirtuin 3, a mitochondrial deacetylase, is activated in HepG2 cells.¹¹⁴ Downregulation of GCN5L1 diminishes acetylation of mitochondrial proteins and stimulates mitochondrial autophagy, which, in turn, slows down respiration and induces stress-resilience in cells.¹¹⁴ This mitochondrial autophagy is both Atg5- and p62-dependent but Parkin-independent, and thus, whether autophagy regulated by GCN5L1 is mitochondria-selective is currently unknown. One mechanism through which the reduced protein acetylation induces mitochondrial autophagy is stimulation of transcription factor EB, a master regulator of lysosomal biogenesis. Interestingly, peroxisome proliferator-activated receptor- γ coactivator-1 α , a key regulator of mitochondrial biogenesis, is upregulated in parallel, so that mitochondrial turnover is stimulated without diminishing the total number of mitochondria.¹¹⁵ Whether such a mechanism also exists in cardiomyocytes and, if so, when it is activated and to what extent it is involved in the clearance of damaged mitochondria in the heart remains to be elucidated. In theory, mitophagy and mitochondrial biogenesis should be coupled. If so, investigating the underlying mechanisms, particularly the role of transcription factor EB in regulating peroxisome proliferator-activated receptor- γ coactivator-1 α/β , NRF1/2 and Tfam, in cardiomyocytes is of great interest.

Mitochondrial Fission and Fusion Proteins

Mitochondria are highly dynamic organelles and their morphology changes continuously through fusion and fission.¹¹⁶ The fusion of mitochondria is regulated by Mfn1 and Mfn2 localized on the OMM and Opa1 on the IMM, whereas fission is regulated by Drp1 (dynamin 1-like), fission 1, and Mif1.¹¹⁷ Although dynamin-related GTPase DNMI1, a yeast homolog of mammalian Drp1, plays an important role in mediating mitophagy in yeast,¹¹⁸ whether fission is prerequisite for mitochondrial autophagy and, if so, how it is coupled to autophagy are not fully understood in mammalian cells.^{11,41} Twig et al reported that fission events often generate uneven daughter mitochondria and that the one with reduced membrane potential has a reduced probability to fuse in INS1 cells.¹¹ The daughter mitochondria that are unable to fuse are characterized by decreased expression levels of Opa1. Inhibition of fission through a dominant negative form of Drp1 or knockdown of fission 1, a factor that is involved in recruiting Drp1

to mitochondria, inhibits mitochondrial autophagy and causes accumulation of dysfunctional mitochondria.¹¹ This study suggests that mitochondrial fission is important to segregate dysfunctional mitochondria for their removal through autophagy. Mouse embryonic fibroblast cells deficient in Opa1 or Mfn1/2 exhibit an increase in fragmented mitochondria and mitochondrial autophagy during starvation, whereas downregulation of Drp1 induces unopposed fusion of mitochondria and prevents mitochondrial autophagy.¹¹⁹ Parkin promotes mitochondrial fission through ubiquitination and degradation of Mfn1 and Mfn2, which, in turn, leads to increases in mitophagy.¹²⁰ Mild and transient oxidative stress induces mitophagy, but not nonselective autophagy, in a Drp1-dependent manner in HeLa cells,¹²¹ and overexpression of fission 1 stimulates mitochondrial fragmentation and mitophagy in mouse embryonic fibroblast cells.¹²² These results are all consistent with the notion that mitochondrial autophagy couples to mitochondrial fission. However, it is possible that interventions used to alter mitochondrial fission and fusion could also directly or indirectly affect mitochondrial autophagy independently of the fission and fusion.¹²³ For example, Drp1 affects general autophagy through its interaction with Bcl-xL in cardiomyocytes.¹²⁴ Even if mitochondrial fission directly affects autophagy, the molecular mechanisms by which fission mediates mitochondrial autophagy remain to be elucidated.

Genetic manipulation of proteins involved in mitochondrial fission/fusion induces both morphological changes and functional impairment in mitochondria in cardiomyocytes. Mitochondrial fragmentation and autophagy induced by overexpression of Bnip3 were both inhibited in the presence of dominant negative Drp1 in adult ventricular cardiomyocytes.¹²⁵ Python mutant mice, which have a single mutation in exon 11 of the *Drp1* gene, exhibit elongated mitochondria, suggesting impairment of mitochondrial fission.¹²⁶ These mice show an $\approx 50\%$ reduction in adenosine triphosphate level in the heart and develop dilated cardiomyopathy.¹²⁶ Although basal cardiac function is impaired in cardiac-specific Drp1 KO mice, a consensus has not been reached as to whether removal of dysfunctional mitochondria by autophagy is reduced in these mice. Ikeda et al reported that genetic deletion of Drp1 suppresses both general autophagy and mitochondrial autophagy in the mouse heart in vivo.¹²⁴ Kageyama et al reported that Drp1 is required for Parkin-independent mitochondrial autophagy in the heart and that Drp1 and Parkin act synergistically to promote mitochondrial homeostasis in the brain.⁴⁸ On the other hand, Song et al showed that Drp1 ablation increases mitophagy and causes generalized loss of mitochondria.¹²⁷

The role of mitochondrial fusion in regulating autophagy and cardiac function seems more complex. Although homozygous mutation of Opa1 is embryonic lethal,¹²⁸ heterozygous Opa1^{+/-} mice exhibit late onset of cardiomyopathy, with a reduced mtDNA copy number and mitochondrial dysfunction.¹²⁸ Whether mitochondrial autophagy is enhanced in this model is unknown. Downregulation of Mfn1 increased the number of small, spherical mitochondria,¹²⁹ whereas downregulation of Mfn2 increased pleiomorphic and enlarged mitochondria in cardiomyocytes.¹³⁰ In both cases, the cardiomyocytes were protected against stress, although another report showed that downregulation of Mfn2

induces mitochondrial dysfunction.⁶⁹ Combined ablation of Mfn1 and Mfn2 in the adult heart induces mitochondrial fragmentation and respiratory dysfunction, resulting in dilated cardiomyopathy.^{131,132} Downregulation of Mfn2 inhibits Parkin-mediated mitophagy, whereas unopposed fission caused by either single or combined downregulation of Mfn1 and Mfn2 may positively affect nonselective autophagy.¹³³ Thus, the overall effects of Mfn1/2 downregulation on mitochondrial clearance by autophagy is unclear. Furthermore, Bhandari et al demonstrated that suppression of mitochondrial fusion rescues the cardiomyopathy induced by Parkin deficiency in *Drosophila*.¹³⁴ These studies indicate that mitochondrial fusion contributes to the cardiac dysfunction of Parkin deficiency by promoting contamination of the healthy mitochondrial pool with damaged mitochondria, independently of mitophagy.

Atypical Forms of Mitochondrial Clearance

Macroautophagy is separated into several specific steps, including induction, recognition, and selection of cytoplasmic substrates, formation of the autophagosome around substrates, autophagosome-lysosome fusion, degradation of the autolysosomal contents, and release of the degradation products into the cytoplasm. The canonical, or conventional, autophagic pathway consists of evolutionarily conserved signaling molecules encoded by Atgs, including Atg4, Atg5, Beclin1 (Atg6), Atg7, Atg12, and Atg16, that govern these steps. On the other hand, increasing lines of evidence suggest that noncanonical autophagic pathways may also be present. Nishida et al revealed that Atg5^{-/-}Atg7^{-/-} double-knockout cells are still able to form autophagosomes and degrade autophagic substrates inside autolysosomes in response to certain stimuli.⁴² During this process of Atg5/Atg7-independent autophagy, termed alternative autophagy, lipidation of LC3 does not occur. Instead, Rab9, a small GTPase involved in membrane trafficking and fusion between the trans-Golgi network and late endosomes, plays a critical role in generating autophagosomes in the alternative autophagic pathway by promoting fusion of the phagophore with vesicles derived from the trans-Golgi network and late endosomes. Recent evidence suggests that unc-51 like autophagy activating kinase 1-dependent, Atg5-independent macroautophagy is the dominant process for removing mitochondria from reticulocytes in the final stage of erythrocyte maturation.⁴⁴ Oxidative stress induces a vesicular transport pathway that selectively removes mitochondrial proteins for delivery to the lysosomes in COS7 cells. This mechanism does not require mitochondrial depolarization and is independent of Atg5 and LC3, indicating that it is distinct from either autophagy or selective mitophagy.⁸⁵ However, the functional significance of alternative autophagy has not yet been demonstrated in the heart. Although conventional macroautophagy is attenuated in Drp1 knockout (KO) mice, Parkin-mediated mitophagy may be stimulated.¹²⁷ Thus, it is possible that multiple mechanisms may mediate degradation of damaged mitochondria.

What Is the Role of Mitophagy or Mitochondrial Autophagy in the Heart at Baseline?

Conditional cardiac-specific *atg5* KO mice, which serve as a loss-of-function model for general autophagy, develop cardiac

dysfunction, suggesting that general autophagy plays an essential role in maintaining the physiological function of the adult heart at baseline.¹³⁵ These mice exhibit accumulation of misfolded proteins and damaged organelles, including mitochondria, consistent with the notion that protein/organelle quality control mediated by autophagy is essential for the heart even under resting conditions. Interestingly, *beclin1*^{+/-} mice, another loss-of-function model of autophagy, do not exhibit any significant baseline cardiac phenotype, despite the fact that stress-induced autophagy is significantly inhibited.¹³⁶ The difference in baseline cardiac phenotype in these mice may be caused by a difference in the level or the timing of autophagy suppression. Atg5 and Beclin1 (Atg6) may regulate qualitatively different types of autophagy.⁴² It is also possible that the development of compensatory mechanisms may vary depending on which Atg is downregulated. Alternatively, the difference may be related to the fact that autophagy-related genes also mediate functions other than autophagy in cells. These issues represent the difficulty and limitations of elucidating the functional significance of mitochondrial autophagy or mitophagy in the heart using a single genetically altered mouse model.

The aforementioned studies suggest the importance of general autophagy in the maintenance of protein/organelle quality control and cardiac function at baseline. However, whether mitochondrial autophagy or mitophagy is critically involved in this process remains unknown. Although Parkin and Pink1 regulate multiple cellular functions besides mitophagy,⁶² their direct involvement in mitophagy means that loss-of-function studies of Parkin and Pink1 can provide valuable information regarding the functional involvement of Pink1-Parkin-mediated mitophagy in the heart. In fact, increasing lines of evidence suggest that Pink1-Parkin-mediated autophagy participates in mitochondrial quality control and the maintenance of cardiac function in the heart at baseline. Cardiomyocytes in which Bnip3 is overexpressed are able to eliminate damaged mitochondria through autophagy via a Parkin-dependent mechanism, indicating that Parkin-mediated mitochondrial autophagy exists in cardiomyocytes.¹²⁵ Parkin knockout mice exhibit normal cardiac function at baseline, but their cardiomyocytes exhibit morphologically disorganized mitochondria with or without mitochondrial dysfunction.^{137,138} Interestingly, clearance of mitochondria by autophagy seems to be maintained at baseline in these mice. This may be as a result of a low rate of mitochondrial degradation by autophagy at baseline, compensation for the lack of Parkin E3 ligase activity by other E3 ligases,¹³⁴ such as tumor necrosis factor-receptor-associated factor 2,¹³⁹ or compensation by Parkin-independent mechanisms of autophagy, such as general autophagy. Song et al suggested that the lack of Parkin-mediated mitophagy in heart-specific Mfn2 KO mice is compensated for by activation of nonselective autophagy in cardiomyocytes,¹³³ thereby allowing cardiomyocytes to maintain mitochondrial quality. This suggests that both mitophagy and general autophagy cooperatively participate in the removal of mitochondria in cardiomyocytes. Currently, molecular mechanisms by which mitochondria are degraded by Mfn2-Parkin-independent autophagy are unknown. However, it would be interesting to

clarify whether these 2 forms of autophagy degrade the same targets or qualitatively or quantitatively different populations of mitochondria, as well as to assess the relative contributions of mitophagy and general autophagy to overall mitochondrial quality control in cardiomyocytes. It should be noted that Parkin deficiency induces significant contractile dysfunction in the heart in *Drosophila*, suggesting that such crosstalk may not exist in lower organisms.¹³⁴ Aging-induced upregulation of p53 sequesters Parkin in the cytosol, thereby inhibiting mitophagy in mice. Because p53-deficient mice exhibit less of a decline in mitochondrial integrity and cardiac reserve during aging, it is possible that endogenous Parkin may play an important role in mediating mitophagy and that Parkin-mediated mitophagy declines with aging, in part because of p53.⁸⁹

Whether endogenous Pink1 plays an essential role in mediating mitophagy in the heart remains to be demonstrated. Because Pink1-mediated phosphorylation of Mfn2 leads to recruitment of Parkin and induction of mitophagy,⁶⁹ it is expected that downregulation of Pink1 inhibits mitophagy. Consistently, mouse hearts lacking Mfn2 develop cardiac dysfunction with age,⁶⁹ suggesting that mitophagy regulated by the Pink1-Mfn2 pathway plays an important role in maintaining cardiac function at baseline. Another study showed, however, that downregulation of Mfn2 exerts protective effects in the heart in response to cell death stimuli and I/R.¹³⁰ Thus, further investigation is required to clarify the role of Mfn2 in mediating mitophagy and mitochondrial quality control in the heart. Genetic deletion of Pink1 induces more severe cardiac effects than that of Parkin,¹⁴⁰ despite the fact that Pink1 and Parkin are thought to work together to mediate mitophagy. This may be because Pink1 has additional functions besides Parkin-mediated autophagy, including phosphorylation of TNF receptor-associated protein 1 (TRAP1), to protect against cell death.¹⁴¹

Role of Mitophagy or Mitochondrial Autophagy in Protection Against Cardiac Stress

The protective effect of mitophagy and mitochondrial autophagy in the heart generally seems more prominent during stress. In response to myocardial infarction (MI), activation of autophagy is observed most prominently in the border zone during the subacute phase (1 week after MI), returning to normal during the chronic phase (3 weeks after MI).¹⁴² The autophagic vacuoles in the border zone are large and contain organelles, including degraded mitochondria. In contrast, autophagy with normal-sized autophagosomes is upregulated in the remote zone during the chronic phase.¹⁴² Enhancing autophagy with rapamycin 2 weeks after coronary ligation ameliorates cardiac dysfunction and maladaptive remodeling, whereas inhibition of autophagy with bafilomycin A1 worsens them.¹⁴² These observations imply the existence of multiple forms of autophagy and suggest that some of them may play a role in mediating the clearance of mitochondria in a phase-dependent fashion after MI. A study using *beclin1*^{+/-} mice also showed that suppression of autophagy is detrimental during the chronic phase of MI. Hoshino et al reported that autophagy peaks during the acute phase of MI (several hours after coronary ligation)

in the border zone, as evaluated with GFP-LC3 transgenic mice.⁵⁰ In this model, mitochondrial autophagy, as indicated by the presence of autophagosomes containing mitochondria, also takes place during the same period.⁵⁰ MI induced by permanent coronary artery ligation upregulates p53 and TP53-induced glycolysis and apoptosis regulator in mice. Genetic deletion of p53 or TP53-induced glycolysis and apoptosis regulator in mice stimulates mitochondrial autophagy and inhibits accumulation of damaged mitochondria and apoptosis, an effect which is abolished by chloroquine treatment. The authors suggest that the beneficial effects of the p53 or TP53-induced glycolysis and apoptosis regulator downregulation may be mediated by mitophagy.⁵⁰ These results are consistent with the negative regulation of mitophagy by cytosolic p53.⁸⁹ Translocation of Parkin to mitochondria and ubiquitination of mitochondrial proteins occur during the acute phase of MI, suggesting that Parkin-mediated mitophagy is stimulated.¹³⁷ Parkin-deficient mice exhibit a decrease in mitochondrial autophagy, accumulation of dysfunctional mitochondria, and a reduced survival rate after MI.¹³⁷ Taken together, these results suggest that either mitophagy or mitochondrial autophagy is stimulated in the heart during the chronic phase of cardiac remodeling and that they are protective for the heart.

During myocardial reperfusion after a period of myocardial ischemia, mitochondria generate ROS, initiating a feed-forward mechanism of oxidative stress, mitochondrial injury, and cell death.¹⁴³ Selective elimination of damaged mitochondria by mitophagy or mitochondrial autophagy is predicted to protect cardiomyocytes during reperfusion. In fact, loss of Pink1 has been reported to increase the infarct size after I/R, although this study did not examine mitochondrial autophagy in the heart.¹⁴⁴ Ischemic preconditioning (IPC) induces the translocation of Parkin to mitochondria, and the IPC-mediated cardioprotective effect is abolished in Parkin KO mice.¹⁴⁵ The translocation of p62 to mitochondria is also impaired in these mice, suggesting that Parkin and p62 act together to induce mitochondrial autophagy in response to IPC.¹⁴⁵ Whether the cardioprotective effect of IPC requires mitophagy rather than the mitophagy-independent functions of Parkin remains to be elucidated. If mitochondrial autophagy does mediate the effect of IPC, investigating how preactivated autophagy protects the heart against I/R may allow identification of novel modalities of cardioprotection. Myocardial I/R injury is exacerbated in cardiac-specific *drp1*^{+/-} mice, in which mitochondrial fission and autophagy are both inhibited.¹²⁴ Thus, suppression of mitophagy and mitochondrial autophagy seems to be detrimental during I/R. Because translocation of Drp1 from the cytosol to mitochondria is stimulated by I/R, enhancement of the quality control mechanism seems to be essential for minimizing myocardial injury during reperfusion. On the other hand, it has been shown previously that Beclin1 haploinsufficiency suppresses autophagy and inhibits I/R injury.¹³⁶ Why haploinsufficiency of Drp1 and Beclin1 exhibit directionally opposite effects on I/R injury is currently unknown. However, one possibility is that Drp1 and Beclin1 may affect general autophagy and mitochondrial autophagy to different extents. Another, less attractive, possibility is that Drp1 and Beclin1

differentially affect cellular functions other than autophagy. Further investigation is required to address this issue.

There are fewer publications addressing mitochondrial autophagy during pressure overload. However, the protein level of Pink1 is known to be decreased after transverse aortic constriction. Because Pink1-deficient mice develop age-dependent hypertrophy and cardiac dysfunction accompanied by mitochondrial dysfunction even without pressure overload,¹⁴⁰ downregulation of Pink1 and consequent suppression of mitochondrial autophagy may be detrimental during pressure overload as well.

Perspective

In summary, the recent progress regarding molecular mechanisms of mitophagy, including how damaged mitochondria are recognized and how they are engulfed by autophagosomes, has greatly improved our understanding of quality control mechanisms in mitochondria. However, our knowledge regarding mitophagy in the heart and the cardiomyocytes therein is limited. First of all, whether mitochondria are degraded by mitophagy or general autophagy requires more studies, using more sensitive and accurate methods to monitor lysosomal degradation of mitochondrial proteins,⁴⁵ together with electron microscopic observation to determine whether autophagosomes contain mitochondria alone or other materials as well. Establishing more reliable methods to monitor mitochondrial autophagy would dramatically advance our understanding of the molecular signaling mechanisms of mitophagy/mitochondrial autophagy. In particular, whether the Pink1/Parkin pathway is involved in mitochondrial autophagy/mitophagy in cardiomyocytes in response to relevant stresses *in vivo* urgently requires more investigation. If the Pink1/Parkin pathway does play an important role in mitophagy/mitochondrial autophagy in cardiomyocytes during relevant stresses, then more investigation is required regarding interactions between the Pink1/Parkin pathway, the receptor-mediated mechanisms, including those mediated by Bnip3, Nix, FUNDC, and cardiolipin, and the protein kinases, such as unc-51 like autophagy activating kinase 1/2, AMP (adenosine monophosphate)-activated protein kinase, mTOR, and CK2. In addition, we cannot exclude the presence of novel mechanisms. For example, it is possible that novel functional homologs of yeast Atg32 and Atg11 may exist. Unbiased approaches are therefore essential to clarify how damaged mitochondria are recognized by autophagy.¹⁴⁶

Increasing lines of evidence suggest that autophagic degradation of mitochondria can be observed even when Parkin, Mfn2, or Drp1, molecules previously shown to mediate mitochondrial autophagy in the heart, are downregulated.^{127,133,138} Thus, mitochondria may be degraded through multiple mechanisms.⁸ If mitochondria are degraded by both mitophagy and general autophagy, identifying an intervention that could conveniently distinguish between these mechanisms would be helpful because general autophagy and mitophagy may have distinct functional significance. If mitochondria are also degraded by the alternative autophagy pathway identified by Nishida et al, some other convenient method would be required to monitor these autophagosomes because they are LC3-independent.⁴² When one mechanism of mitochondrial

degradation is inhibited, another mechanism may be activated to compensate for its loss.^{127,147} If this is shown to be the case, elucidating the interactions between the multiple mechanisms of autophagy may be of great interest.

Although it remains controversial as to whether uncontrolled autophagy can kill cells, a growing line of evidence suggests that autophagy may be detrimental under some conditions.^{148,149} One can speculate that excessive elimination of mitochondria may be detrimental for cardiomyocytes, but whether this can be caused by excessive mitochondrial autophagy or mitophagy and, if so, what would induce such a condition remain unknown.

Increasing lines of evidence suggest that autophagy is intimately involved in the survival and death of cardiomyocytes. It is likely that the beneficial effects of autophagy are partly mediated through elimination of damaged mitochondria and consequent prevention of mitochondrial dysfunction, oxidative stress, and cell death. To introduce modification of autophagy and mitophagy into the treatment of cardiovascular disease, further investigations are needed to better understand when autophagy is activated or inhibited and how it affects the function of mitochondria in response to a wide variety of cardiovascular stress conditions.

Acknowledgments

We thank Christopher D. Brady and Daniela Zablocki for critical reading of the article.

Sources of Funding

This work was supported in part by US Public Health Service Grants HL67724, HL91469, HL 112330, HL102738, and AG23039. This work was also supported by the Fondation Leducq Transatlantic Networks of Excellence.

Disclosures

None.

References

1. Wang Z, Jiang H, Chen S, Du F, Wang X. The mitochondrial phosphatase PGAM5 functions at the convergence point of multiple necrotic death pathways. *Cell*. 2012;148:228–243.
2. Subramanian N, Natarajan K, Clatworthy MR, Wang Z, Germain RN. The adaptor MAVS promotes NLRP3 mitochondrial localization and inflammasome activation. *Cell*. 2013;153:348–361.
3. Mukhopadhyay S, Panda PK, Sinha N, Das DN, Bhutia SK. Autophagy and apoptosis: where do they meet? *Apoptosis*. 2014;19:555–566. doi: 10.1007/s10495-014-0967-2.
4. Dai DF, Rabinovitch PS, Ungvari Z. Mitochondria and cardiovascular aging. *Circ Res*. 2012;110:1109–1124. doi: 10.1161/CIRCRESAHA.111.246140.
5. Oka T, Hikoso S, Yamaguchi O, Taneike M, Takeda T, Tamai T, Oyabu J, Murakawa T, Nakayama H, Nishida K, Akira S, Yamamoto A, Komuro I, Otsu K. Mitochondrial DNA that escapes from autophagy causes inflammation and heart failure. *Nature*. 2012;485:251–255. doi: 10.1038/nature10992.
6. Wallace DC. A mitochondrial paradigm of metabolic and degenerative diseases, aging, and cancer: a dawn for evolutionary medicine. *Annu Rev Genet*. 2005;39:359–407. doi: 10.1146/annurev.genet.39.110304.095751.
7. Wojtovich AP, Nadochiy SM, Brookes PS, Nehrke K. Ischemic preconditioning: the role of mitochondria and aging. *Exp Gerontol*. 2012;47:1–7. doi: 10.1016/j.exger.2011.11.001.
8. Sugiura A, McLelland GL, Fon EA, McBride HM. A new pathway for mitochondrial quality control: mitochondrial-derived vesicles. *EMBO J*. 2014;33:2142–2156. doi: 10.15252/embj.201488104.

9. Lemasters JJ. Selective mitochondrial autophagy, or mitophagy, as a targeted defense against oxidative stress, mitochondrial dysfunction, and aging. *Rejuvenation Res.* 2005;8:3–5. doi: 10.1089/rej.2005.8.3.
10. Zhang Y, Qi H, Taylor R, Xu W, Liu LF, Jin S. The role of autophagy in mitochondria maintenance: characterization of mitochondrial functions in autophagy-deficient *S. cerevisiae* strains. *Autophagy.* 2007;3:337–346.
11. Twig G, Elorza A, Molina AJ, et al. Fission and selective fusion govern mitochondrial segregation and elimination by autophagy. *EMBO J.* 2008;27:433–446. doi: 10.1038/sj.emboj.7601963.
12. Chiang HL, Terlecky SR, Plant CP, Dice JF. A role for a 70-kilodalton heat shock protein in lysosomal degradation of intracellular proteins. *Science.* 1989;246:382–385.
13. Agarrareres FA, Terlecky SR, Dice JF. An intralysosomal hsp70 is required for a selective pathway of lysosomal protein degradation. *J Cell Biol.* 1997;137:825–834.
14. Kraft C, Reggiori F, Peter M. Selective types of autophagy in yeast. *Biochim Biophys Acta.* 2009;1793:1404–1412. doi: 10.1016/j.bbamer.2009.02.006.
15. Komatsu M, Ichimura Y. Selective autophagy regulates various cellular functions. *Genes Cells.* 2010;15:923–933. doi: 10.1111/j.1365-2443.2010.01433.x.
16. Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter proteins. *Autophagy.* 2011;7:279–296.
17. Bolender RP, Weibel ER. A morphometric study of the removal of phenobarbital-induced membranes from hepatocytes after cessation of treatment. *J Cell Biol.* 1973;56:746–761.
18. Klionsky DJ, Cuervo AM, Dunn WA Jr, Levine B, van der Klei I, Seglen PO. How shall I eat thee? *Autophagy.* 2007;3:413–416.
19. Hutchins MU, Veenhuis M, Klionsky DJ. Peroxisome degradation in *Saccharomyces cerevisiae* is dependent on machinery of macroautophagy and the Cvt pathway. *J Cell Sci.* 1999;112 (Pt 22):4079–4087.
20. Bellu AR, Komori M, van der Klei IJ, Kiel JA, Veenhuis M. Peroxisome biogenesis and selective degradation converge at Pex14p. *J Biol Chem.* 2001;276:44570–44574. doi: 10.1074/jbc.M107599200.
21. Kiel JA, Komduur JA, van der Klei IJ, Veenhuis M. Macropexophagy in *Hansenula polymorpha*: facts and views. *FEBS Lett.* 2003;549:1–6.
22. Bernales S, McDonald KL, Walter P. Autophagy counterbalances endoplasmic reticulum expansion during the unfolded protein response. *PLoS Biol.* 2006;4:e423. doi: 10.1371/journal.pbio.0040423.
23. Kraft C, Deplazes A, Sohrmann M, Peter M. Mature ribosomes are selectively degraded upon starvation by an autophagy pathway requiring the Ubp3p/Bre5p ubiquitin protease. *Nat Cell Biol.* 2008;10:602–610. doi: 10.1038/ncb1723.
24. Kanki T, Wang K, Cao Y, Baba M, Klionsky DJ. Atg32 is a mitochondrial protein that confers selectivity during mitophagy. *Dev Cell.* 2009;17:98–109. doi: 10.1016/j.devcel.2009.06.014.
25. Okamoto K, Kondo-Okamoto N, Ohsumi Y. A landmark protein essential for mitophagy: Atg32 recruits the autophagic machinery to mitochondria. *Autophagy.* 2009;5:1203–1205.
26. Kanki T, Kurihara Y, Jin X, Goda T, Ono Y, Aihara M, Hirota Y, Saigusa T, Aoki Y, Uchiyama T, Kang D. Casein kinase 2 is essential for mitophagy. *EMBO Rep.* 2013;14:788–794. doi: 10.1038/embor.2013.114.
27. Lipatova Z, Shah AH, Kim JJ, Mulholland JW, Segev N. Regulation of ER-phagy by a Ypt/Rab GTPase module. *Mol Biol Cell.* 2013;24:3133–3144. doi: 10.1091/mbc.E13-05-0269.
28. Nazarko TY. Atg37 regulates the assembly of the pexophagic receptor protein complex. *Autophagy.* 2014;10:1348–1349. doi: 10.4161/auto.29073.
29. Schuck S, Gallagher CM, Walter P. ER-phagy mediates selective degradation of endoplasmic reticulum independently of the core autophagy machinery. *J Cell Sci.* 2014;127:4078–4088. doi: 10.1242/jcs.154716.
30. Farré JC, Krick R, Subramani S, Thumm M. Turnover of organelles by autophagy in yeast. *Curr Opin Cell Biol.* 2009;21:522–530. doi: 10.1016/j.ccb.2009.04.015.
31. Campbell CL, Thorsness PE. Escape of mitochondrial DNA to the nucleus in *ymel* yeast is mediated by vacuolar-dependent turnover of abnormal mitochondrial compartments. *J Cell Sci.* 1998;111 (Pt 16):2455–2464.
32. Nowikovsky K, Reipert S, Devenish RJ, Schweyen RJ. Mdm38 protein depletion causes loss of mitochondrial K⁺/H⁺ exchange activity, osmotic swelling and mitophagy. *Cell Death Differ.* 2007;14:1647–1656. doi: 10.1038/sj.cdd.4402167.
33. Farré JC, Manjithaya R, Mathewson RD, Subramani S. PpAtg30 tags peroxisomes for turnover by selective autophagy. *Dev Cell.* 2008;14:365–376. doi: 10.1016/j.devcel.2007.12.011.
34. Farré JC, Burkenroad A, Burnett SF, Subramani S. Phosphorylation of mitophagy and pexophagy receptors coordinates their interaction with Atg8 and Atg11. *EMBO Rep.* 2013;14:441–449. doi: 10.1038/embor.2013.40.
35. Kissová I, Deffieu M, Manon S, Camougrand N. Uth1p is involved in the autophagic degradation of mitochondria. *J Biol Chem.* 2004;279:39068–39074. doi: 10.1074/jbc.M406960200.
36. Tal R, Winter G, Ecker N, Klionsky DJ, Abeliovich H. Aup1p, a yeast mitochondrial protein phosphatase homolog, is required for efficient stationary phase mitophagy and cell survival. *J Biol Chem.* 2007;282:5617–5624. doi: 10.1074/jbc.M605940200.
37. Mao K, Wang K, Liu X, Klionsky DJ. The scaffold protein Atg11 recruits fission machinery to drive selective mitochondrial degradation by autophagy. *Dev Cell.* 2013;26:9–18. doi: 10.1016/j.devcel.2013.05.024.
38. Brauer MJ, Saldanha AJ, Dolinski K, Botstein D. Homeostatic adjustment and metabolic remodeling in glucose-limited yeast cultures. *Mol Biol Cell.* 2005;16:2503–2517. doi: 10.1091/mbc.E04-11-0968.
39. Hayashi M, Raimondi A, O'Toole E, Paradise S, Collesi C, Cremona O, Ferguson SM, De Camilli P. Cell- and stimulus-dependent heterogeneity of synaptic vesicle endocytic recycling mechanisms revealed by studies of dynamin 1-null neurons. *Proc Natl Acad Sci U S A.* 2008;105:2175–2180. doi: 10.1073/pnas.0712171105.
40. Hayashi-Nishino M, Fujita N, Noda T, Yamaguchi A, Yoshimori T, Yamamoto A. A subdomain of the endoplasmic reticulum forms a cradle for autophagosome formation. *Nat Cell Biol.* 2009;11:1433–1437. doi: 10.1038/ncb1991.
41. Kim I, Lemasters JJ. Mitochondrial degradation by autophagy (mitophagy) in GFP-LC3 transgenic hepatocytes during nutrient deprivation. *Am J Physiol Cell Physiol.* 2011;300:C308–C317. doi: 10.1152/ajpcell.00056.2010.
42. Nishida Y, Arakawa S, Fujitani K, Yamaguchi H, Mizuta T, Kanaseki T, Komatsu M, Otsu K, Tsujimoto Y, Shimizu S. Discovery of Atg5/Atg7-independent alternative macroautophagy. *Nature.* 2009;461:654–658. doi: 10.1038/nature08455.
43. Zhang J, Randall MS, Loyd MR, Dorsey FC, Kundu M, Cleveland JL, Ney PA. Mitochondrial clearance is regulated by Atg7-dependent and -independent mechanisms during reticulocyte maturation. *Blood.* 2009;114:157–164. doi: 10.1182/blood-2008-04-151639.
44. Honda S, Arakawa S, Nishida Y, Yamaguchi H, Ishii E, Shimizu S. Ulk1-mediated Atg5-independent macroautophagy mediates elimination of mitochondria from embryonic reticulocytes. *Nat Commun.* 2014;5:4004. doi: 10.1038/ncomms5004.
45. Katayama H, Kogure T, Mizushima N, Yoshimori T, Miyawaki A. A sensitive and quantitative technique for detecting autophagic events based on lysosomal delivery. *Chem Biol.* 2011;18:1042–1052. doi: 10.1016/j.chembiol.2011.05.013.
46. Yoshii SR, Kishi C, Ishihara N, Mizushima N. Parkin mediates proteasome-dependent protein degradation and rupture of the outer mitochondrial membrane. *J Biol Chem.* 2011;286:19630–19640. doi: 10.1074/jbc.M110.209338.
47. Hernandez G, Thornton C, Stotland A, Lui D, Sin J, Ramil J, Magee N, Andres A, Quarato G, Carreira RS, Sayen MR, Wolkowicz R, Gottlieb RA. MitoTimer: a novel tool for monitoring mitochondrial turnover. *Autophagy.* 2013;9:1852–1861. doi: 10.4161/auto.26501.
48. Kageyama Y, Hoshijima M, Seo K, Bedja D, Sysa-Shah P, Andrabi SA, Chen W, Höke A, Dawson VL, Dawson TM, Gabrielson K, Kass DA, Iijima M, Sesaki H. Parkin-independent mitophagy requires Drp1 and maintains the integrity of mammalian heart and brain. *EMBO J.* 2014;33:2798–2813. doi: 10.15252/embj.201488658.
49. Edwards JG. Quantification of mitochondrial DNA (mtDNA) damage and error rates by real-time QPCR. *Mitochondrion.* 2009;9:31–35. doi: 10.1016/j.mito.2008.11.004.
50. Hoshino A, Matoba S, Iwai-Kanai E, Nakamura H, Kimata M, Nakaoka M, Katamura M, Okawa Y, Ariyoshi M, Mita Y, Ikeda K, Ueyama T, Okigaki M, Matsubara H. p53-TIGAR axis attenuates mitophagy to exacerbate cardiac damage after ischemia. *J Mol Cell Cardiol.* 2012;52:175–184. doi: 10.1016/j.yjmcc.2011.10.008.
51. Chen H, Vermulst M, Wang YE, Chomyn A, Prolla TA, McCaffery JM, Chan DC. Mitochondrial fusion is required for mtDNA stability in skeletal muscle and tolerance of mtDNA mutations. *Cell.* 2010;141:280–289. doi: 10.1016/j.cell.2010.02.026.
52. Larsen S, Nielsen J, Hansen CN, Nielsen LB, Wibrand F, Stride N, Schroder HD, Boushel R, Helge JW, Dela F, Hey-Mogensen M. Biomarkers of mitochondrial content in skeletal muscle of healthy young human subjects. *J Physiol.* 2012;590:3349–3360. doi: 10.1113/jphysiol.2012.230185.
53. Schwarten M, Mohrlüder J, Ma P, Stoldt M, Thielmann Y, Stangler T, Hersch N, Hoffmann B, Merkel R, Willbold D. Nix directly binds to GABARAP: a possible crosstalk between apoptosis and autophagy. *Autophagy.* 2009;5:690–698.

54. Novak I, Kirkin V, McEwan DG, Zhang J, Wild P, Rozenknop A, Rogov V, Löhr F, Popovic D, Occhipinti A, Reichert AS, Terzic J, Dötsch V, Ney PA, Dikic I. Nix is a selective autophagy receptor for mitochondrial clearance. *EMBO Rep*. 2010;11:45–51. doi: 10.1038/embor.2009.256.
55. Hanna RA, Quinsay MN, Orogo AM, Giang K, Rikka S, Gustafsson ÅB. Microtubule-associated protein 1 light chain 3 (LC3) interacts with Bnip3 protein to selectively remove endoplasmic reticulum and mitochondria via autophagy. *J Biol Chem*. 2012;287:19094–19104. doi: 10.1074/jbc.M111.322933.
56. Liu L, Feng D, Chen G, et al. Mitochondrial outer-membrane protein FUNDC1 mediates hypoxia-induced mitophagy in mammalian cells. *Nat Cell Biol*. 2012;14:177–185. doi: 10.1038/ncb2422.
57. Chu CT, Ji J, Dagda RK, et al. Cardiolipin externalization to the outer mitochondrial membrane acts as an elimination signal for mitophagy in neuronal cells. *Nat Cell Biol*. 2013;15:1197–1205. doi: 10.1038/ncb2837.
58. Kirkin V, Lamark T, Sou YS, et al. A role for NBR1 in autophagosomal degradation of ubiquitinated substrates. *Mol Cell*. 2009;33:505–516. doi: 10.1016/j.molcel.2009.01.020.
59. Geisler S, Holmström KM, Skujat D, Fiesel FC, Rothfuss OC, Kahle PJ, Springer W. PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell Biol*. 2010;12:119–131. doi: 10.1038/ncb2012.
60. Matsumoto G, Wada K, Okuno M, Kurosawa M, Nukina N. Serine 403 phosphorylation of p62/SQSTM1 regulates selective autophagic clearance of ubiquitinated proteins. *Mol Cell*. 2011;44:279–289. doi: 10.1016/j.molcel.2011.07.039.
61. Youle RJ, Narendra DP. Mechanisms of mitophagy. *Nat Rev Mol Cell Biol*. 2011;12:9–14. doi: 10.1038/nrm3028.
62. Scarffe LA, Stevens DA, Dawson VL, Dawson TM. Parkin and PINK1: much more than mitophagy. *Trends Neurosci*. 2014;37:315–324. doi: 10.1016/j.tins.2014.03.004.
63. Jin SM, Lazarou M, Wang C, Kane LA, Narendra DP, Youle RJ. Mitochondrial membrane potential regulates PINK1 import and proteolytic destabilization by PARL. *J Cell Biol*. 2010;191:933–942. doi: 10.1083/jcb.201008084.
64. Narendra DP, Jin SM, Tanaka A, Suen DF, Gautier CA, Shen J, Cookson MR, Youle RJ. PINK1 is selectively stabilized on impaired mitochondria to activate Parkin. *PLoS Biol*. 2010;8:e1000298. doi: 10.1371/journal.pbio.1000298.
65. Greene AW, Grenier K, Aguilera MA, Muise S, Farazifard R, Haque ME, McBride HM, Park DS, Fon EA. Mitochondrial processing peptidase regulates PINK1 processing, import and Parkin recruitment. *EMBO Rep*. 2012;13:378–385. doi: 10.1038/embor.2012.14.
66. Lazarou M, Jin SM, Kane LA, Youle RJ. Role of PINK1 binding to the TOM complex and alternate intracellular membranes in recruitment and activation of the E3 ligase Parkin. *Dev Cell*. 2012;22:320–333. doi: 10.1016/j.devcel.2011.12.014.
67. Okatsu K, Oka T, Iguchi M, et al. PINK1 autophosphorylation upon membrane potential dissipation is essential for Parkin recruitment to damaged mitochondria. *Nat Commun*. 2012;3:1016. doi: 10.1038/ncomms2016.
68. Jin SM, Youle RJ. The accumulation of misfolded proteins in the mitochondrial matrix is sensed by PINK1 to induce PARK2/Parkin-mediated mitophagy of polarized mitochondria. *Autophagy*. 2013;9:1750–1757. doi: 10.4161/auto.26122.
69. Chen Y, Dorn GW II. PINK1-phosphorylated mitofusin 2 is a Parkin receptor for culling damaged mitochondria. *Science*. 2013;340:471–475. doi: 10.1126/science.1231031.
70. Lee S, Sterky FH, Mourier A, Terzioglu M, Cullheim S, Olson L, Larsson NG. Mitofusin 2 is necessary for striatal axonal projections of midbrain dopamine neurons. *Hum Mol Genet*. 2012;21:4827–4835. doi: 10.1093/hmg/dds352.
71. Narendra D, Tanaka A, Suen DF, Youle RJ. Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol*. 2008;183:795–803. doi: 10.1083/jcb.200809125.
72. Kondapalli C, Kazlauskaitė A, Zhang N, Woodroof HI, Campbell DG, Gourlay R, Burchell L, Walden H, Macartney TJ, Deak M, Knebel A, Alessi DR, Muqit MM. PINK1 is activated by mitochondrial membrane potential depolarization and stimulates Parkin E3 ligase activity by phosphorylating Serine 65. *Open Biol*. 2012;2:120080. doi: 10.1098/rsob.120080.
73. Shiba-Fukushima K, Imai Y, Yoshida S, Ishihama Y, Kanao T, Sato S, Hattori N. PINK1-mediated phosphorylation of the Parkin ubiquitin-like domain primes mitochondrial translocation of Parkin and regulates mitophagy. *Sci Rep*. 2012;2:1002. doi: 10.1038/srep01002.
74. Kane LA, Lazarou M, Fogel AI, Li Y, Yamano K, Sarraf SA, Banerjee S, Youle RJ. PINK1 phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *J Cell Biol*. 2014;205:143–153. doi: 10.1083/jcb.201402104.
75. Kazlauskaitė A, Kondapalli C, Gourlay R, Campbell DG, Ritorto MS, Hofmann K, Alessi DR, Knebel A, Trost M, Muqit MM. Parkin is activated by PINK1-dependent phosphorylation of ubiquitin at Ser65. *Biochem J*. 2014;460:127–139. doi: 10.1042/BJ20140334.
76. Koyano F, Okatsu K, Kosako H, et al. Ubiquitin is phosphorylated by PINK1 to activate parkin. *Nature*. 2014;510:162–166. doi: 10.1038/nature13392.
77. Ordureau A, Sarraf SA, Duda DM, Heo JM, Jedrychowski MP, Sviderskiy VO, Olszewski JL, Koerber JT, Xie T, Beausoleil SA, Wells JA, Gygi SP, Schulman BA, Harper JW. Quantitative proteomics reveal a feedforward mechanism for mitochondrial PARKIN translocation and ubiquitin chain synthesis. *Mol Cell*. 2014;56:360–375. doi: 10.1016/j.molcel.2014.09.007.
78. Chan NC, Salazar AM, Pham AH, Sweredoski MJ, Kolawa NJ, Graham RL, Hess S, Chan DC. Broad activation of the ubiquitin-proteasome system by Parkin is critical for mitophagy. *Hum Mol Genet*. 2011;20:1726–1737. doi: 10.1093/hmg/ddr048.
79. Okatsu K, Iemura S, Koyano F, Go E, Kimura M, Natsume T, Tanaka K, Matsuda N. Mitochondrial hexokinase HKI is a novel substrate of the Parkin ubiquitin ligase. *Biochem Biophys Res Commun*. 2012;428:197–202. doi: 10.1016/j.bbrc.2012.10.041.
80. Sarraf SA, Raman M, Guarani-Pereira V, Sowa ME, Huttlin EL, Gygi SP, Harper JW. Landscape of the PARKIN-dependent ubiquitylome in response to mitochondrial depolarization. *Nature*. 2013;496:372–376. doi: 10.1038/nature12043.
81. Yang JY, Yang WY. Bit-by-bit autophagic removal of parkin-labelled mitochondria. *Nat Commun*. 2013;4:2428. doi: 10.1038/ncomms3428.
82. Narendra D, Kane LA, Hauser DN, Fearnley IM, Youle RJ. p62/SQSTM1 is required for Parkin-induced mitochondrial clustering but not mitophagy; VDAC1 is dispensable for both. *Autophagy*. 2010;6:1090–1106.
83. Okatsu K, Saisho K, Shimanuki M, Nakada K, Shitara H, Sou YS, Kimura M, Sato S, Hattori N, Komatsu M, Tanaka K, Matsuda N. p62/SQSTM1 cooperates with Parkin for perinuclear clustering of depolarized mitochondria. *Genes Cells*. 2010;15:887–900. doi: 10.1111/j.1365-2443.2010.01426.x.
84. Kim NC, Tresse E, Kolaitis RM, et al. VCP is essential for mitochondrial quality control by PINK1/Parkin and this function is impaired by VCP mutations. *Neuron*. 2013;78:65–80. doi: 10.1016/j.neuron.2013.02.029.
85. Soubannier V, McLelland GL, Zunino R, Braschi E, Rippstein P, Fon EA, McBride HM. A vesicular transport pathway shuttles cargo from mitochondria to lysosomes. *Curr Biol*. 2012;22:135–141. doi: 10.1016/j.cub.2011.11.057.
86. Soubannier V, Rippstein P, Kaufman BA, Shoubridge EA, McBride HM. Reconstitution of mitochondria derived vesicle formation demonstrates selective enrichment of oxidized cargo. *PLoS One*. 2012;7:e52830. doi: 10.1371/journal.pone.0052830.
87. McLelland GL, Soubannier V, Chen CX, McBride HM, Fon EA. Parkin and PINK1 function in a vesicular trafficking pathway regulating mitochondrial quality control. *EMBO J*. 2014;33:282–295. doi: 10.1002/embj.201385902.
88. Bingol B, Tea JS, Phu L, Reichelt M, Bakalarski CE, Song Q, Foreman O, Kirkpatrick DS, Sheng M. The mitochondrial deubiquitinase USP30 opposes parkin-mediated mitophagy. *Nature*. 2014;510:370–375. doi: 10.1038/nature13418.
89. Hoshino A, Mita Y, Okawa Y, Ariyoshi M, Iwai-Kanai E, Ueyama T, Ikeda K, Ogata T, Matoba S. Cytosolic p53 inhibits Parkin-mediated mitophagy and promotes mitochondrial dysfunction in the mouse heart. *Nat Commun*. 2013;4:2308. doi: 10.1038/ncomms3308.
90. Hollville E, Carroll RG, Cullen SP, Martin SJ. Bcl-2 family proteins participate in mitochondrial quality control by regulating Parkin/PINK1-dependent mitophagy. *Mol Cell*. 2014;55:451–466. doi: 10.1016/j.molcel.2014.06.001.
91. Wang X, Winter D, Ashrafi G, Schlehe J, Wong YL, Selkoe D, Rice S, Steen J, LaVoie MJ, Schwarz TL. PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell*. 2011;147:893–906. doi: 10.1016/j.cell.2011.10.018.
92. Shin JH, Ko HS, Kang H, Lee Y, Lee YI, Pletinkova O, Troconso JC, Dawson VL, Dawson TM. PARIS (ZNF746) repression of PGC-1 α contributes to neurodegeneration in Parkinson's disease. *Cell*. 2011;144:689–702. doi: 10.1016/j.cell.2011.02.010.
93. Kim KY, Stevens MV, Akter MH, Rusk SE, Huang RJ, Cohen A, Noguchi A, Springer D, Bocharov AV, Eggerman TL, Suen DF, Youle RJ, Amar M, Remaley AT, Sack MN. Parkin is a lipid-responsive regulator of fat uptake in mice and mutant human cells. *J Clin Invest*. 2011;121:3701–3712. doi: 10.1172/JCI44736.

94. Lu K, Psakhye I, Jentsch S. Autophagic clearance of polyQ proteins mediated by ubiquitin-Atg8 adaptors of the conserved CUET protein family. *Cell*. 2014;158:549–563. doi: 10.1016/j.cell.2014.05.048.
95. Kubli DA, Ycaza JE, Gustafsson AB. Bnip3 mediates mitochondrial dysfunction and cell death through Bax and Bak. *Biochem J*. 2007;405:407–415. doi: 10.1042/BJ20070319.
96. Chen Y, Lewis W, Diwan A, Cheng EH, Matkovich SJ, Dorn GW II. Dual autonomous mitochondrial cell death pathways are activated by Nix/Bnip3L and induce cardiomyopathy. *Proc Natl Acad Sci U S A*. 2010;107:9035–9042. doi: 10.1073/pnas.0914013107.
97. Rikka S, Quinsay MN, Thomas RL, Kubli DA, Zhang X, Murphy AN, Gustafsson AB. Bnip3 impairs mitochondrial bioenergetics and stimulates mitochondrial turnover. *Cell Death Differ*. 2011;18:721–731. doi: 10.1038/cdd.2010.146.
98. Sandoval H, Thiagarajan P, Dasgupta SK, Schumacher A, Prchal JT, Chen M, Wang J. Essential role for Nix in autophagic maturation of erythroid cells. *Nature*. 2008;454:232–235. doi: 10.1038/nature07006.
99. Ding WX, Ni HM, Li M, Liao Y, Chen X, Stolz DB, Dorn GW II, Yin XM. Nix is critical to two distinct phases of mitophagy, reactive oxygen species-mediated autophagy induction and Parkin-ubiquitin-p62-mediated mitochondrial priming. *J Biol Chem*. 2010;285:27879–27890. doi: 10.1074/jbc.M110.119537.
100. Kubli DA, Quinsay MN, Huang C, Lee Y, Gustafsson AB. Bnip3 functions as a mitochondrial sensor of oxidative stress during myocardial ischemia and reperfusion. *Am J Physiol Heart Circ Physiol*. 2008;295:H2025–H2031. doi: 10.1152/ajpheart.00552.2008.
101. Hamacher-Brady A, Brady NR, Logue SE, Sayen MR, Jinno M, Kirshenbaum LA, Gottlieb RA, Gustafsson AB. Response to myocardial ischemia/reperfusion injury involves Bnip3 and autophagy. *Cell Death Differ*. 2007;14:146–157. doi: 10.1038/sj.cdd.4401936.
102. Quinsay MN, Thomas RL, Lee Y, Gustafsson AB. Bnip3-mediated mitochondrial autophagy is independent of the mitochondrial permeability transition pore. *Autophagy*. 2010;6:855–862.
103. Landes T, Emorine LJ, Courilleau D, Rojo M, Belenguer P, Arnauné-Pelloquin L. The BH3-only Bnip3 binds to the dynamin Opa1 to promote mitochondrial fragmentation and apoptosis by distinct mechanisms. *EMBO Rep*. 2010;11:459–465. doi: 10.1038/embor.2010.50.
104. Zhu Y, Massen S, Terenzio M, Lang V, Chen-Lindner S, Eils R, Novak I, Dikic I, Hamacher-Brady A, Brady NR. Modulation of serines 17 and 24 in the LC3-interacting region of Bnip3 determines pro-survival mitophagy versus apoptosis. *J Biol Chem*. 2013;288:1099–1113. doi: 10.1074/jbc.M112.399345.
105. Regula KM, Ens K, Kirshenbaum LA. Inducible expression of BNIP3 provokes mitochondrial defects and hypoxia-mediated cell death of ventricular myocytes. *Circ Res*. 2002;91:226–231.
106. Yussman MG, Toyokawa T, Odley A, Lynch RA, Wu G, Colbert MC, Aronow BJ, Lorenz JN, Dorn GW II. Mitochondrial death protein Nix is induced in cardiac hypertrophy and triggers apoptotic cardiomyopathy. *Nat Med*. 2002;8:725–730. doi: 10.1038/nm719.
107. Diwan A, Krenz M, Syed FM, Wansapura J, Ren X, Koesters AG, Li H, Kirshenbaum LA, Hahn HS, Robbins J, Jones WK, Dorn GW. Inhibition of ischemic cardiomyocyte apoptosis through targeted ablation of Bnip3 restrains postinfarction remodeling in mice. *J Clin Invest*. 2007;117:2825–2833. doi: 10.1172/JCI32490.
108. Dhingra R, Margulets V, Chowdhury SR, Thliveris J, Jassal D, Fernyhough P, Dorn GW II, Kirshenbaum LA. Bnip3 mediates doxorubicin-induced cardiac myocyte necrosis and mortality through changes in mitochondrial signaling. *Proc Natl Acad Sci U S A*. 2014;111:E5537–E5544. doi: 10.1073/pnas.1414665111.
109. Wu W, Tian W, Hu Z, et al. ULK1 translocates to mitochondria and phosphorylates FUNDC1 to regulate mitophagy. *EMBO Rep*. 2014;15:566–575. doi: 10.1002/embr.201438501.
110. Chen G, Han Z, Feng D, et al. A regulatory signaling loop comprising the PGAM5 phosphatase and CK2 controls receptor-mediated mitophagy. *Mol Cell*. 2014;54:362–377. doi: 10.1016/j.molcel.2014.02.034.
111. Liu X, Ye B, Miller S, Yuan H, Zhang H, Tian L, Nie J, Imae R, Arai H, Li Y, Cheng Z, Shi Y. Ablation of ALCAT1 mitigates hypertrophic cardiomyopathy through effects on oxidative stress and mitophagy. *Mol Cell Biol*. 2012;32:4493–4504. doi: 10.1128/MCB.01092-12.
112. Webster BR, Scott I, Traba J, Han K, Sack MN. Regulation of autophagy and mitophagy by nutrient availability and acetylation. *Biochim Biophys Acta*. 2014;1841:525–534. doi: 10.1016/j.bbali.2014.02.001.
113. Mariño G, Pietrocola F, Eisenberg T, et al. Regulation of autophagy by cytosolic acetyl-coenzyme A. *Mol Cell*. 2014;53:710–725. doi: 10.1016/j.molcel.2014.01.016.
114. Webster BR, Scott I, Han K, Li JH, Lu Z, Stevens MV, Malide D, Chen Y, Samsel L, Connelly PS, Daniels MP, McCoy JP Jr, Combs CA, Gucek M, Sack MN. Restricted mitochondrial protein acetylation initiates mitochondrial autophagy. *J Cell Sci*. 2013;126:4843–4849. doi: 10.1242/jcs.131300.
115. Scott I, Webster BR, Chan CK, Okonkwo JU, Han K, Sack MN. GCN5-like protein 1 (GCN5L1) controls mitochondrial content through coordinated regulation of mitochondrial biogenesis and mitophagy. *J Biol Chem*. 2014;289:2864–2872. doi: 10.1074/jbc.M113.521641.
116. Scorrano L. Keeping mitochondria in shape: a matter of life and death. *Eur J Clin Invest*. 2013;43:886–893. doi: 10.1111/eci.12135.
117. Ikeda Y, Shirakabe A, Brady C, Zablocki D, Ohishi M, Sadoshima J. Molecular mechanisms mediating mitochondrial dynamics and mitophagy and their functional roles in the cardiovascular system. *J Mol Cell Cardiol*. 2015;78:116–122. doi: 10.1016/j.yjmcc.2014.09.019.
118. Kanki T, Wang K, Baba M, Bartholomew CR, Lynch-Day MA, Du Z, Geng J, Mao K, Yang Z, Yen WL, Klionsky DJ. A genomic screen for yeast mutants defective in selective mitochondria autophagy. *Mol Biol Cell*. 2009;20:4730–4738. doi: 10.1091/mbc.E09-03-0225.
119. Rambold AS, Kostelecky B, Elia N, Lippincott-Schwartz J. Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc Natl Acad Sci U S A*. 2011;108:10190–10195. doi: 10.1073/pnas.1107402108.
120. Deng H, Dodson MW, Huang H, Guo M. The Parkinson's disease genes pink1 and parkin promote mitochondrial fission and/or inhibit fusion in *Drosophila*. *Proc Natl Acad Sci U S A*. 2008;105:14503–14508. doi: 10.1073/pnas.0803998105.
121. Frank M, Duvezin-Caubet S, Koob S, Occhipinti A, Jagasia R, Petcherski A, Ruonala MO, Priault M, Salin B, Reichert AS. Mitophagy is triggered by mild oxidative stress in a mitochondrial fission dependent manner. *Biochim Biophys Acta*. 2012;1823:2297–2310. doi: 10.1016/j.bbamer.2012.08.007.
122. Gomes AV, Zong C, Ping P. Protein degradation by the 26S proteasome system in the normal and stressed myocardium. *Antioxid Redox Signal*. 2006;8:1677–1691. doi: 10.1089/ars.2006.8.1677.
123. Zhao T, Huang X, Han L, Wang X, Cheng H, Zhao Y, Chen Q, Chen J, Cheng H, Xiao R, Zheng M. Central role of mitofusin 2 in autophagosome-lysosome fusion in cardiomyocytes. *J Biol Chem*. 2012;287:23615–23625. doi: 10.1074/jbc.M112.379164.
124. Ikeda Y, Shirakabe A, Maejima Y, Zhai P, Sciarretta S, Toli J, Nomura M, Mihara K, Egashira K, Ohishi M, Abellatif M, Sadoshima J. Endogenous drp1 mediates mitochondrial autophagy and protects the heart against energy stress. *Circ Res*. 2015;116:264–278. doi: 10.1161/CIRCRESAHA.116.303356.
125. Lee Y, Lee HY, Hanna RA, Gustafsson AB. Mitochondrial autophagy by Bnip3 involves Drp1-mediated mitochondrial fission and recruitment of Parkin in cardiac myocytes. *Am J Physiol Heart Circ Physiol*. 2011;301:H1924–H1931. doi: 10.1152/ajpheart.00368.2011.
126. Ashrafian H, Docherty L, Leo V, et al. A mutation in the mitochondrial fission gene Dnm1l leads to cardiomyopathy. *PLoS Genet*. 2010;6:e1001000. doi: 10.1371/journal.pgen.1001000.
127. Song M, Mihara K, Chen Y, Scorrano L, Dorn GWn. Mitochondria fusion and fission factors reciprocally orchestrate mitophagic culling in mouse hearts and cultured fibroblasts. *Cell Metab*. 2015;21:273–285. doi: 10.1016/j.cmet.2014.12.011.
128. Chen L, Liu T, Tran A, Lu X, Tomilov AA, Davies V, Cortopassi G, Chiamvimonvat N, Bers DM, Votruba M, Knowlton AA. OPA1 mutation and late-onset cardiomyopathy: mitochondrial dysfunction and mtDNA instability. *J Am Heart Assoc*. 2012;1:e003012. doi: 10.1161/JAHA.112.003012.
129. Papanicolaou KN, Ngho GA, Dabkowski ER, O'Connell KA, Ribeiro RF Jr, Stanley WC, Walsh K. Cardiomyocyte deletion of mitofusin-1 leads to mitochondrial fragmentation and improves tolerance to ROS-induced mitochondrial dysfunction and cell death. *Am J Physiol Heart Circ Physiol*. 2012;302:H167–H179. doi: 10.1152/ajpheart.00833.2011.
130. Papanicolaou KN, Khairallah RJ, Ngho GA, Chikando A, Luptak I, O'Shea KM, Riley DD, Lugus JJ, Colucci WS, Lederer WJ, Stanley WC, Walsh K. Mitofusin-2 maintains mitochondrial structure and contributes to stress-induced permeability transition in cardiac myocytes. *Mol Cell Biol*. 2011;31:1309–1328. doi: 10.1128/MCB.00911-10.
131. Chen Y, Liu Y, Dorn GW II. Mitochondrial fusion is essential for organelle function and cardiac homeostasis. *Circ Res*. 2011;109:1327–1331. doi: 10.1161/CIRCRESAHA.111.258723.
132. Papanicolaou KN, Kikuchi R, Ngho GA, Coughlan KA, Dominguez I, Stanley WC, Walsh K. Mitofusins 1 and 2 are essential for postnatal

- metabolic remodeling in heart. *Circ Res*. 2012;111:1012–1026. doi: 10.1161/CIRCRESAHA.112.274142.
133. Song M, Chen Y, Gong G, Murphy E, Rabinovitch PS, Dorn GW II. Super-suppression of mitochondrial reactive oxygen species signaling impairs compensatory autophagy in primary mitophagic cardiomyopathy. *Circ Res*. 2014;115:348–353. doi: 10.1161/CIRCRESAHA.115.304384.
 134. Bhandari P, Song M, Chen Y, Burelle Y, Dorn GW II. Mitochondrial contagion induced by Parkin deficiency in *Drosophila* hearts and its containment by suppressing mitofusin. *Circ Res*. 2014;114:257–265. doi: 10.1161/CIRCRESAHA.114.302734.
 135. Nakai A, Yamaguchi O, Takeda T, Higuchi Y, Hikoso S, Taniike M, Omiya S, Mizote I, Matsumura Y, Asahi M, Nishida K, Hori M, Mizushima N, Otsu K. The role of autophagy in cardiomyocytes in the basal state and in response to hemodynamic stress. *Nat Med*. 2007;13:619–624. doi: 10.1038/nm1574.
 136. Matsui Y, Takagi H, Qu X, Abdellatif M, Sakoda H, Asano T, Levine B, Sadoshima J. Distinct roles of autophagy in the heart during ischemia and reperfusion: roles of AMP-activated protein kinase and Beclin 1 in mediating autophagy. *Circ Res*. 2007;100:914–922. doi: 10.1161/01.RES.0000261924.76669.36.
 137. Kubli DA, Zhang X, Lee Y, Hanna RA, Quinsay MN, Nguyen CK, Jimenez R, Petrosyan S, Murphy AN, Gustafsson AB. Parkin protein deficiency exacerbates cardiac injury and reduces survival following myocardial infarction. *J Biol Chem*. 2013;288:915–926. doi: 10.1074/jbc.M112.411363.
 138. Piquereau J, Godin R, Deschênes S, Bessi VL, Mofarrahi M, Hussain SN, Burelle Y. Protective role of PARK2/Parkin in sepsis-induced cardiac contractile and mitochondrial dysfunction. *Autophagy*. 2013;9:1837–1851. doi: 10.4161/auto.26502.
 139. Yang KC, Ma X, Liu H, Murphy J, Barger PM, Mann DL, Diwan A. Tumor necrosis factor receptor-associated factor 2 mediates mitochondrial autophagy. *Circ Heart Fail*. 2015;8:175–187. doi: 10.1161/CIRCHEARTFAILURE.114.001635.
 140. Billia F, Hauck L, Konecny F, Rao V, Shen J, Mak TW. PTEN-inducible kinase 1 (PINK1)/Park6 is indispensable for normal heart function. *Proc Natl Acad Sci U S A*. 2011;108:9572–9577. doi: 10.1073/pnas.1106291108.
 141. Pridgeon JW, Olzmann JA, Chin LS, Li L. PINK1 protects against oxidative stress by phosphorylating mitochondrial chaperone TRAP1. *PLoS Biol*. 2007;5:e172. doi: 10.1371/journal.pbio.0050172.
 142. Kanamori H, Takemura G, Goto K, Maruyama R, Tsujimoto A, Ogino A, Takeyama T, Kawaguchi T, Watanabe T, Fujiwara T, Fujiwara H, Seishima M, Minatoguchi S. The role of autophagy emerging in postinfarction cardiac remodeling. *Cardiovasc Res*. 2011;91:330–339. doi: 10.1093/cvr/cvr073.
 143. Lesnefsky EJ, Moghaddas S, Tandler B, Kerner J, Hoppel CL. Mitochondrial dysfunction in cardiac disease: ischemia–reperfusion, aging, and heart failure. *J Mol Cell Cardiol*. 2001;33:1065–1089. doi: 10.1006/jmcc.2001.1378.
 144. Siddall HK, Yellon DM, Ong SB, Mukherjee UA, Burke N, Hall AR, Angelova PR, Ludtmann MH, Deas E, Davidson SM, Mocanu MM, Hausenloy DJ. Loss of PINK1 increases the heart's vulnerability to ischemia-reperfusion injury. *PLoS One*. 2013;8:e62400. doi: 10.1371/journal.pone.0062400.
 145. Huang C, Andres AM, Ratliff EP, Hernandez G, Lee P, Gottlieb RA. Preconditioning involves selective mitophagy mediated by Parkin and p62/SQSTM1. *PLoS One*. 2011;6:e20975. doi: 10.1371/journal.pone.0020975.
 146. Orvedahl A, Sumpster R Jr, Xiao G, et al. Image-based genome-wide siRNA screen identifies selective autophagy factors. *Nature*. 2011;480:113–117. doi: 10.1038/nature10546.
 147. Xu X, Kobayashi S, Chen K, Timm D, Volden P, Huang Y, Gulick J, Yue Z, Robbins J, Epstein PN, Liang Q. Diminished autophagy limits cardiac injury in mouse models of type 1 diabetes. *J Biol Chem*. 2013;288:18077–18092. doi: 10.1074/jbc.M113.474650.
 148. Elgendy M, Sheridan C, Brumatti G, Martin SJ. Oncogenic Ras-induced expression of Noxa and Beclin-1 promotes autophagic cell death and limits clonogenic survival. *Mol Cell*. 2011;42:23–35. doi: 10.1016/j.molcel.2011.02.009.
 149. Liu Y, Shoji-Kawata S, Sumpster RM Jr, Wei Y, Ginet V, Zhang L, Posner B, Tran KA, Green DR, Xavier RJ, Shaw SY, Clarke PG, Puyal J, Levine B. Autosis is a Na⁺,K⁺-ATPase-regulated form of cell death triggered by autophagy-inducing peptides, starvation, and hypoxia-ischemia. *Proc Natl Acad Sci U S A*. 2013;110:20364–20371. doi: 10.1073/pnas.1319661110.

Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Molecular Mechanisms of Mitochondrial Autophagy/Mitophagy in the Heart Toshiro Saito and Junichi Sadoshima

Circ Res. 2015;116:1477-1490

doi: 10.1161/CIRCRESAHA.116.303790

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2015 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circres.ahajournals.org/content/116/8/1477>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>